


MONITORING *the* FUTURE

NATIONAL SURVEY RESULTS
ON DRUG USE
1975–2016

2016 Overview

Key Findings on Adolescent Drug Use

Lloyd D. Johnston
Patrick M. O'Malley
Richard A. Miech
Jerald G. Bachman
John E. Schulenberg



MONITORING THE FUTURE

NATIONAL SURVEY RESULTS ON DRUG USE

2016 Overview Key Findings on Adolescent Drug Use

by

Lloyd D. Johnston, Ph.D.
Patrick M. O'Malley, Ph.D.
Richard A. Miech, Ph.D.
Jerald G. Bachman, Ph.D.
John E. Schulenberg, Ph.D.

The University of Michigan
Institute for Social Research

Sponsored by:
The National Institute on Drug Abuse
National Institutes of Health

This publication was written by the principal investigators and staff of the Monitoring the Future project at the Institute for Social Research, the University of Michigan, under Research Grant R01 DA 001411 from the National Institute on Drug Abuse.

The findings and conclusions in this report are those of the authors and do not necessarily represent the views of the sponsor.

Public Domain Notice

All material appearing in this volume is in the public domain and may be reproduced or copied, whether in print or non-print media including derivatives, without permission from the authors. If you plan to modify the material, please contact the Monitoring the Future Project at mtfinformation@umich.edu for verification of accuracy. Citation of the source is appreciated, including at least the following: Monitoring the Future, Institute for Social Research, The University of Michigan.



<http://creativecommons.org/licenses/by/4.0/>

Recommended Citation

Johnston, L. D., O'Malley, P. M., Miech, R. A., Bachman, J. G., & Schulenberg, J. E. (2017). *Monitoring the Future national survey results on drug use, 1975-2016: Overview, key findings on adolescent drug use*. Ann Arbor: Institute for Social Research, The University of Michigan.

Institute for Social Research
The University of Michigan
Ann Arbor, Michigan
Printed January 2017

Contents

Click on any item below (in [blue](#)) to go directly to that page.

Introduction.....	1
Study Design and Methods	3
Summary of Key Findings	5
Specific Results by Drug	
Any Illicit Drug.....	9
Marijuana	11
Synthetic Marijuana	13
Inhalants.....	15
LSD	17
Cocaine	19
Crack.....	21
Amphetamines and Other Stimulant Drugs	23
Methamphetamine and Crystal Methamphetamine (Ice)	25
Heroin	27
Other Narcotic Drugs, Including OxyContin and Vicodin	29
Tranquilizers	31
Sedatives (Barbiturates)	33
MDMA (Ecstasy, Molly) and Other “Club Drugs”	35
Alcohol.....	37
Cigarettes	39
Smokeless Tobacco.....	41
Vaping.....	43
Small Cigars, Cigarillos, Large Cigars, and Tobacco Using a Hookah.....	45
Steroids	47
Subgroup Differences	49
Lessons Learned.....	52
Tables	
1. Trends in Lifetime Prevalence of Use of Various Drugs for Grades 8, 10, and 12 Combined.....	54
2. Trends in Annual Prevalence of Use of Various Drugs for Grades 8, 10, and 12 Combined.....	56
3. Trends in 30-Day Prevalence of Use of Various Drugs for Grades 8, 10, and 12 Combined.....	58

4.	Trends in Daily Prevalence of Use of Selected Drugs for Grades 8, 10, and 12 Combined.....	60
5.	Trends in Lifetime Prevalence of Use of Various Drugs in Grades 8, 10, and 12	62
6.	Trends in Annual Prevalence of Use of Various Drugs in Grades 8, 10, and 12	68
7.	Trends in 30-day Prevalence of Use of Various Drugs in Grades 8, 10, and 12	76
8.	Trends in 30-Day Prevalence of Daily Use of Various Drugs in Grades 8, 10, and 12	82
9.	Trends in Harmfulness of Drugs as Perceived by 8th Graders.....	87
10.	Trends in Harmfulness of Drugs as Perceived by 10th Graders.....	90
11.	Trends in Harmfulness of Drugs as Perceived by 12th Graders.....	93
12.	Trends in Disapproval of Drug Use in Grade 8.....	97
13.	Trends in Disapproval of Drug Use in Grade 10.....	100
14.	Trends in Disapproval of Drug Use in Grade 12.....	103
15.	Trends in Availability of Drugs as Perceived by 8th Graders	107
16.	Trends in Availability of Drugs as Perceived by 10th Graders	109
17.	Trends in Availability of Drugs as Perceived by 12th Graders	111

Introduction

Monitoring the Future (MTF) is a long-term study of American adolescents, college students, and adult high school graduates through age 55. It has been conducted annually by the University of Michigan's Institute for Social Research since its inception in 1975 and is supported under a series of investigator-initiated, competitive research grants from the National Institute on Drug Abuse.

The need for a study such as MTF is clear. Substance use by American young people has proven to be a rapidly changing phenomenon, requiring frequent assessments and reassessments. Since the mid-1960s, when it burgeoned in the general youth population, illicit drug use has remained a major concern for the nation. Smoking, drinking, and illicit drug use are leading causes of morbidity and mortality during adolescence as well as later in life. How vigorously the nation responds to teenage substance use, how accurately it identifies the emerging substance abuse problems, and how well it comes to understand the effectiveness of policy and intervention efforts largely depend on the ongoing collection of valid and reliable data. MTF is uniquely designed to generate such data in order to provide an accurate picture of what is happening in this domain and why, and the study has served that function well for the past 42 years. Policy discussions in the scientific literature and media, in government, education, public health institutions, and elsewhere have been informed by the ready availability of extensive and consistently accurate information from the study relating to a large and ever-growing number of substances. Similarly, the work of organizations and agencies providing prevention and treatment services is informed by MTF.

The 2016 MTF survey involved about 45,500 students in 8th-, 10th-, and 12th grades enrolled in 372 secondary schools nationwide. The first published results based on the 2016 survey are presented in this report. Recent trends in the use of licit and illicit drugs are emphasized, as well as trends in the levels of perceived risk and personal disapproval associated with each drug. This project has shown these beliefs and attitudes to be particularly important in explaining trends in use.

In addition, trends in the perceived availability of each drug are presented, which at times have proven important to explaining changes in usage levels for some drugs.

A synopsis of the design and methods used in the study and an overview of the key results from the 2016 survey follow this introductory section. We then provide a separate section for each individual drug class, including figures that show trends in the overall proportions of students at each grade level (a) using the drug, (b) seeing a “great risk” associated with its use (perceived risk), (c) disapproving of its use (disapproval), and (d) saying that it would be “fairly easy” or “very easy” to get if they wanted to (perceived availability). For 12th graders, annual data are available since 1975 and for 8th and 10th graders since 1991, the first year they were included in the study.

The tables at the end of this report provide the statistics underlying the figures; in addition, they present data on lifetime, annual, 30-day, and (for selected drugs) daily prevalence.¹ For the sake of brevity, we present these prevalence statistics here in tabular form only for the 1991–2016 interval, but statistics on 12th graders going back to 1975 are available in other MTF publications. For each prevalence period, the tables indicate which one-year changes from 2015 to 2016 are statistically significant. (In the text below, ‘s’ indicates $p \leq .05$, ‘ss’ indicates $p \leq .01$, ‘sss’ indicates $p \leq .001$, and ‘ns’ indicates not statistically significant). The graphic depictions of multiyear trends often reveal gradual change that may not reach significance in a given one-year interval but nevertheless may be shown to be real over a longer time frame.

An extensive analysis of the study's findings on secondary school students may be found in *Volume I*, the second publication in this series, published at the end of May each year.² *Volume I* contains a more detailed description of the study's methodology, as well as chapters on grade of initiation, attitudes toward drugs, the social milieu, and a summary of other publications from the study that year (mostly journal articles). *Volume I* also contains an appendix

¹Prevalence refers to the proportion or percentage of the sample reporting use of the given substance on one or more occasions in a given time interval—e.g., lifetime, past 12 months, or past 30 days. For most drugs, the prevalence of daily use refers to reported use on 20 or more occasions in the past 30 days, except for cigarettes and smokeless tobacco, for which actual daily use is measured, and for binge drinking, defined as having 5+ drinks on at least one occasion in the prior two weeks. E-cigarettes and some tobacco products are measured on number of days used in past 30 days.

²The most recent publication of *Volume I* is Miech, R. A., Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (2016). *Monitoring the Future national survey results on drug use, 1975–2015: Volume I, Secondary school students*. Ann Arbor: Institute for Social Research, The University of Michigan, 636 pp. Available at: http://monitoringthefuture.org/pubs/monographs/mtf-vol1_2015.pdf

report because the follow-up data from those populations become available later in the year. Those findings will be covered in *Volume II*, the third monograph in this annual series, published at the end of July each year.³

Two annual MTF Occasional Papers are published each year in conjunction with *Volumes I* and *II*, providing trends in use for various demographic subgroups.⁴

A fourth monograph, *HIV/AIDS Risk and Protective Behaviors Among Young Adults*, dealing with national trends in HIV/AIDS-related risk and protective behaviors among young adults 21 to 40 years old, was added to the series beginning in 2010.⁵ It is published in October of

each year. From 2005 to 2009, these findings were reported as part of *Volume II*.

For the publication years prior to 2010, the volumes in these annual series are available from the NIDA Drug Publications Research Dissemination Center (877-NIDA-NIH, drugpubs.drugabuse.gov) and can also be found on the MTF website. Beginning with the 2010 publication date, the volumes are available at the MTF website immediately upon publication. Further information on the study, including its latest press releases, a listing of all publications, and freely accessible reports may also be found at www.monitoringthefuture.org.

³The most recent publication of *Volume II* is Johnston, L. D., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., and Miech, R. A. (2016). *Monitoring the Future national survey results on drug use, 1975–2015: Volume II, College students & adults ages 19–55*. Ann Arbor: Institute for Social Research, The University of Michigan, 427 pp. Available at: http://www.monitoringthefuture.org/pubs/monographs/mtf-vol2_2015.pdf

⁴Johnston, L. D., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., & Miech, R. A. (2016). *Demographic subgroup trends among adolescents in the use of various licit and illicit drugs 1975–2015* (Monitoring the Future Occasional Paper No. 86). Ann Arbor, MI: Institute for Social Research, University of Michigan, 552 pp. Available at: <http://monitoringthefuture.org/pubs/occapers/mtf-occ86.pdf>

Johnston, L. D., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., & Miech, R. A. (2016). *Demographic subgroup trends among young adults in the use of various licit and illicit drugs 1989–2015* (Monitoring the Future Occasional Paper No. 87). Ann Arbor, MI: Institute for Social Research, University of Michigan, 109 pp. Available at: <http://monitoringthefuture.org/pubs/occapers/mtf-occ87.pdf>

⁵The most recent publication in the *HIV/AIDS monograph series* is Johnston, L. D., O'Malley, P. M., Bachman, J. G., Schulenberg, J. E., Patrick, M. E., & Miech, R. A. (2016). *HIV/AIDS: Risk and protective behaviors among adults ages 21–40 in the U.S., 2004–2015*. Ann Arbor: Institute for Social Research, The University of Michigan, 123 pp. Available at: http://monitoringthefuture.org/pubs/monographs/mtf-hiv-aids_2015.pdf

Study Design and Methods

Monitoring the Future's main data collection involves a series of large, annual surveys of nationally representative samples of public and private secondary school students throughout the coterminous United States. Every year since 1975, such samples of 12th graders have been surveyed. In 1991, the study was expanded to include comparable, independent national samples of 8th and 10th graders. The year 2016 marked the 42nd survey of 12th graders and the 26th survey of 8th and 10th graders.

Sample Sizes

In 2016 about 45,500 students in 372 secondary schools participated in the study, with sample sizes in 8th, 10th, and 12th grades of about 17,600, 15,200, and 12,600, respectively. The number of cases upon which a particular statistic is based may be less than the total sample size. Multiple questionnaire forms are distributed randomly at each grade level to increase coverage of attitudinal and behavioral domains relevant to substance use. To reduce burden on the respondents, not all questions are contained in all forms. The tables here contain notes on the number of forms used for each statistic if less than the total sample is used.

Field Procedures

University of Michigan staff members administer the questionnaires to students, usually in the student classroom during a regular class period. Participation is voluntary. Parents are notified well in advance of the survey administration and are provided the opportunity to decline their child's participation. Questionnaires are self-completed and are formatted for optical scanning. Procedures are kept consistent over time.

In 8th and 10th grades the questionnaires are completely anonymous, and in 12th grade they are confidential (name and address information is gathered separately from the 12th grade questionnaire to permit the longitudinal follow-up surveys of random subsamples of participants after high school). Extensive procedures are followed to protect the confidentiality of the participants and their data. All procedures are reviewed and approved on an annual basis by the University of Michigan's Institutional Review Board (IRB) for compliance with federal guidelines for the treatment of human subjects.

Measures

A standard set of three questions is used to determine usage levels for most of the drugs. For example, we ask, "On how many occasions (if any) have you used marijuana... (a)...in your lifetime? (b)...during the last 12 months? (c)...during the last 30 days?" Each of the

three questions is answered on the same answer scale: 0, 1–2, 3–5, 6–9, 10–19, 20–39, and 40 or more occasions.

For the psychotherapeutic drugs (amphetamines, sedatives [barbiturates], tranquilizers, and narcotics other than heroin), respondents are instructed to include only use "...on your own—that is, without a doctor telling you to take them." A similar qualification is used in the question on use of anabolic steroids, OxyContin, Vicodin, and several other drugs.

For cigarettes, respondents are asked two questions about use. First, they are asked, "Have you ever smoked cigarettes?" The answer categories are "never," "once or twice," "occasionally but not regularly," "regularly in the past," and "regularly now." The second question asks, "How frequently have you smoked cigarettes during the past 30 days?" The answer categories are "not at all," "less than one cigarette per day," "one to five cigarettes per day," and about one-half, one, one and one half, and two packs or more per day.

Smokeless tobacco questions parallel those for cigarettes. There are also questions recently added about electronic vaporizers, e-cigarettes, small cigars, and a number of other tobacco products. In general, their use is asked on a prevalence/frequency scale for either the last 12 months or the last 30 days.

Alcohol use is measured using the three questions illustrated above for marijuana. A parallel set of three questions asks about the frequency of being drunk. Binge drinking is assessed with the question, "How many times (if any) have you had five or more drinks in a row" over the past two weeks? Extreme binge, now also called high-intensity, drinking among 12th graders is assessed with similar questions about consuming ten or more and fifteen or more drinks in a row.

In general, we try to keep measures consistent across time. When a change is warranted, we usually splice the older and newer measures for at least one year to permit an assessment of the effect of the change.

Perceived risk is measured by the question, "How much do you think people risk harming themselves (physically or in other ways), if they..." try or use a drug—for example, "...try marijuana once or twice." The answer categories are "no risk," "slight risk," "moderate risk," "great risk," and "can't say, drug unfamiliar." Parallel questions refer to using marijuana "occasionally" and "regularly."

Disapproval is measured by the question “Do YOU disapprove of people doing each of the following?” followed by “trying marijuana once or twice,” for example. Answer categories are “don’t disapprove,” “disapprove,” and “strongly disapprove.” In the 8th- and 10th-grade questionnaires, a fourth category—“can’t say, drug unfamiliar”—is provided and included in the calculation of percentages.

Perceived availability is measured by the question “How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?” Answer categories are “probably impossible,” “very difficult,” “fairly difficult,” “fairly easy,” and “very easy.” For 8th and 10th graders, an additional answer category—“can’t say, drug unfamiliar”—is provided and included in the calculation of percentages.

Summary of Key Findings

Before delving into the 2016 findings, a note about three kinds of trends we reference: Monitoring the Future (MTF) is designed to detect age effects, secular trends (sometimes referred to as period effects), and cohort effects in substance use and related attitudes and beliefs. Age effects (similar changes at similar ages seen across multiple class cohorts) are common during adolescence, and we typically find that use, as well as positive attitudes and beliefs about use, increase across 8th, 10th, and 12th grades. When changes over time in substance use and related attitudes and beliefs are parallel across the three grades, they reflect secular trends, which are also common.

Cohort effects pertain to differences in substance use and related attitudes and behaviors among those born at different times that are maintained as the birth cohorts age (or in this case, as class-in-school cohorts, which are strongly correlated with age). Such cohort effects sometimes drive changes in substance use prevalence at the population level. For example, much of the decline in the prevalence of U.S. cigarette smoking has its roots in youth cohorts that did not take up smoking and then continued to abstain from smoking as they aged into adulthood. As subsequent youth cohorts continued to avoid smoking and then grew older, these cohorts contributed to a decline in the overall population prevalence of smoking. Cohort effects can also act in the opposite direction, with newer cohorts taking up a substance and continuing to use it as they get older. One important contribution of the MTF study has been the specification of cohort effects that emerged starting in the early 1990s, when an increase in youth substance use occurred for many drugs.

MTF allows detection of cohort effects at an early age through comparison of substance use prevalence of 8th, 10th, and 12th graders relative to each other. Often 8th grade substance use is a bellwether, and year-to-year changes that are unique to 8th grade can signify an emerging increase or decrease in substance use at later grade levels with some time lag.

The analyses and associated tables that follow present substance use trends for all three grades separately, as well as trends in key attitudes, beliefs, and perceived availability. In a number of cases we provide insight into the age and cohort effects and secular trends that underlie trends in use and in key attitudes and beliefs.

An additional set of tables provides an overview of drug

use trends for the three grades combined (Tables 1–4). This information gives a summary of the general nature of secular trends over the last several years, though it obscures any cohort effects that may be occurring. Also, for simultaneous trends that are in the same direction and magnitude across all three grades, these combined analyses provide greater statistical power to detect secular trends that are statistically significant.

Declines in Use of a Number of Drugs in 2016

Perhaps the most striking finding in 2016 is that across the broad spectrum of drugs (more than 50 classes and subclasses) most decreased and hardly any exhibited a statistically significant increase.⁶

In many cases these decreases in use were continuing the declines seen in 2014 and 2015. **Cigarettes** and **alcohol** continued to show significant declines, reaching their lowest levels in the history of the study. With regard to illicit drugs, annual prevalence declined in the three grades combined for the use of **any illicit drug, any illicit drug other than marijuana, synthetic marijuana, MDMA** (ecstasy, Molly), **cocaine, crack, amphetamines, Adderall** specifically, **methamphetamine, heroin, Rohypnol**, some psychotherapeutic drugs used without medical supervision (**Vicodin, tranquilizers**), and **steroids**.

The decline in annual prevalence of using **any illicit drug** is largely due to the fact that the annual prevalence of **marijuana**, which tends to drive the overall index, showed a decline in the two lower grades in 2016. (The 2.4 percentage point decline to 9.4% in 8th grade was highly significant; the 1.5 percentage point decline in 10th grade to 23.9% was not.). Among 12th graders, however, the annual prevalence of marijuana has held quite steady for several years, increasing by a non-significant 0.7% to 35.6% in 2016. For the three grades combined, annual prevalence of marijuana showed a significant decline of 1.1 percentage points in 2016.

Although use of marijuana declined (or in the case of the 12th graders held steady) in 2016 marijuana attitudes among students continued to move toward greater acceptance. This fact is contradictory to the association that has existed between perceived risk and actual use for many years, in which they usually move in opposite directions with a one-year lag between a change in perceived risk and a change in annual prevalence. (This is a topic we will address in future publications.) *Perceived risk* of smoking marijuana regularly declined only slightly

⁶ The few exceptions were annual use of bath salts among 8th graders only, annual use of over-the-counter cough and cold medicines to get high among 8th graders,

and 30-day use of heroin with a needle among 10th graders; and none of those changes were large or part of a pattern of continuing increase.

further in 2016 in two of the grades, following a sharp decline in all three grades in recent years. In all three grades, the percentage seeing great risk of smoking marijuana regularly is at or near the lowest point ever recorded in the study—58%, 44%, and 31% in grades 8, 10, and 12, respectively. *Disapproval* of smoking marijuana regularly was unchanged in 8th grade (at 82%) and 10th grade (at 74%) in 2016, but declined non-significantly in 12th grade (to 69%). Although the 2016 levels are at or near historic lows, they still indicate relatively high levels of disapproval of smoking marijuana regularly. Indeed the decline in disapproval or regular marijuana use since about 2008 has been considerably less than the decline in perceived risk. Reported *availability* of marijuana continued a longer-term decline among 8th and 10th graders in 2016.

Use of *any illicit drug other than marijuana* declined slightly in all three grades, significantly so in 8th grade. Annual prevalence in 2016 is 5%, 10%, and 14% in 8th, 10th, and 12th grades, respectively—the lowest levels since all three grades were included in the study in 1991.

Additional drugs with declining annual prevalence include *synthetic marijuana* (which in 2011 was the second most widely used drug after marijuana), *MDMA* (ecstasy, Molly), *inhalants*, *heroin*, *cocaine*, *crack*, *amphetamines*, and *steroids*. *Methamphetamine* use declined significantly at 10th grade but remained unchanged in the other two grades.

The *psychotherapeutic drugs* warrant special attention, given that they now make up a significantly larger part of the overall U.S. drug problem than was true 10–15 years ago. This is in part because of increases in nonmedical use of many prescription drugs over that period, and in part because use of a number of street drugs has declined substantially since the mid- to late-1990s. It seems likely that young people are less concerned about the dangers of using these prescription drugs outside of medical regimen because they are widely used for legitimate purposes. (Indeed, the low levels of perceived risk for sedatives and amphetamines observed among 12th graders illustrate this point.) Also, prescription psychotherapeutic drugs are now being advertised directly to the consumer, which implies that they are both widely used and safe.

Fortunately, the use of most of these drugs has either leveled or begun to decline in the past few years. The proportion of 12th graders misusing any of these prescription drugs (i.e., amphetamines, sedatives, tranquilizers, or narcotics other than heroin) in the prior year continued to decline in 2016 (-1.0%, not significant) to 12%, down from a high of 17% in 2005. *Amphetamine* use without a doctor's orders—which generally has been the second most widely used class of illicit drugs after

marijuana—continued a gradual decline in 2016 in all grades, though the one-year declines did not reach statistical significance. Use of *narcotics* other than heroin without a doctor's orders (reported only for 12th grade) also continued a gradual decline begun after 2009, when annual prevalence was 9.2%; it was 4.8% after a non-significant decline of 0.6 percentage points in 2016. Given the epidemic of narcotics use in older populations along with concurrent rise in medical emergencies and deaths, it is particularly good news that young people are moving away from the use of these drugs.

Illicit Drugs Holding Steady in 2016

The use of a number of drugs showed little or no change from 2015 to 2016. These include *hallucinogens*, *LSD* specifically, *hallucinogens other than LSD*, *salvia*, *tranquilizers*, *heroin* use without a needle, *crystal methamphetamine*, and the club drugs *GHB*, *rohypnol*, and *ketamine*.

Illicit Drugs Showing Any Increase in 2016

The use of so-called “*bath salts*” (synthetic stimulant cathinones) rose significantly (but just by 0.5% to 0.9%) among 8th graders in 2016 but remained unchanged in the upper grades. Similarly, *cough and cold medicines used to get high* (which usually contain dextromethorphan) rose significantly among 8th graders to an annual prevalence of 2.6%, but not among the upper grades.

Tobacco and Alcohol Use

As in 2015, *cigarette smoking* and *alcohol* use have continued their long declines and are now at the lowest levels in the history of the survey. *Thirty-day* prevalence of cigarette use reached a peak in 1996 at grades 8 and 10, capping a rapid climb from the 1991 levels (when data were first gathered on these grades). Between 1996 and 2016, current smoking fell dramatically in these grades, by 87% and 84%, respectively. For 12th graders, peak 30-day prevalence occurred in 1997 at 37% and has shown a more modest decline since then. It is at 11% in 2016—a seven-tenths decline since the peak—with a continuing decline in smoking prevalence in 2016. A similar decline is statistically significant at 8th and 10th grades. Because of the strong cohort effect that we have consistently observed for cigarette smoking, we have predicted use at 12th grade to continue to show declines, as the lighter-using cohorts of 8th and 10th graders become 12th graders.

Initiation of *cigarette* use also continues its long-term and extremely important decline. *Lifetime* prevalence declined between 2015 and 2016 in all three grades: to 10% in 8th grade (-3.5, sss), to 18% in 10th grade (-2.4, s), and to 29% in 12th grade (-2.8, s). The fact that fewer young people now initiate cigarette smoking is an important reason for the large declines in their current use. The proportion of students who have *ever tried* cigarettes

has fallen from peak levels reached in 1996 or 1997 by roughly three quarters, two thirds, and one half in the three grades, respectively.

Overall increases in perceived risk and disapproval appear to have contributed to the downturn in cigarette use. Perceived risk of smoking one or more packs of cigarettes per day increased substantially and steadily in all grades from 1995 through 2004, with 62%, 68%, and 74% of 8th, 10th, and 12th graders seeing great risk in 2004. Since then, changes have been small and uneven, and the corresponding figures in 2016 are only slightly changed, at 61%, 72%, and 77%. Disapproval of smoking one or more packs of cigarettes per day has increased somewhat steadily in all three grades since 1996 and has reached very high levels. In 2016 disapproval stands at 88%, 89%, and 85% in grades 8, 10, and 12, respectively.

It seems likely that some of the attitudinal change surrounding cigarettes is attributable to the considerable adverse publicity aimed at the tobacco industry in the 1990s, as well as a reduction in cigarette advertising and an increase in antismoking campaigns reaching youth.

Various other attitudes toward smoking became more unfavorable during that interval as well, though most have since leveled off. For example, among 8th graders, the proportions saying that they “prefer to date people who don’t smoke” rose from 71% in 1996 to 81% by 2004, where it remained through 2016. Similar changes occurred in 10th and 12th grades. Thus, at the present time, smoking is likely to make an adolescent less attractive to the great majority of potential romantic age-mates. Likewise, most of the other negative connotations of smoking and smokers have leveled off in the past few years after rising previously. In addition to changes in attitudes and beliefs about smoking, price almost surely also played an important role in the decline in use. Cigarette prices rose appreciably in the late 1990s and early 2000s as cigarette companies tried to cover the costs of the 1998 Master Settlement Agreement, and as many states increased excise taxes on cigarettes. A significant increase in the federal tobacco tax passed in 2009 may have contributed to the continuation of the decline in use since then.

Cigarillos. One consequence of the rise in cigarette prices is that it may have shifted some adolescents to less expensive alternatives, like cigarillos (little or small cigars), which are taxed at a lower rate than cigarettes. Taking into account this form of smoking of tobacco raises the 30-day prevalence of students smoking tobacco—by about three-fourths among 8th and 10th graders and by more than half among 12th graders—over what it would be if just cigarette smoking were counted. It does appear, however, that the prevalence of using

small cigars is also in decline, with 16% of 12th graders in 2016 reporting any *past-year* use, down substantially from 23% in 2010. Of note is the fact that the majority of users of small cigars in each grade smoke flavored ones.

Annual prevalence of smoking tobacco using a **Hookah** (water pipe) had been increasing steadily until 2014 among 12th graders (8th and 10th graders are not asked about this practice), reaching 23% in 2014; but use declined non-significantly by three percentage points to 20% in 2015 and declined significantly (by 6.9 percentage points) to reach 13% in 2016.

Smokeless tobacco. From the mid-1990s to the early 2000s, smokeless tobacco use declined substantially, but a rebound in use developed from the mid-2000s through 2010. Since 2010, prevalence levels have declined modestly in all three grades. Perceived risk and disapproval appear to have played important roles in the earlier decline in smokeless tobacco use. In all three grades, perceived risk and disapproval rose fairly steadily from 1995 through 2004, accompanying the declines in use. However, there was not much change between 2004 and 2010, suggesting that other factors may have led to the increases in smokeless tobacco use during that time interval. : perhaps including increased promotion of these products, a proliferation of types of smokeless tobacco products available, and increased restrictions on places where cigarette smoking is permitted. The decline in smokeless tobacco use since 2010 (including a significant decline among 12th graders in 2015 and 10th graders in 2016) may be attributable, at least in part, to the 2009 increase in federal taxes on tobacco. Perceived risk has not changed appreciably since 2010 at any grade level.

Alcohol remains the substance most widely used by today’s teenagers. Despite recent declines, six out of every ten students (61%) have consumed alcohol (more than just a few sips) by the end of high school, and about a quarter (23%) have done so by 8th grade. In fact, nearly half (46%) of 12th graders and one in eleven (9%) 8th graders in 2016 reported having been drunk at least once in their life.

Alcohol use began a substantial decline in the 1980s. To some degree, alcohol trends have tended to parallel the trends in illicit drug use. These include a modest increase in binge drinking (defined as having five or more drinks in a row at least once in the past two weeks) in the early to mid-1990s, though it was a proportionally smaller increase than was seen for cigarettes and most of the illicit drugs. Fortunately, binge drinking rates leveled off in the early 2000s, just about when the illicit drug rates began to turn around, and in 2002, a drop in **drinking** and **drunkenness** resumed in all grades. Gradual declines continued into 2016, which marked the lowest levels for

alcohol use and drunkenness ever recorded by the survey in the three grades combined. All three grades showed further decline in 2016. The declines in the percentage reporting having been drunk at least once are quite substantial, down from the peak year since 1991 by about 70%, 50%, and 30% for grades 8, 10, and 12.

Vaping

Vaping involves the inhalation of vapors (sometimes including nicotine) using devices such as e-cigarettes, “mods,” and e-pens. Thirty-day prevalence of vaping fell significantly in each grade in 2016. It declined by 1.8 percentage points in 8th grade, by 3.3 percentage points in 10th grade, and by 3.8 percentage points in 12th grade, to levels of 6%, 11%, and 13% in the respective grades.

This marks the first reversal of vaping prevalence, which grew rapidly from near zero prevalence in 2011 to one of the most common forms of adolescent substance use. Despite the decline in 2016 the prevalence of vaping remains substantially higher than the use of any other tobacco product, including cigarettes. Whether teen

vaping has peaked or only paused is an issue that MTF will be able to determine in the coming years.

The percentage of students who associated vaping with “great risk” increased slightly as vaping prevalence declined. *E-cigarettes* are the most commonly used vaping device, and from 2015 to 2016 the percentage of adolescents who believe that regular e-cigarette use poses a great risk of harm increased from 16% to 18% in 12th grade, from 17% to 19% in 10th grade, and from 19% to 21% in 8th grade. Even after these increases, e-cigarettes have some of the lowest levels of perceived risk of any substance.

One reason for low levels of perceived risk may be that the majority of users say that they vape only flavoring not nicotine, on their most recent occasion of use. This is consistent with the fact that only a very few of them (from 5% to 10% of users) say they are using e-cigarettes to help them quit regular cigarette use. The most common reason given for use is “to see what it’s like,” while the second most common reason given is “because it tastes good.”

Any Illicit Drug

MTF routinely reports three different indexes of illicit drug use—any illicit drug,⁷ any illicit drug other than marijuana, and any illicit drug including inhalants. In this section we discuss only the first two; the statistics for all three may be found in Tables 5–7.

In order to make direct comparisons over time, we have kept the definitions and measurement of these indexes constant. The levels of prevalence of each of the indexes could be somewhat affected by the inclusion of newer substances. Typically, the effects would be minimal, primarily because most individuals using newer ones are also using the more prevalent drugs included in the indexes. The major exception has been inhalants, the use of which is quite prevalent in the lower grades, so in 1991 a special index that includes inhalants was added.

Trends in Use

In the late 20th century, U.S. adolescents reached extraordinarily high levels of illicit drug use by U.S. as well as international standards. The trends in *lifetime* use of **any illicit drug** are shown in the first (upper left) panel on the facing page.⁸ In 1975, when MTF began, the majority of young people (55%) had used an illicit drug by the time they left high school. This figure rose to two thirds (66%) in 1981 before a long and gradual decline to 41% by 1992—the low point. After 1992—in what we have called the “relapse phase” in the drug epidemic—the proportion rose considerably to a recent high point of 55% in 1999; it then declined gradually to 47% in 2009, and has remained between 48% and 50% since 2011.

Trends for *annual*, as opposed to lifetime, prevalence are shown in the second (upper right) panel. They are quite parallel to those for lifetime prevalence, but at a lower level. Among 8th graders, a gradual and continuing falloff occurred after 1996. Peak rates since 1991 were reached in 1997 in the two upper grades and declined little for several years. Between 2001 and 2007 all three grades showed declines, but the annual use rates in all three grades rose some through 2012. In 2016, 8th and 10th grades showed some decline, while

annual prevalence among 12th grader showed essentially no change.

Because marijuana is much more prevalent than any other illicit drug, trends in its use tend to drive the index of any illicit drug use. Thus we also report an index that excludes marijuana and shows the proportions of students who use any of the other illicit drugs. The proportions who have used **any illicit drug other than marijuana** in their *lifetimes* are shown in the third panel (lower left). In 1975 over one third (36%) of 12th graders had tried some illicit drug other than marijuana. This figure rose to 43% by 1981, then declined for over a decade to a low of 25% in 1992. An increase followed in the 1990s as the use of a number of drugs rose steadily, and it reached 30% by 1997. (In 2001 it was 31%, but this apparent upward shift in the estimate was an artifact due to a change in the question wording for “other hallucinogens” and tranquilizers.⁹) Lifetime prevalence among 12th graders then fell slightly to 24% by 2009, before dropping to 21% by 2016. The fourth (lower right) panel presents the *annual* prevalence data for any illicit drug other than marijuana, which shows a pattern of change over the past few years similar to the index of any illicit drug use, but with much less pronounced change since 1991. It dropped fairly steadily and gradually in all three grades in recent years but leveled in 2013 before dropping to 14% among 12th graders by 2016. In fact, prevalence declined in all three grades in 2016—significantly so in 8th grade.





Overall, these data reveal that while use of individual drugs (other than marijuana) may fluctuate widely, the proportion using *any* of them is much more stable. In other words, the proportion of students prone to using such drugs and willing to cross the normative barriers to such use changes more gradually. The usage rate for each individual drug, on the other hand, reflects many more rapidly changing determinants specific to that drug, such as how widely its psychoactive potential is recognized, how favorable the reports of its supposed benefits are, how risky its use is seen to be, how acceptable it is in the peer group, how accessible it is, and so on.

⁷Footnote ‘a’ to Tables 5 through 8 provides the exact definition of any illicit drug.

⁸This is the only set of figures in this *Overview* presenting lifetime use statistics. Lifetime statistics for all drugs may be found in Table 5.

⁹The term psychedelics was replaced with hallucinogens, and shrooms was added to the list of examples, resulting in somewhat more respondents indicating use of this class of drugs. For tranquilizers, Xanax was added to the list of examples given, slightly raising the reported prevalence of use.

Any Illicit Drug and Any Illicit Drug Other than Marijuana: Trends in Lifetime and Annual Use
 Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

*In 2001, a revised set of questions on other hallucinogen use and tranquilizer use were introduced. In 2013, a revised set of questions on amphetamine use was introduced. Data for any illicit drug other than marijuana were affected by these changes.

Marijuana

Marijuana has been the most widely used illicit drug throughout MTF's 42-year history. It can be taken orally, mixed with food or drink, and smoked, including in a concentrated form as hashish—the use of which is much more common in Europe.¹⁰ The great majority of recreational use in the U.S. involves smoking it in rolled cigarettes (“joints”), in pipes or water pipes (“bongs”), in hollowed-out cigars (“blunts”), or more recently in a vaporizer. Newer methods also include smoking or eating different forms of resin extracts like hash oil, honey oil, or shatter—a solid form.

Trends in Use

Annual marijuana prevalence peaked among 12th graders in 1979 at 51%, following a rise that began during the 1960s. Then use declined fairly steadily for 13 years, bottoming at 22% in 1992—a decline of more than half. The 1990s, however, saw a resurgence of use. After a considerable increase (one that actually began among 8th graders a year earlier than among 10th and 12th graders), annual prevalence rates peaked in 1996 at 8th grade and in 1997 at 10th and 12th grades. After these peak years, use declined among all three grades through 2007 or 2008. After these declines, an upturn occurred in use in all three grades, lasting for three years in the lower grades and longer in grade 12. Annual marijuana prevalence among 8th graders increased in use from 2007 to 2010, decreased slightly from 2010 to 2012, and then declined significantly in 2016. Among 10th graders, use increased somewhat from 2008 to 2013 and then declined after that. Among 12th graders, use increased from 2006 to 2011 and then held level through 2016. As shown in Table 8, *daily* use increased in all three grades after 2007, reaching peaks in 2011 (at 1.3% in 8th), 2013 (at 4.0% in 10th), and 2011 (at 6.6% in 12th), before declining slightly since. Daily prevalence rates in 2016 were 0.7%, 2.5%, and 6.0%, respectively, with one in seventeen 12th graders smoking daily.

Perceived Risk

The proportion of students seeing great risk from smoking marijuana regularly fell during the rise in use in the 1970s and again during the subsequent rise in use in the 1990s. Indeed, for 10th and 12th grades, perceived risk declined a year before use rose in the upturn of the 1990s, making perceived risk a leading indicator of

change in use. (The same may have happened for 8th grade as well, but we lack data starting early enough to know.) The decline in perceived risk halted in 1996 in 8th and 10th grades; the increases in use in 10th and 12th grades ended a year or two later, again making perceived risk a leading indicator of trends in use. From 1996 to 2000, perceived risk held fairly steady, and the decline in use in the upper grades stalled. After some decline prior to 2002, perceived risk increased a bit in all grades through 2004 as use decreased. Since 2004 in 8th grade, 2005 in 12th grade, and 2008 in 10th grade, perceived risk has fallen substantially, presaging some resurgence in marijuana use lasting three to five years; but no increase in perceived risk preceded the recent leveling of use. Rather, perceived risk has continued a steep decline since the mid-2000s without a concomitant further rise in overall use. Disapproval and availability may be constraining factors offsetting the effects of risk. Recent, sharp declines in the use of “gateway drugs”—in particular cigarette smoking, with which marijuana use has been highly correlated—may also be playing a role

Disapproval





Personal disapproval of trying marijuana has declined some since 2007 or 2008 in all three grades, but still remains quite high with 82%, 74%, and 69% in 8th, 10th, and 12th grades still saying that they disapprove of regular use. Disapproval fell considerably among 8th graders between 1991 and 1996 and among 10th and 12th graders between 1992 and 1997—by 17, 21, and 19 percentage points, respectively, over those intervals of increasing use and declining perceived risk. As is often the case, perceived risk fell before disapproval. Since 2008 there has been some decline in disapproval.

Availability

Since the MTF study began in 1975, between 80% and 90% of 12th graders each year have said that they could get marijuana fairly easily or very easily if they wanted some, with that figure standing at 81% in 2016. Marijuana has been considerably less readily available to 8th graders, with 35% in 2016 reporting it to be fairly or very easy to get. Availability is intermediate for the 10th graders, with 64% reporting easy access in 2016. Availability has declined appreciably, especially among the younger adolescents, but marijuana remains readily available to most 12th graders.

¹⁰Compared with Europe, American teens have high rates of illicit drug use <http://monitoringthefuture.org/pressreleases/16ESPADpr.pdf>

Marijuana: Trends in Annual Use, Risk, Disapproval, and Availability
 Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

Synthetic Marijuana

Synthetic marijuana has generally been sold over the counter under such labels as Spice and K-2. It usually contains some herbal materials that have been sprayed with one or more of the designer chemicals that fall into the cannabinoid family. Until March 2011, these drugs were not scheduled by the Drug Enforcement Administration (DEA), so they were readily and legally available on the Internet and in convenience stores, head shops, gas stations, etc. However, the DEA scheduled some of the most widely used chemicals beginning March 1, 2011, making their possession and sale no longer legal; subsequent laws have expanded the list of banned chemicals, but producers keep tweaking the chemical formula to avoid legal control. These drugs can be dangerous both because the active ingredients keep changing and because those ingredients have never undergone testing to determine their effects on humans.

Trends in Use

MTF first addressed the use of synthetic marijuana in its 2011 survey by asking 12th graders about their use in the prior 12 months (which would have covered a considerable period of time prior to the drugs being scheduled). Annual prevalence was found to be 11.4%, making synthetic marijuana the second most widely used class of illicit drug after marijuana itself among 12th graders at that time. Despite the DEA's intervention, use among 12th graders remained unchanged in 2012 at 11.3%, which suggests either that

compliance with the new scheduling had been limited or that producers of these products succeeded in continuing to change their chemical formulas to avoid using the ingredients that had been scheduled. In 2012, for the first time, 8th and 10th graders were asked about their use of synthetic marijuana; their annual prevalence rates were 4.4% and 8.8%, respectively. Use in all 3 grades dropped in 2013, and the decline was sharp and significant among 12th graders, while the declines were significant for both 10th and 12th graders in 2014 and continued through 2016, with significant declines in both 10th and 12th grades. Annual prevalence in 2016 was down to 2.7%, 3.3%, and 3.5% for the three grades, reflecting a dramatic drop in use.





Perceived Risk

All three grades were asked whether they associated great risk with trying synthetic marijuana once or twice. As can be seen on the facing page, the level of perceived risk for experimental use was quite low in 2012 (between 24% and 25%) but has risen some, particularly among 12th graders, to 36% in 2016. (Likely the percent would be higher if those answering "Can't say, Drug unfamiliar" were excluded.) The availability of these drugs over the counter probably has had the effect of communicating to teens that they must be safe, though they are not.

Disapproval and Availability have not been measured for this class of drugs.

Synthetic Marijuana: Trends in Annual Use and Risk

Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

Inhalants

Inhalants are any gases or fumes that can be inhaled for the purpose of getting high. These include many household products—the sale and possession of which is legal—including glue, nail polish remover, gasoline, solvents, butane, and propellants used in certain commercial products such as whipped cream dispensers. Unlike nearly all other classes of drugs, their use is most common among younger adolescents and tends to decline as youth grow older. The use of inhalants at an early age may reflect the fact that many inhalants are cheap, readily available (often in the home), and legal to buy and possess. The decline in use with age likely reflects their coming to be seen as “kids’ drugs,” in addition to the fact that a number of other drugs become available to older adolescents, who are also more able to afford them.

Trends in Use

Inhalant use (excluding the use of nitrite inhalants) by 12th graders rose gradually from 1976 to 1987, which was somewhat unusual because most other forms of illicit drug use were in decline during the 1980s. Use of inhalants rose among 8th and 10th graders from 1991, when those grades were first included in the study, through 1995; it rose among 12th graders from 1992 to 1995. All grades then exhibited a fairly steady and substantial decline in use through 2001 or 2002. After 2001 the grades diverged somewhat in their trends: 8th graders showed a significant increase in use for two years, followed by a decline from 2004 to 2013, and a leveling in 2014, before resuming the decline in 2015 and 2016; 10th graders showed an increase after 2003 but a considerable decline since 2007; and 12th graders showed a brief increase from 2003 to 2005 but also a considerable decline since then. For the three grades combined, annual use declined significantly in both 2012 and 2013, held steady in 2014 and then declined further in 2015 and 2016.

Perceived Risk

Only 8th and 10th graders have been asked questions about the degree of risk they associate with inhalant

use. Relatively low proportions think that there is a “great risk” in using an inhalant once or twice. However, significant increases in this belief were observed between 1995 and 1996 in both 8th and 10th grades, probably due to an anti-inhalant advertising initiative launched by The Partnership for a Drug-Free America. That increase in perceived risk marked the beginning of a long and important decline in inhalant use, when no other drugs showed a turnaround in use. However, the degree of risk associated with inhalant use declined steadily between 2001 and 2008 among both 8th and 10th graders, perhaps explaining the increase in use in 2003 among 8th graders and in 2004 in the upper grades. The hazards of inhalant use were communicated during the mid-1990s, but generational forgetting of those hazards has likely taken place as replacement cohorts who were too young to get that earlier message now comprise the nation’s adolescents. The decline in perceived risk is worrisome, though perceived risk has not changed much since about 2008. In this case, the decline in perceived risk (between 2001 and 2008) did not translate into a large surge in use, but it may leave future class cohorts at risk for a resurgence of inhalant use.





Disapproval

Over 80% of 8th and 10th grade students say that they would disapprove of even trying an inhalant. (The question was not asked of 12th graders.) There was a very gradual upward drift in this attitude among 8th and 10th graders from 1995 through about 2001, with a gradual falloff since then in both grades. For 8th graders there has been some decline in disapproval of trying inhalants since 2012. Since 2013 it has dropped among 10th graders as well, including significant declines in 2015 and a continuing decline in 2016.

Availability

Respondents have not been asked about the availability of inhalants, because we assume that these products are universally available to young people in these age ranges.

Inhalants: Trends in Annual Use, Risk, and Disapproval
 Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

For some years, LSD was the most widely used drug within the larger class of hallucinogens. This was no longer true for some following years, due to sharp decreases in its use combined with an increasing use of psilocybin. (Statistics on overall hallucinogen use and on use of hallucinogens other than LSD are shown in the tables at the end of this report.) Now overall hallucinogen use and use of hallucinogens other than LSD are about equivalent due to a drop in the use of the other hallucinogens.

Trends in Use

Annual prevalence of LSD use among 12th graders has been below 10% since MTF began. Use declined some for the first 10 years among 12th graders, likely continuing a decline that had begun before 1975. Use was fairly level in the latter half of the 1980s but, as was true for a number of other drugs, rose in all three grades between 1991 and 1996. Between 1996 and 2006 or so, use declined in all three grades, with particularly sharp declines between 2001 and 2003. Since then use has remained at very low levels although there has been a slight increase in the upper grades since 2013.

Perceived Risk

We think it likely that perceived risk for LSD use increased during the early 1970s, before MTF began, as concerns grew about possible neurological and genetic effects (most of which were never scientifically confirmed) as well as “bad trips” and “flashbacks.” However, there was some decline in perceived risk in the late 1970s, after which it remained fairly level among 12th graders through most of the 1980s. A substantial decline occurred in all grades in the early 1990s as use rose. Since about 2000, perceived risk declined steadily and substantially among 8th graders until 2007, when it leveled; it declined considerably among 10th graders before leveling around 2002, dropping through 2007, and then leveling after that. Among 12th graders, perceived risk has held fairly steady after 2002, at least until 2014 when some decline began. The decline in the lower grades suggests that younger teens may be less knowledgeable about this drug’s effects than their predecessors—through what we have called “generational forgetting”—making them vulnerable to a resurgence in use. (The percentages who respond “can’t say, drug unfamiliar” to questions about LSD have risen in recent years,

consistent with the notion of “generational forgetting.”)

The decline of LSD use in recent years, despite a fall in perceived risk, suggests that some factors other than a change in underlying attitudes and beliefs were contributing to the downturn—prior to 2001 some displacement by ecstasy may have been a factor, while more recently a decline in availability (discussed below) likely is a factor.





Disapproval

Disapproval of LSD use was quite high among 12th graders through most of the 1980s but began to decline after 1991 along with perceived risk. All three grades exhibited a decline in disapproval through 1996, with disapproval of experimentation dropping 11 percentage points between 1991 and 1996 among 12th graders. After 1996 a slight increase in disapproval emerged among 12th graders, accompanied by a leveling among 10th graders and some further decline among 8th graders. Since 2001, disapproval of LSD use has diverged among the three grades, declining considerably among 8th graders, declining less among 10th graders, and increasing significantly among 12th graders. Note, however, that the percentages of 8th and 10th graders who respond with “can’t say, drug unfamiliar” increased through 2008; thus the base for disapproval has shrunk, suggesting that the real decline of disapproval among the younger students is less than it appears here. There has been rather little change in disapproval over the past eight years (or more years in the case of 12th graders.)

Availability

Reported availability of LSD by 12th graders fell considerably from 1975 to 1979, declined a bit further until 1986, and then began a substantial rise, reaching a peak in 1995. LSD availability also rose somewhat among 8th and 10th graders in the early 1990s, reaching a peak in 1995 or 1996. Since those peak years, there has been considerable falloff in availability in all three grades, quite possibly in part because fewer students have LSD-using friends from whom they could gain access. There was also very likely a decrease in supply due to the closing of a major LSD-producing lab by the Drug Enforcement Administration in 2000. It is clear that attitudinal changes cannot explain the recent declines in use.

LSD: Trends in Annual Use, Risk, Disapproval, and Availability
 Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

Cocaine was used almost exclusively in powder form for some years, though “freebasing” emerged for a while. The early 1980s brought the advent of crack cocaine. Our original questions did not distinguish among different forms of cocaine or modes of administration. Since 1987, though, we have asked separate questions about the use of crack and “cocaine other than crack,” which has consisted almost entirely of powder cocaine use. Data on cocaine use in general (i. e., all forms of cocaine) are presented in the figures in this section, and results for crack alone are presented in the next section.

Trends in Use

There have been some important changes in the levels of overall cocaine use over the life of MTF. Use among 12th graders originally burgeoned in the late 1970s and remained fairly stable through the first half of the 1980s before starting a precipitous decline after 1986. Annual prevalence among 12th graders dropped by about three quarters between 1986 and 1992. Between 1992 and 1999, use reversed course again during the relapse phase of the overall drug epidemic and doubled before declining by 2000. Use also rose among 8th and 10th graders after 1992 before reaching peak levels in 1998 and 1999. Over the last sixteen years, use has declined in all three grades; annual 12th-grade use stands at a historic low of just 2.3% in 2016, with use by 8th and 10th graders still lower, at 0.8% and 1.3%.

Perceived Risk

Questions about the dangers of cocaine in general (without specifying any particular form of cocaine) have been asked only of 12th graders. The results tell a fascinating story. They show that perceived risk for experimental use fell in the latter half of the 1970s (when use was rising), stayed level in the first half of the 1980s (when use was level), and then jumped very sharply in a single year (by 14 percentage points between 1986 and 1987), just when the substantial decline in use began. The year 1986 was marked by a national media frenzy over crack cocaine and also by the widely publicized cocaine-related death of Len Bias, a National Basketball Association first-round draft pick. Bias’ death was originally reported as resulting from his first experience with cocaine. Though that was later proven to be incorrect, the

message had already “taken.” We believe that this event helped to persuade many young people that use of cocaine at any level is dangerous, no matter how healthy the individual.¹¹ Perceived risk continued to rise through 1991 as the fall in use continued. Perceived risk declined modestly from 1991 to 2000, and use rose from 1992–2000. Perceived risk has leveled in recent years at far higher levels than existed prior to 1987, and there was a gradual upward drift for about six years in grades 8 and 10, before leveling. For the 12th graders perceived risk also increased for about six years before leveling after 2013. There is as yet little evidence of generational forgetting of cocaine’s risks. For 12th graders, survey questions on both risk and disapproval referred to cocaine in general, until 1986. After that they referred to cocaine powder and crack separately, as did the questions asked of 8th and 10th graders. The question change seemed to matter rather little in the results.

Disapproval





Disapproval of cocaine use by 12th graders followed a cross-time pattern similar to that for perceived risk, although its seven-percentage-point jump in 1987 was not quite as pronounced. Some decline from 1991 to 1997 was followed by a period of stability. Subsequent years showed a gradual increase in disapproval in all three grades. This upward drift ended in recent years, but disapproval of even trying cocaine remains very high and is above 85% in all grades in 2016.

Availability

The proportion of 12th graders saying that it would be “fairly easy” or “very easy” for them to get cocaine if they wanted some was 33% in 1977, rose to 48% by 1980 as use rose, and held fairly level through 1982; it increased steadily to 59% by 1989 (in a period of rapidly declining use). Perceived availability then fell back to about 47% by 1994. Since around 1997, perceived availability of cocaine has fallen considerably in all three grades. Among 12th graders it stood at 29% in 2016—about half of its peak level in 1989. Note that the pattern of change does not map well onto the pattern of actual use, suggesting that changes in overall availability have not been a major determinant of use—particularly during the sharp decline in use in the late 1980s.

¹¹ Among 12th graders trends in perceived risk in Table 8 show a particularly sharp rise from 34% in 1986 to 48% in 1987 for trying cocaine once or twice.

Cocaine (including Crack): Trends in Annual Use, Risk, Disapproval, and Availability
 Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

*Prior to 1991, data reported here is based on questions on use of cocaine in general. Starting in 1991, data based on questions on use of cocaine powder specifically.

Crack

Several indirect indicators suggest that crack use grew rapidly in the period 1983–1986, beginning before we had direct measures of its use. In 1986 a single usage question was included in one of the five 12th-grade questionnaire forms, asking those who indicated any cocaine use in the prior 12 months if they had used crack. The results from that question represent the first data point in the first panel on the facing page. After that, three questions about crack use covering the usual three prevalence periods were introduced into several questionnaire forms.

Trends in Use

Clearly crack use rose rapidly in the early 1980s, judging by the 4% annual prevalence reached in 1986; but after 1986 there was a precipitous drop in crack use among 12th graders; the drop continued through 1991. After 1991 for 8th and 10th graders (when data were first available) and after 1993 for 12th graders, all three grades showed a slow, steady increase in use through 1998 during the relapse phase of the overall drug epidemic. Since 1999, annual prevalence dropped by about 75% in 8th grade, 82% in 10th grade, and 70% in 12th grade. Today use of crack is at historic lows in all three grades. As with many drugs, the decline at 12th grade lagged behind those in the lower grades due to a cohort effect.

Perceived Risk

By the time we added questions about the perceived risk of using crack in 1987, crack was already seen by 12th graders as one of the most dangerous illicit drugs: 57% saw a great risk in even trying it. This compared to 54% for heroin, for example. Perceived risk for crack rose still higher through 1990, reaching 64% of 12th graders who said they thought there was a great risk in taking crack once or twice. (Use was dropping during that interval.) After 1990 some falloff in perceived risk began, well before crack use began to increase in 1994, making perceived risk again a leading indicator. Between 1991 and 1998 there was a considerable falloff in this belief in grades 8 and 10, as use rose steadily. Perceived risk leveled in 2000 in

grades 8 and 12 and a year later in grade 10. We think that the declines in perceived risk for crack and cocaine during the 1990s may well reflect an example of generational forgetting wherein the class cohorts that were in adolescence when the adverse consequences were most obvious (i.e., in the mid-1980s) were replaced by cohorts who were less knowledgeable about the dangers. By 2016 perceived risk for crack had leveled in all three grades.

Disapproval





Disapproval of crack use was not assessed until 1990, when it was at a very high level, with 92% of 12th graders saying that they disapproved of even trying it. Disapproval of crack use declined slightly but steadily in all three grades from 1991 through about 1997. After 1997, disapproval in all three grades rose back to high levels by 2012 before beginning a slight decline.

Availability

Crack availability did not change dramatically in the early years for which data are available. It began a sustained decline after 1995 among 8th graders, after 1999 among 10th graders, and after 2000 among 12th graders. Since 2000, availability has declined considerably, particularly in the upper grades.

NOTE: The distinction between crack cocaine and other forms of cocaine (mostly powder) was made several years after the study's inception. The figures on the facing page begin their trend lines when these distinctions were introduced. Figures are not presented here for the "other forms of cocaine" measures, simply because the trend curves look extremely similar to those for crack. (All statistics are contained in the tables presented later.) Although the trends are very similar, the absolute levels of use, risk, etc., are somewhat different. Usage levels tend to be higher for cocaine powder compared to crack, and the levels of perceived risk a bit lower, while disapproval has been close for the two different forms of cocaine and relative availability has varied (Tables 15 through 17).

Crack: Trends in Annual Use, Risk, Disapproval, and Availability
Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

Amphetamines and Other Stimulant Drugs

Amphetamines, a class of psychotherapeutic stimulants, had a relatively high prevalence of use in the youth population for many years. The behavior reported here excludes any use under medical supervision. Amphetamines are controlled substances—they are not legally bought or sold without a doctor’s prescription—but some are diverted from legitimate channels, and some are manufactured and/or imported illegally. There are other controlled stimulants that are also included, like Ritalin which is used to treat ADHD, as is Adderall, the most prevalent of the amphetamines.

Trends in Use

The use of these stimulants rose in the last half of the 1970s, reaching a peak in annual prevalence of 26% in 1981 (likely exaggerated due to commonly used “look-alikes”)—two years after marijuana use peaked. From 1981 to 1992, 12th graders reported a steady and very substantial decline in their use, reaching 7%.

As with many other illicit drugs, these stimulants made a comeback in the 1990s. Use peaked in the lower two grades by 1996. Since then, use declined steadily in 8th grade and sporadically in 10th grade. Only after 2002 did it begin to decline in 12th grade. The decline paused in 2008 for 8th graders and 2009/2010 for 12th graders, and then resumed. The 12th-grade decline began in 2003 but reversed from 2009 to 2013. In 2013 the amphetamines/stimulants prevalence question text was changed in half of the questionnaire forms. The 2013 report used data from the changed forms only, to be comparable to the 2014 measure. In 2014 the remaining forms were changed; the 2014 and subsequent data presented here are for all the forms. From 2009 to 2013 use rose in the upper grades, likely due to use intended to assist with academic performance. Since 2013 there has been a downward drift in annual prevalence but a steeper decline in 30-day prevalence (significant in the upper grades).

See Table 6 for the trends in annual use of two specific amphetamines—Ritalin and Adderall. Since it was first measured in 2001, Ritalin use has declined by 70% to 80% in all three grades. Adderall use declined in the lower grades since it was first measured in 2009; but annual prevalence increased significantly in 12th grade between 2009 (5.4%) and 2013 (7.4%) where it remained in 2015 before falling to 6.2% in 2016.

Perceived Risk

Only 12th graders are asked about the amount of risk they associate with amphetamine/stimulant use. For a few years, changes in perceived risk were not correlated with changes in usage levels (at the aggregate level). Specifically, in the interval 1981–1986, risk was quite stable even though use fell considerably, likely as a result of some displacement by increasing cocaine use. There was, however, a decrease in risk during the period 1975–1981 (when use was rising), some increase in perceived risk in 1986–1991 (when use was falling), and some decline in perceived risk from 1991 to 1995 (in advance of use rising again). Perceived risk has generally been rising in recent years, very likely contributing to the decline in use that occurred among 12th graders after 2002; it appears to have leveled after 2007. In 2011 the examples of specific amphetamines provided in the text of the questions on perceived risk, disapproval, and availability were updated with the inclusion of Adderall and Ritalin. This led to some discontinuities in the trend lines in 2011. (Levels of perceived risk and disapproval lowered as a result.) Based on the revised question, some decline has occurred since 2013.





Disapproval

Disapproval of amphetamine/stimulant use is asked in 12th grade only. Relatively high proportions of 12th graders have disapproved of even trying amphetamines/stimulants throughout the life of the study. Disapproval did not change in the late 1970s despite an increase in use. From 1981 to 1992, disapproval rose gradually and substantially from 71% to 87% as perceived risk rose and use declined. In the mid-1990s disapproval declined along with perceived risk, but it increased fairly steadily from 1996 through 2009 before leveling. There has been a slight falloff since 2013.

Availability

In 1975, amphetamines/stimulants had a high level of reported availability. The level fell by about 10 percentage points by 1977, drifted up a bit through 1980, jumped sharply in 1981, and then began a long, gradual decline through 1991. There was a modest increase in availability at all three grade levels in the early 1990s as use rose, followed by a long-term decline which continued through 2016.

Amphetamines: Trends in Annual Use, Risk, Disapproval, and Availability
Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

*In 2013 the question text was changed on two of the questionnaire forms for 8th and 10th graders and four of the questionnaire forms for 12th graders, and changed on the remaining forms in 2014. Beginning in 2013, data presented here include only the changed forms.

**In 2011 the list of examples was changed from uppers, pep pills, bennies, speed to uppers, speed, Adderall, Ritalin, etc. These changes likely explain the discontinuity in the 2011 results.

Methamphetamine and Crystal Methamphetamine (Ice)

One subclass of amphetamines is called methamphetamine (“speed”). This subclass has been around for a long time and gave rise to the phrase “speed kills” in the 1960s. Probably because of the reputation it got at that time as a particularly dangerous drug, it was not popular for some years, so we did not include a full set of questions about its use in MTF’s early questionnaires. One form of methamphetamine, crystal methamphetamine or “ice,” grew in popularity in the 1980s. It comes in crystallized form, as the name implies, and the chunks can be heated and the fumes inhaled, much like crack cocaine.

Trends in Use

For most of the life of the study, the only question about *methamphetamine* use has been contained in one of the six 12th-grade questionnaire forms. Respondents who indicated using any type of amphetamines in the prior 12 months were asked in a sequel question to indicate on a pre-specified list the types they had used during that period. Methamphetamine was one type on the list, and data exist on its use since 1976. (The rates are not graphed here until 1990.) In 1976, annual prevalence using this measure was 1.9%; it then roughly doubled to 3.7% by 1981 (the peak year), before declining for over a decade all the way down to 0.4% by 1992. Use then rose again in the mid-1990s, as did use of a number of drugs, reaching 1.3% by 1998. In other words, it has followed a cross-time trajectory fairly similar to that for amphetamines as a whole. No questions have yet been added to the study on perceived risk, disapproval, or availability with regard to overall methamphetamine use.

In 1990, in the 12th-grade questionnaires only, we introduced our usual set of three questions for *crystal methamphetamine*, measuring lifetime, annual, and 30-day use. Among 12th graders in 1990, 1.3% indicated any use in the prior year; use climbed to 3.0% by 1998, and has generally been declining since then, reaching an all-time low of 0.5% in 2015 and then 0.8% in 2016. This variable is charted on the first panel of the facing page.

Responding to the growing concern about methamphetamine use in general—not just crystal methamphetamine use—we added a full set of three

questions about the use of any methamphetamine to the 1999 questionnaires for all three grade levels. These questions yield a somewhat higher annual prevalence for 12th graders: 4.3% in 2000, compared to the sum of the methamphetamine and crystal methamphetamine answers in the other, branching question format, which totaled 2.8%. It would appear, then, that the long-term method we had been using for tracking methamphetamine use probably yielded an understatement of the absolute prevalence level, perhaps because some proportion of methamphetamine users did not correctly categorize themselves initially as amphetamine users (even though methamphetamine was given as one of the examples of amphetamines). We think it likely that the shape of the trend curve was not distorted, however.

The newer questions for *methamphetamine* (not graphed here) show annual prevalence rates in 2016 of 0.4% for both 8th and 10th graders, and 0.6% for 12th graders. In every grade, 2016 prevalence is at the lowest level ever recorded by the survey. The 2016 levels for all three grades are down considerably from the first measurement taken in 1999, when they were 3.2%, 4.6%, and 4.7% (see Table 6). So, despite growing public concern about the methamphetamine problem in the United States, use actually has shown a fairly steady and substantial decline since 1999, at least among secondary school students. (A similar decline in methamphetamine use did not begin to appear among college students and young adults generally until after 2004, likely reflecting a cohort effect. See [Volume II](#) in this series for data on adults through age 55.)





Other Measures

Data on perceived risk and availability for crystal methamphetamine, specifically, may be found on the facing page.

Clearly, the perceived risk of crystal methamphetamine use has risen considerably since 2003, very likely explaining much of the decline in use since then. Perceived risk then leveled after 2013. Perceived availability generally has been falling in all three grades since 2006, perhaps in part because there are many fewer crystal methamphetamine users from whom to gain access.

Crystal Methamphetamine (Ice): Trends in Annual Use, Risk, and Availability

Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

Heroin

For many decades, heroin—a derivative of opium—was administered primarily by injection into a vein. However, in the 1990s the purity of available heroin reached very high levels, making other modes of administration (e.g., snorting, smoking) practical alternatives. Thus, in 1995 we introduced questions that asked separately about using heroin with and without a needle to determine whether non-injection use explained the upsurge in heroin use we observed. The usage statistics presented on the facing page are based on heroin use by any method, but data on the two specific types of administration are provided in the tables at the end of this report.

Trends in Use

The annual prevalence of heroin use among 12th graders fell by half between 1975 and 1979, from 1.0% to 0.5%. The rate then held amazingly steady until 1994. Use rose in the mid- and late-1990s, along with the use of most drugs; it reached peak levels in 1996 among 8th graders (1.6%), in 1997 among 10th graders (1.4%), and in 2000 among 12th graders (1.5%), suggesting a cohort effect. Following those peak levels, use declined, with annual prevalence in all three grades fluctuating between 0.7% and 0.9% from 2005 through 2010. Since then, annual prevalence in the three grades combined has declined, from 0.8% to 0.3% in 2016. In 2016, use reached its lowest levels since 1991 in all three grades (0.3% in each).

Because the questions about use with and without a needle were not introduced until the 1995 survey, they did not encompass much of the period of increasing heroin use. Responses to the new questions showed that, by then, about equal proportions of all 8th-grade users were taking heroin by each method of ingestion and some—nearly a third of users—were using both means. At 10th grade, a somewhat higher proportion of all users took heroin without a needle, and at 12th grade, the proportion was higher still. Much of the increase in overall heroin use after 1995 occurred in the proportions using it without injecting, which we strongly suspect was true in the immediately preceding period of increase as well. Likewise, much of the decrease since the recent peak levels has been due to decreasing use of heroin without a needle. In 2012, there were significant decreases in use of heroin without a needle for 8th and 12th graders, and very slight declines since then in 8th and 10th grades.

Use with a needle has fallen considerably in all three

grades since the mid-1990s; annual prevalence in 2016 stood at 0.2%, 0.3%, and 0.3%, respectively, including significant declines in 8th and 10th grades from the 2014 to 2015 prevalence levels, but no further change in 2016. The proportional declines were greatest in the lower grades. While a heroin epidemic continues among adults, our data—as well as those from NSDUH—suggest that use has grown primarily among young adults and not among adolescents.

Perceived Risk

Students have long seen heroin to be one of the most dangerous drugs, which helps to account for both the consistently high level of personal disapproval of use (see below) and the quite low prevalence of use. Nevertheless, perceived risk levels have changed some over the years. Between 1975 and 1986, perceived risk gradually declined, even though use dropped and then stabilized in that interval. Then there was a big spike in 1987 (when perceived risk for cocaine also jumped dramatically), where it held for four years. In 1992, perceived risk dropped to a lower plateau again, presaging an increase in use a year or two later. Perceived risk rose in the latter half of the 1990s, and use leveled off and then declined. Risk at 12th grade is still rising, but has been level for some time at 8th and 10th grades. Perceived risk of use without a needle rose in 8th and 10th grades between 1995 and 1997, foretelling an end to the increase in that form of use. Note that perceived risk has served as a leading indicator of use for this drug as well as a number of others. During the 2000s, perceived risk was relatively stable.





Disapproval

There has been little fluctuation in the very high levels of disapproval of heroin use over the years, though it did rise gradually between 2000 and 2010. The small changes that have occurred have been generally consistent with changes in perceived risk and use.

Availability

The proportion of 12th-grade students saying they could get heroin fairly easily if they wanted some remained around 20% through the mid-1980s. It then increased considerably from 1986 to 1992 before stabilizing at about 35% from 1992 through 1998. From the mid- to late-1990s through 2014, perceived availability of heroin declined gradually but substantially in all three grades before leveling in 2014 or 2015.

Heroin: Trends in Annual Use, Risk, Disapproval, and Availability
 Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

*Prior to 1995, the questions asked about heroin use in general. Since 1995, the questions have asked about heroin use without a needle.

Other Narcotic Drugs, Including OxyContin and Vicodin

There are a number of narcotic drugs other than heroin—all controlled substances. Many are analgesics that can be prescribed by physicians and dentists for pain. Like heroin, many are derived from opium, but there are also a number of synthetic analogues in use today, with OxyContin and Vicodin being two of the major ones.

Throughout the life of the MTF study, we have asked about the use of any narcotic drug other than heroin without specifying which one. Examples of drugs in the class are provided in the question stem. In one of the six 12th-grade questionnaire forms, however, respondents indicating that they had used any narcotic in the past 12 months were then asked to check which of a fairly long list of such drugs they used. Table E-4 in Appendix E of [Volume I](#) of this annual monograph series provides trends in their annual prevalence data. In the late 1970s, opium and codeine were among the narcotics most widely used. In recent years Vicodin, codeine, Percocet, OxyContin, and hydrocodone have been the most prevalent.

Trends in Use

Use is reported for 12th graders only, because we considered the data from 8th and 10th graders to be of questionable validity. As shown in the first panel of the facing page, 12th graders' use of narcotics other than heroin generally trended down from about 1977 through 1992, dropping considerably. After 1992 use rose rather steeply as all forms of substance use were increasing, with annual prevalence nearly tripling from 3.3% in 1992 to 9.5% in 2004, before leveling through about 2009. Since then, use has been declining, particularly since 2009.

In 2002, the question was revised to add Vicodin, OxyContin, and Percocet to the examples given, which clearly had the effect of increasing reported prevalence, as may be seen in the first panel on the facing page. So the extent of the increase over the full time span likely is exaggerated, although probably not by much, because these drugs came onto the scene later, during the rise.

They simply were not being fully reported until the late 1990s. Narcotics had become one of the most widely used classes of illicit drugs by 2004, when annual prevalence reached 9.5%.

Use rates for two narcotics of recent interest—OxyContin and Vicodin—are presented in the second and third panels on the facing page, in a departure from the usual arrangement. There are no data for disapproval and only limited data on perceived risk (since 2012) for the two drugs, showing low and stable risk levels.

OxyContin use increased some in all grades from 2002 (when it was first measured) through roughly 2009, though the trend lines have been irregular. Since 2009 or 2010, the prevalence rate has dropped in all grades. Annual prevalence in 2016 was 0.9%, 2.1%, and 3.4% in grades 8, 10, and 12, respectively. Use of **Vicodin**, on the other hand, remained fairly steady at somewhat higher levels from 2002—the first year it was measured—until 2009, after which it declined substantially in all grades. In 2016, annual prevalence rates continued to decline and were 0.8%, 1.7%, and 2.9% for 8th, 10th and 12th graders respectively.

Availability


Questions were asked about the availability of narcotics other than heroin, taken as a class. Perceived availability increased gradually among 12th graders for more than a decade (from 1978 through 1989), even as reported use was dropping. Perceived availability then rose further for another decade (from 1991 through 2001) as use rose quite sharply before leveling by about 2000 and then declining after 2006. In contrast, perceived availability had declined among 8th and 10th graders since the late 1990s. (In all three grades, a change in question wording in 2010 to include OxyContin and Vicodin as examples presumably accounts for the jump in reported availability that year.) Availability has declined further in all three grades since 2010.

Narcotics other than Heroin and OxyContin and Vicodin Specifically:


Trends in Annual Use and Availability

Grades 8, 10, 12


Use of Narcotics other than Heroin
% who used any narcotics other than heroin
in last 12 months*




OxyContin Use
% who used OxyContin in last 12 months



Vicodin Use
% who used Vicodin in last 12 months



Availability of Narcotics other than Heroin**
% saying "fairly easy" or "very easy" to get



Source. The Monitoring the Future study, the University of Michigan.

*Beginning in 2002, a revised set of questions on other narcotics use was introduced in which Talwin, laudanum, and paregoric were replaced as examples given with Vicodin, OxyContin, and Percocet.

**In 2010 the list of examples was changed from methadone, opium to Vicodin, OxyContin, Percocet, etc.

Tranquilizers

Tranquilizers are psychotherapeutic drugs that are legally sold only by prescription. They are central nervous depressants and, for the most part, comprise benzodiazepines (minor tranquilizers), although some non-benzodiazepines have been introduced. Respondents are instructed to exclude any medically prescribed use from their answers. At present, *Xanax* is the tranquilizer most commonly used by 12th graders (only 12th graders are asked to indicate which specific tranquilizers they used). (See Table E-3 in appendix E of [Volume I](#) in this series for details.) Valium, Klonopin, and Ativan are other tranquilizers, used at somewhat lower levels. In 2001, the examples given in the tranquilizer question were modified to reflect changes in the drugs in common use—Miltown was dropped and Xanax was added. As the first panel on the facing page shows, this caused a modest increase in the reported level of tranquilizer use in the upper grades, so we have broken the trend line to reflect the point of redefinition.

Trends in Use

During the late 1970s and all of the 1980s, tranquilizers fell steadily and substantially from popularity, with 12th graders' use declining by three fourths over the 15-year interval between 1977 and 1992. Their use then increased, as happened with many other drugs during the 1990s. Annual prevalence more than doubled among 12th graders, rising steadily through 2002, before leveling. Use also rose steadily among 10th graders, but began to decline some in 2002. Use peaked much earlier among 8th graders in 1996 and then declined slightly for two years. Tranquilizer use remained relatively stable among 8th graders through 2010 at considerably lower levels than the upper two grades. Use in 8th grade showed declines in 2011 and

2012 before stabilizing again. From 2002 to 2005, there was some decline among 10th graders, followed by a leveling, then a resumption of the decline through 2014 before drifting up again. Among 12th graders, there was a very gradual decline from 2002 through 2007, before leveling and then decreasing in 2010 and again in 2013. This staggered pattern of change suggests that a cohort effect has been at work. There has been little further change since 2013. In 2016, the prevalence of use of these prescription-type drugs was somewhat lower than their recent peak levels, with annual prevalence rates of 1.7%, 4.1%, and 4.9% in grades 8, 10, and 12, respectively.





Perceived Risk and Disapproval

Data have not been collected on perceived risk and disapproval for tranquilizers, primarily due to questionnaire space limitations.

Availability

As the number of 12th graders reporting non-medically prescribed tranquilizer use fell dramatically during the 1970s and 1980s, so did the proportion saying that tranquilizers would be fairly or very easy to get. Whether declining use caused the decline in availability or vice versa is unclear. However, 12th graders' perceived availability has continued to fall since then, even as use rebounded in the 1990s; it is now down by eight tenths over the life of the study—from 72% in 1975 to 15% by 2016 saying that tranquilizers would be fairly or very easy to get if they wanted some. Availability has fallen fairly continuously since 1991 in the lower grades as well, though not as sharply. In 2016 all three grades showed a slight upward movement in availability (significantly so in 8th grade).

Tranquilizers: Trends in Annual Use and Availability Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

*Beginning in 2001, a revised set of questions on tranquilizer use was introduced in which Xanax replaced Miltown in the list of examples.

Sedatives (Barbiturates)

Like tranquilizers, sedatives are prescription-controlled psychotherapeutic drugs that act as central nervous system depressants. They are used to assist sleep and relieve anxiety.

Though for many years respondents have been asked specifically about their use of barbiturate sedatives, they likely have been including other classes of sedatives in their answers. In 2004, the question on use was revised to say “sedatives/barbiturates”—a change that appeared to have no impact on reported levels of use. Respondents are told for what purposes sedatives are prescribed and are instructed to exclude from their answers any use under medical supervision. Usage data are reported only for 12th graders because we believe that 8th- and 10th-grade students tend to over-report use, perhaps including in their answers their use of nonprescription sleep aids or other over-the-counter drugs.

Trends in Use

As with tranquilizers, the use of sedatives (barbiturates) fell steadily among 12th graders from the mid-1970s through the early 1990s. From 1975 to 1992, annual prevalence fell by three fourths, from 10.7% to 2.8%. As with many other drugs, a gradual, long-term resurgence in sedative use occurred after 1992, but unlike the case with most illegal drugs, sedative (barbiturate) use continued to rise steadily through 2005, well beyond the point at which the use of most illegal drugs began falling. (Recall that tranquilizer use also continued to rise into the early 2000s.) Use has declined some since 2005, and by 2016, the annual prevalence rate was down by about six-tenths from its recent peak, falling to 3.0%. The sedative methaqualone (known as Quaaludes) was included in the MTF study from the very beginning, and was never as popular as barbiturates; use rates have generally been declining since 1975, reaching an annual prevalence of just 0.5% in 2007, about where it remained through 2012, after which the question was dropped.

Perceived Risk

Trying sedatives (barbiturates) was never seen by most students as very dangerous; and it is clear from the upper

right panel on the facing page that changes in perceived risk cannot explain the trends in use that occurred from 1975 through 1986, when perceived risk was actually declining along with use. But then perceived risk shifted up some through 1991 while use was still falling. It dropped back some through 1995, as use was increasing, and then remained relatively stable for a few years. Perceived risk has generally been at quite low levels, which may help to explain why the use of this class of psychotherapeutic drugs (and likely others) continued to grow in the first half of the decade of the 2000s. However, perceived risk began to rise a bit after 2000, foretelling the decline in use that began after 2005. When the term “sedatives” was changed to “sedatives/barbiturates” in 2004, the trend line shifted down slightly, but perceived risk continued to climb gradually through 2013, before turning down. Prior to that point use declined as perceived risk rose.





Disapproval

Like many illicit drugs other than marijuana, sedative (barbiturate) use has received the disapproval of most high school seniors since 1975, with some variation in disapproval rates that have moved consistently with usage patterns. The change in question wording in 2004 appeared to lessen disapproval slightly. There has been a modest increase in disapproval since 2000, although that appears to have stopped in 2014 and was followed by a slight decrease afterwards.

Availability

As the fourth panel on the facing page shows, the perceived availability of sedatives (barbiturates) has generally been declining during most of the life of the study, except for one upward shift that occurred in 1981—a year in which look-alike drugs became more widespread. (The change in question text in 2004 appears to have had the effect of increasing reported availability among 12th graders but not among students in the lower grades.) Perceived availability for sedatives (barbiturates) continued its long-term decline into 2015; in 2016, it declined significantly among 10th graders, with no significant change among 8th and 12th graders.

Sedatives (Barbiturates): Trends in Annual Use, Risk, Disapproval, and Availability
 Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

*In 2004 the question text was changed. Barbiturates was changed to Sedatives, including barbiturates and "have you taken barbiturates..." was changed to "have you taken sedatives..." In the list of examples downs, downers, goofballs, yellows, reds, blues, rainbows were changed to downs, or downers, and include Phenobarbital, Tuinal, and Seconal.

**In 2004 the question text was changed from barbiturates to sedatives/barbiturates and the list of examples was changed from downers, goofballs, reds, yellows, etc. to just downers. These changes likely explain the discontinuity in the 2004 results.

MDMA (Ecstasy, Molly) and Other “Club Drugs”

“Club drugs,” so called because they have been popular at nightclubs and raves, include LSD, MDMA (known as ecstasy, and more recently, Molly), methamphetamine, GHB (gamma-hydroxybutyrate), ketamine (special K), and Rohypnol. (For discussion of LSD and methamphetamine, see prior pages.) We focus here initially on MDMA (ecstasy, Molly) and treat the other drugs in the last section below.

Trends in MDMA (Ecstasy, Molly) Use

Ecstasy (3, 4-methylenedioxymethamphetamine or MDMA) is used more for its mildly hallucinogenic properties than for its stimulant properties. Questions on ecstasy use were added to the surveys in 1996.

In 1996, annual prevalence of ecstasy use was 4.6% in both 10th and 12th grades—considerably higher than among college students (2.8%) and young adults (1.7%)—but use declined over the next two years. Use then rose sharply, bringing annual prevalence up to 3.5%, 6.2%, and 9.2% for 8th, 10th, and 12th graders by 2001. From 2001 to 2005, use declined substantially, down to 1.7%, 2.6%, and 3.0%, respectively. Following some irregular changes in recent years, in 2014 compared to 2005, use was down slightly in 8th grade (to 0.9%) and 10th grade (to 2.3%) and up slightly in 12th grade (to 3.6%). “Molly,” reputedly a purer form of MDMA, received much attention in 2013. Because that term was not used in the 2013 questionnaires, it is not clear whether students included it in their answers about ecstasy use that year. The inclusion of Molly as an example in some of the 2014 questionnaires seemed to make a modest difference in reported prevalence. (The 2014 data reported here show one point based on the unmodified questionnaires and another based on the modified ones.) Since 2014, the change had been downward and significantly so in 2016 in all three grades, despite the inclusion of Molly.

Perceived Risk

In 2001, 12th graders’ perceived risk of ecstasy use jumped by 8 percentage points and in 2002, by another seven. Significant increases occurred in 2003 for all grades. This sharp rise in perceived risk likely caused the drop in use, as we predicted. From 2004 to 2011, we saw a troubling drop in perceived risk (first among 8th and 10th, and then among 12th graders), corresponding to the increase in use in the upper two grades and then in all three grades. This suggests a generational forgetting of the dangers of ecstasy use. In 2012, only 8th graders showed much further decline. The rebound in use after 2004

might be explained by the sizable drop in perceived risk. The addition of Molly as an example caused a considerable jump in perceived risk after 2013 in grades 8 and 10, suggesting that they see it as more dangerous than ecstasy.

Disapproval

Disapproval of ecstasy use declined some after 1998 but increased significantly in all three grades in 2002, perhaps due to the rise in perceived risk. The rise in disapproval continued through 2003 for 8th, 2004 for 10th, and 2006 for 12th graders, suggesting some cohort effect. After those peaks, disapproval dropped sharply among 8th graders and less among 10th graders before leveling, and it did not drop among 12th graders until 2010—again suggesting a cohort effect. The erosion in perceived risk and disapproval—which was sharpest among 8th graders—left these groups more vulnerable to a possible rebound in use; some rebound appears to have occurred during the past decade.





Availability

The figure shows a dramatic rise in 12th graders’ perceived availability of ecstasy after 1991, particularly between 1999 and 2001, consistent with informal reports about growing importation of the drug. Perceived availability then declined considerably in all grades, including significant declines in 2016 at 10th and 12th grades. Decreased availability may help account for the declines in use in the past few years.

Rohypnol, GHB, and Ketamine

Rohypnol and **GHB** are labeled date rape drugs because they can have amnesiac effects and can be added to food or drink without a victim’s knowledge. By 2011, both drugs had shown significant declines since their peak levels of annual use (Table 6). In 2011, annual prevalence for Rohypnol use was 0.8%, 0.6%, and 1.3%, and for GHB use, 0.6%, 0.5%, and 1.4% in grades 8, 10, and 12, respectively. Annual prevalence for another club drug, **ketamine**, had also shown significant declines, and was at 0.8%, 1.2%, and 1.7% in 2011. Questions about GHB and ketamine use were dropped from the surveys of 8th and 10th graders in 2012. In 2016, annual prevalence among 12th graders for Rohypnol, GHB, and ketamine was 1.1%, 0.9%, and 1.2%, respectively. Annual prevalence of Rohypnol was 0.5% for 8th graders and 0.5% for 10th graders. No questions about risk, disapproval, or availability are asked for these drugs.

Ecstasy (MDMA): Trends in Annual Use, Risk, Disapproval, and Availability
 Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

*In 2014/2015, revised sets of questions on ecstasy were introduced in which molly was added to the description. This likely explains the discontinuity in the results for those years.

Alcohol

Alcohol has been widely used by young people in the U.S. for a very long time. In 2016, the proportions of 8th, 10th, and 12th graders who reported drinking an alcoholic beverage in the 30-day period prior to the survey were 7%, 20%, and 33%, respectively. Various measures of alcohol use are presented in the tables at the end of this report. Here we focus on episodic heavy or “binge” drinking (i.e., having five or more drinks in a row on one or more occasions in the prior two weeks) because heavy alcohol consumption is probably of greatest concern from a public health perspective. In 2016 lifetime, annual, 30-day, and binge drinking measures of alcohol use were at historic lows over the life of the study in all three grades (8, 10 and 12).

Trends in Use

Among 12th graders, binge drinking peaked in 1979 along with overall illicit drug use. The prevalence of binge drinking then declined substantially from 41% in 1983 to 28% in 1992, a drop of almost one third (also the low point of any illicit drug use). Although illicit drug use rose sharply in the 1990s, binge-drinking rose by only a small fraction, and that rise was followed by some decline at all three grades. By 2016, proportional declines since the recent peaks reached in the 1990s were 75%, 60%, and 51% for grades 8, 10, and 12, respectively (Table 8). The observed prevalence of binge drinking continued its decline from 2015 to 2016 (significant in 8th grade), to 2016 rates of 3%, 10%, and 16% for the 3 grades.

It should be noted that there is no evidence of any displacement effect in the aggregate between alcohol and marijuana—a hypothesis frequently heard. The two drugs have moved much more in parallel over the decades than in opposite directions, at least until about a five-year period in the 2000s, during which alcohol continued to decline while marijuana reversed course and rose. Moreover, these two behaviors have consistently been positively correlated at the individual level.

Perceived Risk

Across the past four decades, since the MTF study began,

the majority of 12th graders have not viewed binge drinking on weekends as carrying a great risk. However, an increase from 36% to 49% occurred between 1982 and 1992. A decline to 43% followed by 1997 as use rose, before it stabilized. Since 2003, perceived risk has risen some in all grades, at least through 2011 or 2012. These changes are consistent with changes in actual binge drinking. We believe that the public service advertising campaigns in the 1980s against drunk driving, as well as those that urged use of designated drivers when drinking, contributed to the increase in perceived risk of binge drinking generally. Drunk driving by 12th graders declined during that period by an even larger proportion than binge drinking. Also, we showed that increases in the minimum drinking age during the 1980s were followed by reductions in drinking and increases in perceived risk associated with drinking, policy-driven effects that may still be deterring alcohol use among adolescents.





Disapproval

Disapproval of weekend binge drinking moved fairly parallel with perceived risk, suggesting that such drinking (and very likely the drunk-driving behavior associated with it) became increasingly unacceptable in the peer group. Note that the rates of disapproval and perceived risk for binge drinking are higher in the lower grades than in 12th grade. As with perceived risk, disapproval increased appreciably in all grades, though it leveled in 2015.

Availability

Perceived availability of alcohol, which until 1999 was asked only of 8th and 10th graders, was very high and mostly steady in the early 1990s. Since 1996, however, there have been substantial declines in 8th and 10th grades. For 12th grade, availability has declined only modestly with 85% in 2016 still saying that alcohol would be fairly or very easy to get. Overall, it appears that states, communities, and parents have been successful in reducing access to alcohol, particularly among the younger teens.

Alcohol: Trends in Binge Drinking, Risk, Disapproval, and Availability
 Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

Cigarettes

Cigarette smoking is the leading cause of preventable disease and mortality in the United States, and is usually initiated in adolescence. That makes what happens with cigarette smoking in adolescence particularly important to study.

Trends in Use

Differences in smoking rates between various birth cohorts (or, in this case, school class cohorts) tend to stay with those cohorts throughout the life cycle. This means that it is critical to prevent smoking very early. It also means that the trends in a given historical period may differ across various grade levels as changes in use occurring earlier in adolescence work their way up the age spectrum (i.e., “cohort effects”).

Among 12th graders, 30-day prevalence of smoking reached a peak in 1976 at 39%. (The peak likely occurred considerably earlier at lower grade levels as these same class cohorts passed through them in previous years.) After about a one-quarter drop in 12th-grade 30-day prevalence between 1976 and 1981, the rate remained remarkably stable until 1992 (28%). In the 1990s, smoking began to rise sharply, after 1991 among 8th and 10th graders and after 1992 among 12th graders. Over the next four to five years, smoking rates increased by about one half in the lower two grades and by almost one third in grade 12—very substantial increases to which MTF drew considerable public attention. Smoking peaked in 1996 for 8th and 10th graders and in 1997 for 12th graders before beginning a fairly steady and substantial decline that continued through 2004 for 8th and 10th graders. Between the peak levels in the mid-1990s and 2004, 30-day prevalence of smoking declined by 56% in 8th grade, 47% in 10th, and 32% in 12th. This important decline in adolescent smoking decelerated after about 2002. Still, by 2016, 30-day prevalence levels had fallen from peak levels by 87%, 84%, and 71% in grades 8, 10, and 12, respectively. An increase in 2009 in federal taxes on cigarettes (from \$0.39 to \$1.01 per pack) may have contributed to the recent decline in use. Of particular importance, smoking initiation by 8th graders declined by four fifths from a peak of 49% in 1996 to 10% by 2016.

Perceived Risk

Among 12th graders, the proportion seeing great risk in pack-a-day smoking rose before and during the first period of decline in use in the late 1970s. It leveled in 1980 (before use leveled), declined a bit in 1982, but then started to rise again gradually for five years. (It is possible that cigarette advertising effectively offset the influence of rising perceptions of risk during that period.) Perceived risk fell some in the early 1990s at all three grade levels as use increased sharply. Since then, there has generally been an increase (though not entirely consistently over the years) in perceived risk, reaching in 2015 the highest levels yet observed in grades 8 and 10 and close to the highest in grade 12. Risk has fallen back some in 8th and 10th grades over the past year. Note the differences in the extent of perceived risk among grade levels. There is a clear age effect: by the time most youngsters fully appreciate the hazards of smoking, many already have initiated the behavior.


Disapproval

Disapproval rates for smoking have been fairly high throughout the study and, unlike perceived risk, are higher in the lower grade levels. Among 12th graders, there was a gradual increase in disapproval of smoking from 1976 to 1986, followed by some erosion over the decade through 1997. After 1997, disapproval rose for some years in all three grades, but leveled briefly after 2006 or 2007, before rising even more. We measure a number of other smoking-related attitudes; these became increasingly negative, but leveled off six or seven years ago (see Table 3 in the [2016 MTF press release on teen tobacco use](#)). Though disapproval has continued to increase, some attitudes and beliefs about cigarette smoking are no longer moving in a direction that would discourage use, suggesting that external changes in the environment may be required to further reduce youth smoking.

Availability

Since 1996, availability has declined considerably among 8th and 10th graders. Some 46% of 8th graders and 63% of 10th graders now say that cigarettes would be very easy or fairly easy to get, down from 78% in 1992 among 8th graders and 91% in 1995 among 10th graders.

Cigarettes: Trends in 30-Day Use, Risk, Disapproval, and Availability
 Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

Smokeless Tobacco

Traditionally, smokeless tobacco has come in two forms: “snuff” and “chew.” Snuff is finely ground tobacco usually sold in tins, either loose or in packets. It is held in the mouth between the lip or cheek and the gums. Chew is a leafy form of tobacco, usually sold in pouches. It too is held in the mouth and may, as the name implies, be chewed. In both cases, nicotine is absorbed by the mucous membranes of the mouth. These forms are sometimes called “spit” tobacco because users expectorate the tobacco juices and saliva (stimulated by the tobacco) that accumulate in the mouth. “*Snus*” (rhymes with goose) is a relatively new variation on smokeless tobacco, as are some other *dissolvable tobacco* products that literally dissolve in the mouth. Given that snus appeared to be gaining in popularity, separate items regarding the use in the past 12 months of snus and dissolvable tobacco were added to the 12th-grade surveys in 2011 and to the 8th- and 10th-grade surveys in 2012. In addition, in 2011 snus and dissolvable tobacco were added as examples in the long-standing question on smokeless tobacco.

Trends in Use

The use of smokeless tobacco by teens had been decreasing gradually, and 30-day prevalence is now about half of the recent peak levels in the mid-1990s, though there was a reversal of the declines from about 2007 through 2010. Among 8th graders, 30-day prevalence declined from a 1994 peak of 7.7% to 3.2% in 2007. It reached a low of 2.8% in 2013, and then fell even lower to 2.5% in 2016. Among 10th graders, use declined from a 1994 peak of 10.5% to 4.9% by 2004, and then rose to 6.4% in 2013 before dropping again to 3.5% in 2016. Among 12th graders, 30-day use declined from a 1995 peak of 12.2% to 6.1% by 2006 then rose to 8.5% in 2010, before falling back to 6.6% in 2016. Thirty-day prevalence of daily use of smokeless tobacco fell gradually but appreciably for some years. Daily usage rates in 2016 were 0.6%, 1.0%, and 2.7% in grades 8, 10, and 12, respectively—down substantially from peak levels recorded in the 1990s—but most of the declines occurred in the 1990s, not since.

Smokeless tobacco use among American young people is almost exclusively a male behavior. Among males, the 30-day prevalence rates in 2016 were 3.6%, 5.8%, and

11.9% in grades 8, 10, and 12, versus 1.4%, 1.3%, and 1.5% for females. The respective current daily use rates for males were 1.2%, 1.7%, and 5.1% compared to 0.1%, 0.2%, and 0.3% for females.

Annual prevalence in 2016 for *snus* was 2.2%, 3.0%, and 5.8% among 8th, 10th, and 12th graders, respectively, reflecting a decline since 2012 in all three grades. For *dissolvable tobacco*, the corresponding figures were 0.7%, 0.9%, and 1.1%, reflecting little change since 2012.

Perceived Risk

The most recent low point in the level of perceived risk for smokeless tobacco was 1995 in all three grades (though for 12th graders it was considerably lower in the mid-1980s). For a decade following 1995, there was a gradual but substantial increase in proportions saying that there is a great risk in using smokeless tobacco regularly. It thus appears that one important reason for the appreciable declines in smokeless tobacco use during the latter half of the 1990s was that an increasing proportion of young people were persuaded of the dangers of using it. However, the increases in perceived risk ended by 2004 in 12th grade, and it has declined some in the interval since then in all grades. The decline could be due to generational forgetting of the dangers of use, the increased marketing of snus and other smokeless products, and/or public statements about smokeless tobacco use being relatively less dangerous than cigarette smoking. By 2016, perceived risk leveled in all three grades.





Disapproval

Only 8th and 10th graders are asked about their personal disapproval of using smokeless tobacco regularly. The most recent low points for disapproval in both grades were 1995 and 1996. Disapproval rose among 8th graders from 74% in 1996 to 82% in 2005, about where it was in 2016 (81%). For 10th graders, disapproval rose from 71% in 1996 to 82% in 2008, also about where it was in 2016 (81%).

Availability

There are no questions on perceived availability of smokeless tobacco.

Smokeless Tobacco: Trends in 30-Day Use, Risk, and Disapproval
 Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

Vaping

Vaping involves the use of a battery-powered device to heat a liquid or plant material so that it releases chemicals in an aerosol, vapor, or mist that users inhale. Examples of vaping devices include e-cigarettes, “mods,” and e-pens. The vaping process reduces the number of harmful chemicals that users ingest in comparison to traditional, combustion-based forms of smoking. The vapor may contain nicotine, the active ingredients of marijuana, flavored propylene glycol, and/or flavored vegetable glycerin. The liquid that is vaporized comes in hundreds of flavors, many of which (e.g., bubble gum and milk chocolate cream) are likely to be attractive to younger teens.

Since 2015, MTF has included questions about vaping with any device, and asks respondents if they have used an “electronic vaporizer such as e-cigarettes.” Since 2014, the survey has included questions about the use of the specific vaping device of an e-cigarette.

Trends in Use

Vaping declined in 2016, which marks the first significant reversal of a rapid rise in adolescent vaping. Previous to 2016 prevalence of vaping had grown from near-zero levels in 2011 to one of the most common forms of adolescent substance use by 2015. From 2015 to 2016, the percentage of students who vaped in the last 30 days declined by 2 or 3 percentage points to 6%, 11%, and 13% among 8th, 10th, and 12th graders, respectively. Each of these declines was statistically significant.

A similar decline was also present in use of e-cigarettes, which are specific vaping devices, in the last 30 days. From 2015 to 2016, e-cigarette prevalence declined significantly in all three grades-- from 10% to

6% among 8th-grade students, from 14% to 10% among 10th grade students, and from 16% to 12% among 12th grade students.

Despite these declines, vaping continued to have higher use among teens than traditional tobacco cigarettes or any other tobacco product. As a point of comparison, prevalence for tobacco cigarettes was 2.6%, 4.9%, and 10.5% among 8th, 10th, and 12th grade students respectively. Note that in 8th and 10th grades vaping is more than twice as common as use of regular cigarettes.

Perceived Risk

Perceived risk of vaping has increased as prevalence has declined. E-cigarettes are by far the most common vaping device, and the percentage of adolescents who believe that regular e-cigarette use poses a risk of harm increased from 19% to 21% in 8th grade, from 17% to 19% in 10th grade, and from 16% to 18% in 12th grade. However, even after these increases e-cigarettes have one of the lowest levels of perceived risk for regular use of all drugs, including alcohol.

Disapproval





Disapproval of regular use of e-cigarettes is also relatively low, compared to most other substances. However, it did rise in 2016 from 65% to 67% in 8th grade and from 60% to 65% in 10th grade (the increase was statistically significant in 10th grade but not in 8th grade; the question is not asked of 12th graders.)

Availability

Data on availability of vaping devices or e-cigarettes have not been gathered so far.

Electronic Vaporizers: Trends in 30-Day Use

Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.

Small Cigars, Cigarillos, Large Cigars, and Tobacco Using a Hookah

Twelfth graders were first asked about smoking small cigars and smoking tobacco using a hookah (water pipe) in 2010. These questions were not asked of 8th and 10th graders. Only the prevalence and frequency of use in the past 12 months were reported; we use this prevalence period, requiring only a single question (which we call a “tripwire” question) to determine whether additional questions on the substance may be warranted in future surveys. Small cigar and hookah use are charted on the facing page.

Smoking Tobacco Using a Hookah. The past 12 months prevalence of hookah use had been rising since it was first measured in 2010, from 17.1% in 2010 to 22.9% in 2014; but it declined sharply in 2015 and 2016 to 13.0%. Only about 7% of the 12th-grade students in 2016 indicated use on more than two occasions during the prior 12 months, suggesting that a considerable amount of hookah use is light or experimental. (Males are only slightly more likely than females to use hookahs, at 15% for males and 11% for females in 2016.)

Small Cigars. Small or little cigars are the approximate size and shape of a cigarette, but they are classified as cigars because they are wrapped in brown paper, which contains some tobacco leaf, rather than in white paper. In 2016, the annual prevalence for small or little cigars (our question uses the term “small cigars”) was 16%. Smoking small cigars has declined significantly since 2010, when annual prevalence was 23%. Unlike hookah smoking, use of small cigars shows a sizable gender difference: the 2016 annual prevalence for 12th grade males was 23% compared to 9% for females. The increases in the federal taxes on tobacco products, instituted in 2009, may well have played a role in decreasing the use of small cigars. The increase on a pack of small cigars fell under the same regulations as regular cigarettes (rising from \$0.39 to \$1.01 per pack). Some producers of small cigars subsequently increased

the weight of their cigars slightly (taxation is based on weight, with cigars falling into a higher weight class with a lower tax rate) in order to avoid the higher taxes placed on cigarettes and to remove them from FDA control under current law. Eight percent of 12th graders indicated having used small cigars on more than two occasions during the past year, and only 1% on more than 20 occasions, so they tend to be smoked much less frequently than regular cigarettes. Some small cigars are flavored, which is likely to make them more attractive to young people. A concern in the public health community is that these products will have the effect of reversing the hard-won gains in reducing cigarette smoking among youth. Small cigars contain nicotine and combust tobacco in a similar way, and therefore carry similar dangers.


Small (Little) Cigars and Cigarillos. In a set of questions introduced in 2014 we asked about the use in the prior 30 days of little cigars OR cigarillos. (Cigarillos lie between little cigars and large cigars in size—length and thickness—and are wrapped in tobacco leaf like large cigars. They fall into the lower federal taxation bracket than cigarettes.) The distinction is made between flavored and unflavored (regular) little cigars or cigarillos, and it shows that the flavored ones are more widely used by teens. There was no significant change between 2014 and 2015 in the 30-day prevalence of either type, but in 2016, there were declines in all 3 grades, significant in 8th and 12th (Table 7). Thirty-day prevalence in 2016 was 2.8%, 4.9%, and 9.5% for flavored and 1.9%, 3.0%, and 6.1% for regular small cigars or cigarillos in grades 8, 10, and 12, respectively.

Large Cigars. A question on the 30-day prevalence of smoking large cigars also was added in 2014. The rates were 1.5%, 2.3%, and 6.5% in 2016—with all three grades showing declines in 2016 (significant in 8th and 10th grades).


Small Cigars and Tobacco using a Hookah: Trends in Annual Use

Grade 12


Small Cigar Use
% who used in last 12 months




Use of Tobacco with a Hookah
% who used in last 12 months



Disapproval
% disapproving of using once or twice



Availability
% saying "fairly easy" or "very easy" to get



Source. The Monitoring the Future study, the University of Michigan.

Steroids

Unlike all other drugs discussed in this Overview, anabolic steroids are not usually taken for their psychoactive effects, though they may have some, but rather for muscle and strength development. However, they are similar to most other drugs studied here in two respects: they are controlled substances for which there is an illicit market, and they can have adverse consequences for the user. Questions about steroid use were added beginning in 1989. Respondents are asked: “Steroids, or anabolic steroids, are sometimes prescribed by doctors to promote healing from certain types of injuries. Some athletes, and others, have used them to try to increase muscle development. The question asks, “On how many occasions (if any) have you taken steroids on your own—that is, without a doctor telling you to take them?” In 2006, the question text was changed slightly in some questionnaire forms—the phrase “to promote healing from certain types of injuries” was replaced by “to treat certain conditions.” The resulting data did not show any effect from this rewording. In 2007, the remaining forms were changed in the same manner.

Trends in Use

Anabolic steroids have been used predominately by males; therefore, data based on all respondents can mask the higher rates and larger fluctuations that occur among males. (For example, in 2016, annual prevalence rates were 0.5%, 0.9%, and 1.3% for boys in grades 8, 10, and 12, compared with 0.5%, 0.5%, and 0.6% for girls.) Between 1991 and 1998, the overall annual prevalence rate was fairly stable among 8th and 10th graders, ranging between 0.9% and 1.2%. In 1999, however, use jumped from 1.2% to 1.7% in both 8th and 10th grades. (Almost all of that increase occurred among boys, increasing from 1.6% in 1998 to 2.5% in 1999 in 8th grade and from 1.9% to 2.8% in 10th grade. Thus, rates among boys increased by about half in a single year. By 2016 among all 8th graders, steroid use had declined by about two thirds to 0.5%. Among 10th graders, use continued to increase, reaching 2.2% in 2002, but then declined by about two thirds to 0.7% by 2016. In 12th grade, there was a different trend story. With data going back to 1989, we can see that steroid use first fell from 1.9% overall in 1989 to 1.1% in 1992—the low point. From 1992 to 2000, there was a more gradual increase in use, reaching 1.7% in 2000. In 2001, use rose significantly

among 12th graders to 2.4% (possibly reflecting a cohort effect). Twelfth graders’ use decreased significantly in 2005 to 1.5%, then stayed fairly level through 2015 (1.7%), and then declined significantly in 2016 to 1.0%. Use is now down from recent peak levels by about two-thirds among 8th and 10th graders, and about six-tenths among 12th graders. (The use of androstenedione—a steroid precursor—has also declined sharply since 2001, most sharply through 2007. It was classified as a Schedule II controlled substance in 2005 by the DEA.)

Perceived Risk

Perceived risk and disapproval were asked of 8th and 10th graders for only a few years. All grades seemed to have a peak in perceived risk around 1993. The longer-term data from 12th graders show a ten percentage-point drop between 1998 and 2000. A change this sharp is quite unusual and highly significant, suggesting that some particular event or events in 1998—quite possibly publicity about use of androstenedione by a famous home-run-hitting baseball player—made steroids seem less risky. It seems likely that perceived risk dropped substantially in the lower grades as well, consistent with the sharp upturn in their use that year. By 2006, perceived risk for 12th graders was up to 60%, with little change until 2013 when it showed a significant 4.4 percentage point decline, reaching 54%, the lowest point ever. It stands at 55% in 2016.





Disapproval

Among 12th graders, disapproval of steroid use has been quite high for some years. Between 1998 and 2003, there was a modest decrease, though not as dramatic as the drop in perceived risk. From 2003 to 2008, disapproval rose some—as perceived risk rose and use declined—then leveled and declined from 2012 through 2014, before leveling.

Availability

Perceived availability of steroids was relatively high prior to 2001 or 2002, but it declined appreciably at all grades through 2016 (2015 in the case of the 8th graders) reaching the lowest levels recorded by the study. A number of steroids have been scheduled by the DEA, no doubt contributing to the drop in availability.

Steroids: Trends in Annual Use, Risk, Disapproval, and Availability
 Grades 8, 10, 12



Source. The Monitoring the Future study, the University of Michigan.
 *Question discontinued in 8th- and 10th-grade questionnaires in 1995.

Subgroup Differences

Understanding the important subgroup variations in substance use among the nation's youth allows for more informed considerations of substance use etiology and prevention. It also helps to prioritize prevention and treatment efforts. In this section, we present a brief overview of some of the major demographic subgroup differences.

Space does not permit a full discussion or documentation of the many subgroup differences of the drugs covered in this report. However, the forthcoming *Volume I* in this series contains tables providing the 2016 subgroup prevalence levels for all of the classes of drugs discussed here; Chapters 4 and 5 in *Volume I* have in-depth discussion and interpretation of those subgroup differences. Comparisons are made by gender, race/ethnicity, college plans, region of the country, community size, and socioeconomic level (as measured by educational level of the parents). In addition, an annual Monitoring the Future Occasional Paper provides tables giving cross-time trends in the subgroup prevalence levels for all of the classes of drugs discussed here and, importantly, charts showing the subgroup trends for all drugs. This Occasional Paper, *Demographic subgroup trends among adolescents in the use of various licit and illicit drugs 1975-2016*, is Number 88 in the series and contains data through 2016. It is available on the MTF website: <http://www.monitoringthefuture.org/pubs/occpapers/mtf-occ88.pdf>. The graphs in the occasional paper present easily accessible views of trends and comparisons while the occasional paper's tables provide the specific numbers behind the figures.

Gender

Generally, males have somewhat higher rates of illicit drug use than females (especially higher rates of frequent use), most notably by 12th grade. Males in all three grades have much higher rates of smokeless tobacco and steroid use and, in the upper grades, higher rates of use of small cigars, large cigars, dissolvable tobacco, and snus specifically. The primary exception may be found in the misuse of prescription drugs like amphetamines, sedatives, and tranquilizers, where females have tended to have higher rates of use than males in the early grades. But misuse of prescription narcotics males: their use is reported only at grade 12 and males have consistently had higher rates of use. For most drugs, though, the gender differences among 8th graders are very small, with females fairly consistently

reporting slightly higher rates than males through 2015; in 2016 males were equal to or higher than females in the use of several drugs. Among 10th graders, males have generally, though not always, reported higher rates than females.

Among 12th graders, for many years males consistently reported distinctly higher 30-day alcohol usage rates than females; however, the differences have been narrowing and in 2016 females have slightly higher prevalences in 8th and 10th grades and only a slightly lower one in 12th grade (32% vs. 35%). Gender differences in binge drinking have followed a similar pattern—females report higher or the same rates in 8th grade, males somewhat higher rates in 10th grade until 2016, and males higher rates in 12th grade (though again the gap is narrowing).

Gender differences in 30-day cigarette smoking among 8th and 10th graders have generally been minimal, but 10th grade males have reported slightly higher rates than females in recent years. Among 12th graders, females generally had higher rates of smoking than males through 1990, but since then males have generally had the higher rates (13% vs. 8% in 2016).

The gender differences in substance use appear to emerge for many drugs as students grow older. In 8th grade, females have higher rates of use for some drugs, such as inhalants and amphetamines. Prevalence rates for both genders then increase with age (with the single exception of inhalants), but the increase is often sharper among males. At each grade level, usage rates for both genders generally tend to move much in parallel across time for the various substances, and the absolute differences between the genders tend to be largest in the historical periods in which overall prevalence rates are highest.

Race/Ethnicity

Among the most dramatic and interesting subgroup differences are those found among the three largest racial/ethnic groups—Whites, African Americans, and Hispanics. For a number of years White students had substantially higher rates of using any illicit drug than did African American students, but the differences have narrowed in recent years as a result of increasing marijuana use among African American students and a leveling among White students. (Marijuana use tends to drive the overall index of any illicit drug use and in

2016 is significantly higher among African American students than among White students in 8th grade and somewhat higher in 10th grade.) Still, African American students have tended to have lower levels of use for certain licit and illicit drugs at all three grade levels—in particular for hallucinogens, synthetic marijuana, and all forms of prescription drugs used without a doctor's orders. For 12th graders, crack and heroin use among African Americans has been higher than among Whites in recent years; and their use of bath salts is now higher in all three grades than bath salts use by either Whites or Hispanics.

African American students' use of alcohol and cigarettes tends to be significantly lower than Whites in all three grades. In fact, African Americans' use of cigarettes has been dramatically lower than Whites' use—a difference that emerged largely during the life of the study (i.e., since 1975).

Hispanic students generally have had rates of use that place them between the other two groups in 12th grade—usually closer to the rates for Whites than for African Americans. In the last few years, however, Hispanics have attained the highest reported rates of use of any illicit drug in all three grades—in large part due to their increase in marijuana use. Indeed, both African Americans and Hispanics have shown a considerably greater increase in marijuana use than Whites, at least until 2016 when their use fell significantly in both grades 8 and 10. In 12th grade Hispanics have the highest use rates for a number of substances—marijuana, synthetic marijuana, inhalants, hallucinogens, LSD, cocaine, crack, methamphetamine, and crystal methamphetamine. In 8th grade, Hispanics have tended to report the highest rates of the three racial/ethnic groups on nearly all classes of drugs. Like African American students, Hispanic students generally have lower rates than White students of misusing any of the prescription drugs, particularly in the upper grades.

Again, we refer the reader to [OP 88](#) for a detailed picture of these complex subgroup differences and how they have changed over the years.

College Plans

While in high school, those students who are not college-bound (a decreasing proportion of the total youth population over the longer term) are considerably more likely to be at risk for using illicit drugs, drinking heavily, and particularly smoking

cigarettes. Again, these differences are largest in periods of highest prevalence. In the lower grades, the college-bound had a greater increase in cigarette smoking than did their non-college-bound peers in the early to mid-1990s; but the college-bound also showed a considerably larger decline since then, leaving them with dramatically lower smoking rates at present than they had in the 1990s.

Region of the Country

The differences associated with region of the country are so sufficiently varied and complex that we cannot do justice to them here. In the past, the Northeast and West tended to have the highest proportions of students using any illicit drug, and the South, the lowest; however, these rankings have not applied to many of the specific drugs and do not apply to all grades today. The cocaine epidemic of the early 1980s was much more pronounced in the West and Northeast than in the other two regions, although the differences decreased as the overall epidemic subsided. The upsurge of ecstasy use in 1999 occurred primarily in the Northeast, but that drug's newfound popularity then spread to the three other regions of the country. While the South and West have generally had lower rates of drinking among students than the Northeast and the Midwest, those differences have narrowed somewhat in recent years and are now fairly small in all three grades. Cigarette smoking rates have generally been lowest in the West; but in 2016, after substantial declines in cigarette smoking in all three grades, the regional differences are smaller.

Population Density

There have not been very large or consistent differences in overall illicit drug use associated with population density since MTF began, helping to demonstrate just how universal the illicit drug phenomenon has been in this country. Use of any illicit drug has tended to be lowest in the more rural areas at 12th grade over most of the life of the study; and use of any illicit drug other than marijuana generally has been lower in large cities in 12th grade. Crack and heroin use have generally not been concentrated in urban areas, as is commonly believed, meaning that no parents and schools should assume that their children are immune to these threats simply because they do not live in a city. Since the late 1990s, students in non-urban areas have emerged with much higher smoking rates than others. For alcohol use there have not been large differences as a function of population density.

Socioeconomic Level

The average level of education of the student's parents, as reported by the student, is used as a proxy for socioeconomic status of the family. For many drugs the differences in use by socioeconomic class are very small, and the trends have been highly parallel. One very interesting difference occurred for cocaine, the use of which was *positively* associated with socioeconomic level in the early 1980s, meaning that higher parental education levels were associated with higher prevalence of cocaine use. However, with the advent of crack, which offered cocaine at a lower price, that association nearly disappeared by 1986.

Cigarette smoking showed a similar narrowing of class differences, but in this case a large *negative* association with socioeconomic level diminished considerably

between roughly 1985 and 1993. In more recent years, that negative association has re-emerged in the lower grades as use declined faster among students from more educated families. We believe that the removal of the Joe Camel ad campaign, which seemed to reach males from educated families in particular, may have played a role in this.

With regard to alcohol, in recent years there has been essentially no association between parental education and binge drinking among 12th graders, nor among 10th graders in 2016, but a negative correlation among 8th graders has been fairly consistent. Similarly, while binge drinking in 8th and 10th grades is negatively correlated with parental education, in 12th grade there is virtually no association.

Lessons Learned

Implications for Prevention

The wide divergence in historical trajectories of the various drugs over time helps to illustrate that, to a considerable degree, the determinants of use are often specific to each drug. These determinants include both perceived benefits and perceived adverse outcomes that young people come to associate with each drug, as well as peer norms about their use and the availability of each drug.

Unfortunately, word of the supposed benefits of using a drug usually spreads much faster than information about the adverse consequences. Supposed benefits take only rumor and a few testimonials, the spread of which have been hastened and expanded greatly by the media and in particular the Internet. It usually takes much longer for the evidence of adverse consequences (e.g., adverse reactions, death, disease, overdose, addiction) to cumulate, be recognized, and then be disseminated. Thus, when a new drug comes onto the scene, it has a considerable “honeymoon period” during which its benefits are alleged and its consequences are not yet known. We believe that ecstasy illustrated this dynamic. Synthetic marijuana and so-called “bath salts” are two more recent examples. “Vaping” may be in a honeymoon period today.

Although encouraging the avoidance or delay of *any* type of substance use is likely beneficial, especially at young ages, prevention efforts also need to be drug-specific. That is, to a considerable degree, prevention must occur drug by drug because people will not necessarily generalize the adverse consequences of the use of one drug to the use of others. Many beliefs and attitudes held by young people are drug specific. The figures in this *Overview* on perceived risk and disapproval for the various drugs—attitudes and beliefs that we have shown to be important in explaining many drug trends over the years—amply illustrate this assertion. These attitudes and beliefs are at quite different levels for the various drugs and, more importantly, often trend quite differently over time.

Marijuana is one drug that is affected by some very specific policies, including medicalization and legalization of recreational use by adults. The effects on youth behaviors and attitudes of recent changes in a number of states will need to be carefully monitored to determine their longer-term effects. Currently, marijuana does not hold the same appeal for youth as it did in the past, and today’s annual prevalence among 12th graders of 36% is considerably lower than rates exceeding 50%

observed in the 1970s. However, if states that legalize recreational marijuana allow marijuana advertising and marketing, then prevalence could rebound and approach or even surpass past levels.

“Generational Forgetting” Helps Keep the Drug Epidemic Going

Another point worth keeping in mind is that there tends to be a continuous flow of new drugs onto the scene and of older ones being rediscovered by young people. Many drugs have made a comeback years after they first fell from popularity, often because knowledge among youth of their adverse consequences faded as generational replacement took place. We call this process “generational forgetting.” Examples include LSD and methamphetamine, two drugs used widely in the 1960s that made a comeback in the 1990s after their initial popularity faded as a result of their adverse consequences becoming widely recognized during periods of high use. Heroin, cocaine, PCP, and crack are some others that have followed a similar pattern. LSD, inhalants, and ecstasy have all shown some effects of generational forgetting in recent years—that is, perceived risk has declined appreciably for those drugs, particularly among the younger students—which puts future cohorts at greater risk of having a resurgence in use. In the case of LSD, perceived risk among 8th graders has declined noticeably, and more students are saying that they are not familiar with the drug. It would appear that a resurgence in availability (which declined very sharply after about 2001, most likely due to the DEA closing a major lab in 2000) could generate another increase in use.

As for newly emerging drugs, examples include nitrite inhalants and PCP in the 1970s; crack and crystal methamphetamine in the 1980s; Rohypnol, GHB, and ecstasy in the 1990s; dextromethorphan, and salvia in the early 2000s; and “bath salts,” “synthetic marijuana,” and “vaping” more recently. The frequent introduction of new drugs (or new forms or new modes of administration of older drugs, as illustrated by crack, crystal methamphetamine, and non-injected heroin) helps keep this nation’s drug problem alive. Because of the lag times described previously, the forces of containment are always playing catch-up with the forces of encouragement and exploitation. Organized efforts to reduce the grace period experienced by new drugs would seem to be among the most promising responses for minimizing the damage they will cause. Such efforts regarding ecstasy by the National Institute on Drug Abuse and others appeared to pay off.

As for other approaches to prevention, it may be useful to emphasize that almost any new drug should be considered dangerous because such drugs are made and sold by people unconcerned with possible adverse consequences on users. Those who manufacture synthetic drugs are constantly changing the chemical formulations in order to

skirt laws prohibiting their sale, and they make no effort to assess the safety of each new formulation. As a result there are many drugs on the market with little or no information about their adverse effects, and many injuries and deaths result from their use. If young people understood this, they might be less likely to use.

TABLE 1
Trends in Lifetime Prevalence of Use of Various Drugs for Grades 8, 10, and 12 Combined
(Entries are percentages.)

	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
Any Illicit Drug ^b	30.4	29.8	32.1	35.7	38.9	42.2	43.3	42.3	41.9	41.0	40.9	39.5	37.5	36.4	35.7
Any Illicit Drug other than Marijuana ^b	19.7	19.7	21.2	22.0	23.6	24.2	24.0	23.1	22.7	22.1‡	23.2	21.1	19.8	19.3	18.6
Any Illicit Drug including Inhalants ^b	36.8	36.3	38.8	41.9	44.9	47.4	48.2	47.4	46.9	46.2	45.5	43.7	41.9	41.3	41.0
Marijuana/Hashish	22.7	21.1	23.4	27.8	31.6	35.6	37.8	36.5	36.4	35.3	35.3	34.0	32.4	31.4	30.8
Inhalants	17.0	16.9	18.2	18.6	19.4	19.1	18.6	18.1	17.5	16.4	15.3	13.6	13.4	13.7	14.1
Hallucinogens	6.1	6.3	7.0	7.7	8.9	10.0	10.2	9.5	9.0	8.5‡	9.2	7.6	6.9	6.3	5.9
LSD	5.5	5.7	6.5	6.9	8.1	8.9	9.1	8.3	7.9	7.2	6.5	5.0	3.7	3.0	2.6
Hallucinogens other than LSD	2.4	2.5	2.7	3.6	3.9	4.8	4.9	4.8	4.4	4.5‡	6.7	6.0	5.8	5.6	5.4
Ecstasy (MDMA) ^c , original	—	—	—	—	—	4.9	5.2	4.5	5.3	7.2	8.0	6.9	5.4	4.7	4.0
Revised	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Cocaine	4.6	4.0	4.1	4.5	5.1	6.0	6.6	7.0	7.2	6.5	5.9	5.7	5.3	5.5	5.5
Crack	2.0	1.9	2.0	2.5	2.8	3.2	3.4	3.8	3.8	3.5	3.2	3.2	2.9	2.9	2.8
Other cocaine	4.1	3.5	3.6	3.9	4.2	5.2	5.9	6.1	6.3	5.6	5.1	4.8	4.5	4.7	4.7
Heroin	1.1	1.3	1.3	1.6	1.9	2.1	2.1	2.2	2.2	2.1	1.7	1.7	1.5	1.5	1.5
With a needle	—	—	—	—	1.1	1.2	1.1	1.1	1.3	1.0	0.9	0.9	0.9	0.9	0.9
Without a needle	—	—	—	—	1.3	1.7	1.7	1.6	1.6	1.8	1.3	1.3	1.3	1.2	1.1
Amphetamines ^b	12.9	12.5	13.8	14.3	15.2	15.5	15.2	14.5	14.0	13.5	13.9	13.1	11.8	11.2	10.3
Methamphetamine	—	—	—	—	—	—	—	—	6.5	6.2	5.8	5.3	5.0	4.5	3.9
Tranquilizers	5.5	5.3	5.4	5.5	5.8	6.5	6.6	6.9	7.0	6.9‡	7.9	7.9	7.3	7.1	6.8
Alcohol	80.1	79.2‡	68.4	68.4	68.2	68.4	68.8	67.4	66.4	66.6	65.5	62.7	61.7	60.5	58.6
Been drunk	46.3	44.9	44.6	44.3	44.5	45.1	45.7	44.0	43.7	44.0	43.4	40.5	38.9	39.4	38.4
Flavored alcoholic beverages	—	—	—	—	—	—	—	—	—	—	—	—	—	54.7	54.7
Cigarettes	53.5	53.0	54.0	54.6	55.8	57.8	57.4	56.0	54.5	51.8	49.1	44.2	40.8	39.6	37.4
Smokeless Tobacco	—	26.2	25.6	26.3	26.0	25.7	22.7	21.1	19.4	17.9	16.6	15.2	14.1	13.6	13.8
Steroids	1.9	1.8	1.8	2.1	2.1	1.8	2.1	2.3	2.8	3.0	3.3	3.3	3.0	2.5	2.1

Table continued on next page.

TABLE 1 (continued)
Trends in Lifetime Prevalence of Use of Various Drugs for Grades 8, 10, and 12 Combined

(Entries are percentages.)

	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015–2016 change	Peak year–2016 change Absolute change	Peak year–2016 change Proportional change (%) ^a	Low year–2016 change Absolute change	Low year–2016 change Proportional change
Any Illicit Drug ^b	34.0	32.7	32.6	33.2	34.4	34.7	34.1	36.0†	34.9	34.3	<u>32.6</u>	-1.7 s	-2.3 ss	-6.5	—	—
Any Illicit Drug other than Marijuana ^b	18.2	17.7	16.8	16.5	16.8	16.1	15.5	16.8†	15.8	15.1	<u>14.3</u>	-0.8	-1.5 s	-9.4	—	—
Any Illicit Drug including Inhalants ^b	39.3	38.0	37.9	37.9	38.8	38.7	37.9	39.3†	37.9	37.4	<u>34.9</u>	-2.6 ss	-3.1 sss	-8.1	—	—
Marijuana/Hashish	28.9	<u>27.9</u>	<u>27.9</u>	29.0	30.4	31.0	30.7	32.0	30.5	30.0	28.6	-1.4 s	-9.2 sss	-24.3	+0.7	+2.6
Inhalants	13.7	13.5	13.1	12.5	12.1	10.6	10.0	8.9	8.8	7.5	<u>6.5</u>	-1.0 ss	-12.9 sss	-66.6	—	—
Hallucinogens	5.7	5.8	5.6	5.3	5.8	5.7	5.0	5.0	4.3	4.3	<u>4.3</u>	0.0	-4.9 sss	-53.6	—	—
LSD	2.5	2.6	2.7	2.5	2.8	2.7	2.5	2.6	<u>2.4</u>	2.8	3.1	+0.2	-6.0 sss	-66.3	+0.6 ss	+26.8
Hallucinogens other than LSD	5.2	5.1	4.8	4.7	5.0	4.9	4.3	4.1	3.5	3.1	<u>3.0</u>	-0.1	-3.7 sss	-55.3	—	—
Ecstasy (MDMA) ^c , original	4.3	4.5	4.1	4.6	5.5	5.5	4.6	4.7	3.5	—	—	—	—	—	—	—
Revised	—	—	—	—	—	—	—	—	5.0	4.0	<u>3.1</u>	-0.9	-2.0 sss	-39.2	—	—
Cocaine	5.3	5.2	4.8	4.2	3.8	3.4	3.3	3.1	2.9	2.7	<u>2.3</u>	-0.4	-4.8 sss	-67.5	—	—
Crack	2.6	2.5	2.2	2.0	1.9	1.6	1.5	1.5	1.3	1.3	<u>1.0</u>	-0.2 s	-2.8 sss	-73.4	—	—
Other cocaine	4.7	4.6	4.1	3.7	3.4	3.1	2.9	2.7	2.5	2.3	<u>2.1</u>	-0.3	-4.2 sss	-66.9	—	—
Heroin	1.4	1.4	1.3	1.4	1.4	1.2	1.0	1.0	0.9	0.7	<u>0.6</u>	-0.1	-1.6 sss	-73.2	—	—
With a needle	0.9	0.8	0.8	0.8	0.9	0.8	0.6	0.7	0.7	0.5	<u>0.4</u>	-0.1	-0.9 sss	-67.8	—	—
Without a needle	1.0	1.0	0.9	0.9	1.0	0.9	0.7	0.7	0.6	0.5	<u>0.4</u>	0.0	-1.3 sss	-75.4	—	—
Amphetamines ^b	10.1	9.5	8.6	8.6	8.9	8.6	8.3	10.5†	9.7	9.1	<u>8.1</u>	-0.9 ss	-1.6 sss	-16.0	—	—
Methamphetamine	3.4	2.5	2.5	2.2	2.2	1.8	1.6	1.5	1.4	1.1	<u>0.8</u>	-0.2	-5.7 sss	-87.3	—	—
Tranquilizers	7.0	6.7	6.3	6.5	6.6	6.0	5.8	5.2	5.3	<u>5.2</u>	5.5	+0.3	-2.3 sss	-29.8	+0.3	-29.8
Alcohol	57.0	56.3	55.1	54.6	53.6	51.5	50.0	48.4	46.4	45.2	<u>41.9</u>	-3.3 sss	-26.8 sss	-39.0	—	—
Been drunk	37.6	36.6	35.1	35.9	34.2	32.5	32.8	31.7	29.2	28.2	<u>26.4</u>	-1.8 ss	-19.9 sss	-43.0	—	—
Flavored alcoholic beverages	53.1	51.3	49.3	47.9	46.7	44.5	42.7	41.1	38.8	37.4	<u>33.8</u>	-3.5 sss	-20.8 sss	-38.1	—	—
Cigarettes	35.0	33.3	31.3	31.2	30.9	28.7	27.0	25.6	22.9	21.1	<u>18.2</u>	-2.9 sss	-39.5 sss	-68.4	—	—
Smokeless Tobacco	13.3	12.9	12.3	13.5	14.5	13.8	13.5	12.8	12.1	11.3	<u>10.3</u>	-1.0	-16.0 sss	-60.9	—	—
E-Vaporizers	—	—	—	—	—	—	—	—	—	29.9	<u>26.6</u>	-3.3 sss	-3.3 sss	-11.0	—	—
Steroids	2.0	1.8	1.6	1.5	1.5	1.5	1.4	1.5	1.4	1.5	<u>1.3</u>	-0.2	-2.0 sss	-60.9	—	—

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. '†' indicates a change in the question text. When a question change occurs, peak levels after that change are used to calculate the peak year to current year difference.

Values in bold equal peak levels since 1991. Values in italics equal peak level before wording change. Underlined values equal lowest level since recent peak level.

Level of significance of difference between classes: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aThe proportional change is the percent by which the most recent year deviates from the peak year [or the low year] for the drug in question. So, if a drug was at 20% prevalence in the peak year and declined to 10% prevalence in the most recent year, that would reflect a proportional decline of 50%.

^bIn 2013, for the questions on the use of amphetamines, the text was changed on two of the questionnaire forms for 8th and 10th graders and four of the questionnaire forms for 12th graders. This change also impacted the any illicit drug indices. Data presented here include only the changed forms beginning in 2013.

^cIn 2014, the text was changed on one of the questionnaire forms for 8th, 10th, and 12th graders to include "molly" in the description. The remaining forms were changed in 2015. Data for both versions of the question are presented here.

TABLE 2
Trends in Annual Prevalence of Use of Various Drugs for Grades 8, 10, and 12 Combined

(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Any Illicit Drug ^c	20.2	19.7	23.2	27.6	31.0	33.6	34.1	32.2	31.9	31.4	31.8	30.2	28.4	27.6	27.1
Any Illicit Drug other than Marijuana ^c	12.0	12.0	13.6	14.6	16.4	17.0	16.8	15.8	15.6	15.3‡	16.3	14.6	13.7	13.5	13.1
Any Illicit Drug including Inhalants ^c	23.5	23.2	26.7	31.1	34.1	36.6	36.7	35.0	34.6	34.1	34.3	32.3	30.8	30.1	30.1
Marijuana/Hashish	15.0	14.3	17.7	22.5	26.1	29.0	30.1	28.2	27.9	27.2	27.5	26.1	24.6	23.8	23.4
Synthetic marijuana	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Inhalants	7.6	7.8	8.9	9.6	10.2	9.9	9.1	8.5	7.9	7.7	6.9	6.1	6.2	6.7	7.0
Hallucinogens	3.8	4.1	4.8	5.2	6.6	7.2	6.9	6.3	6.1	5.4‡	6.0	4.5	4.1	4.0	3.9
LSD	3.4	3.8	4.3	4.7	5.9	6.3	6.0	5.3	5.3	4.5	4.1	2.4	1.6	1.6	1.5
Hallucinogens other than LSD	1.3	1.4	1.7	2.2	2.7	3.2	3.2	3.1	2.9	2.8‡	4.0	3.7	3.6	3.6	3.4
Ecstasy (MDMA) ^d , original	—	—	—	—	—	3.1	3.4	2.9	3.7	5.3	6.0	4.9	3.1	2.6	2.4
Revised	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Salvia	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Cocaine	2.2	2.1	2.3	2.8	3.3	4.0	4.3	4.5	4.5	3.9	3.5	3.7	3.3	3.5	3.5
Crack	1.0	1.1	1.2	1.5	1.8	2.0	2.1	2.4	2.2	2.1	1.8	2.0	1.8	1.7	1.6
Other cocaine	2.0	1.8	2.0	2.3	2.8	3.4	3.7	3.7	4.0	3.3	3.0	3.1	2.8	3.1	3.0
Heroin	0.5	0.6	0.6	0.9	1.2	1.3	1.3	1.2	1.3	1.3	0.9	1.0	0.8	0.9	0.8
With a needle	—	—	—	—	0.7	0.7	0.7	0.7	0.7	0.5	0.5	0.5	0.5	0.5	0.5
Without a needle	—	—	—	—	0.9	0.9	1.0	0.9	1.0	1.1	0.7	0.7	0.6	0.7	0.7
OxyContin	—	—	—	—	—	—	—	—	—	—	—	2.7	3.2	3.3	3.4
Vicodin	—	—	—	—	—	—	—	—	—	—	—	6.0	6.6	5.8	5.7
Amphetamines ^c	7.5	7.3	8.4	9.1	10.0	10.4	10.1	9.3	9.0	9.2	9.6	8.9	8.0	7.6	7.0
Ritalin	—	—	—	—	—	—	—	—	—	—	4.2	3.8	3.5	3.6	3.3
Adderall	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Methamphetamine	—	—	—	—	—	—	—	—	4.1	3.5	3.4	3.2	3.0	2.6	2.4
Bath salts (synthetic stimulants)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Tranquilizers	2.8	2.8	2.9	3.1	3.7	4.1	4.1	4.4	4.4	4.5‡	5.5	5.3	4.8	4.8	4.7
OTC Cough/Cold Medicines	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Rohypnol	—	—	—	—	—	1.1	1.1	1.1	0.8	0.7	0.9‡	0.8	0.8	0.9	0.8
GHB ^b	—	—	—	—	—	—	—	—	—	1.4	1.2	1.2	1.2	1.1	0.8
Ketamine ^b	—	—	—	—	—	—	—	—	—	2.0	1.9	2.0	1.7	1.3	1.0
Alcohol	67.4	66.3‡	59.7	60.5	60.4	60.9	61.4	59.7	59.0	59.3	58.2	55.3	54.4	54.0	51.9
Been drunk	35.8	34.3	34.3	35.0	35.9	36.7	36.9	35.5	36.0	35.9	35.0	32.1	31.2	32.5	30.8
Flavored alcoholic beverages	—	—	—	—	—	—	—	—	—	—	—	—	—	44.5	43.9
Alcoholic beverages containing caffeine	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Dissolvable tobacco products	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Snus	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids	1.2	1.1	1.0	1.2	1.3	1.1	1.2	1.3	1.7	1.9	2.0	2.0	1.7	1.6	1.3

Table continued on next page.

TABLE 2 (continued)
Trends in Annual Prevalence of Use of Various Drugs for Grades 8, 10, and 12 Combined
(Entries are percentages.)

	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015–2016 change	Peak year–2016 change		Low year–2016 change	
													Absolute change	Proportional change (%) ^a	Absolute change	Proportional change
Any Illicit Drug ^c	25.8	24.8	24.9	25.9	27.3	27.6	27.1	28.6‡	27.2	26.8	<u>25.3</u>	-1.4 s	-1.9 s	-6.9	—	—
Any Illicit Drug other than Marijuana ^c	12.7	12.4	11.9	11.6	11.8	11.3	10.8	11.4‡	10.9	10.5	<u>9.7</u>	-0.8 s	-1.2 s	-11.0	—	—
Any Illicit Drug including Inhalants ^c	28.7	27.6	27.6	28.5	29.7	29.8	29.0	30.5‡	28.5	28.4	<u>26.3</u>	-2.1 ss	-2.2 ss	-7.8	—	—
Marijuana/Hashish	22.0	<u>21.4</u>	21.5	22.9	24.5	25.0	24.7	25.8	24.2	23.7	22.6	-1.1	-7.5 sss	-24.8	+1.2	+5.8
Synthetic marijuana	—	—	—	—	—	—	8.0	6.4	4.8	4.2	<u>3.1</u>	-1.0 sss	-4.9 sss	-60.7	—	—
Inhalants	6.9	6.4	6.4	6.1	6.0	5.0	4.5	3.8	3.6	3.2	<u>2.6</u>	-0.5 s	-7.5 sss	-74.0	—	—
Hallucinogens	3.6	3.8	3.8	3.5	3.8	3.7	3.2	3.1	2.8	2.8	<u>2.8</u>	0.0	-3.2 sss	-53.5	—	—
LSD	<u>1.4</u>	1.7	1.9	1.6	1.8	1.8	1.6	1.6	1.7	1.9	2.0	+0.1	-4.4 sss	-69.0	+0.6 ss	+39.5
Hallucinogens other than LSD	3.3	3.3	3.2	3.0	3.3	3.1	2.7	2.5	2.1	1.9	<u>1.8</u>	-0.1	-2.2 sss	-55.2	—	—
Ecstasy (MDMA) ^d , original	2.7	3.0	2.9	3.0	3.8	3.7	2.5	2.8	2.2	—	—	—	—	—	—	—
Revised	—	—	—	—	—	—	—	—	3.4	2.4	<u>1.8</u>	-0.6 sss	-1.6 sss	-46.8	—	—
Salvia	—	—	—	—	3.5	3.6	2.7	2.3	1.4	1.2	<u>1.2</u>	0.0	-2.4 sss	-67.1	—	—
Cocaine	3.5	3.4	2.9	2.5	2.2	2.0	1.9	1.8	1.6	1.7	<u>1.4</u>	-0.3 s	-3.0 sss	-68.3	—	—
Crack	1.5	1.5	1.3	1.2	1.1	1.0	0.9	0.8	0.7	0.8	<u>0.6</u>	-0.2 ss	-1.8 sss	-75.6	—	—
Other cocaine	3.1	2.9	2.6	2.1	1.9	1.7	1.7	1.5	1.5	1.5	<u>1.2</u>	-0.2	-2.8 sss	-69.1	—	—
Heroin	0.8	0.8	0.8	0.8	0.8	0.7	0.6	0.6	0.5	0.4	<u>0.3</u>	-0.1 s	-1.0 sss	-77.4	—	—
With a needle	0.5	0.5	0.5	0.5	0.6	0.5	0.4	0.4	0.4	0.3	<u>0.3</u>	0.0	-0.4 sss	-62.4	—	—
Without a needle	0.6	0.7	0.6	0.5	0.6	0.5	0.4	0.4	<u>0.3</u>	0.3	<u>0.2</u>	-0.1	-0.9 sss	-82.5	—	—
OxyContin	3.5	3.5	3.4	3.9	3.8	3.4	2.9	2.9	2.4	2.3	<u>2.1</u>	-0.2	-1.8 sss	-45.9	—	—
Vicodin	6.3	6.2	6.1	6.5	5.9	5.1	4.3	3.7	3.0	2.5	<u>1.8</u>	-0.7 ss	-4.8 sss	-72.5	—	—
Amphetamines ^c	6.8	6.5	5.8	5.9	6.2	5.9	5.6	7.0‡	6.6	6.2	<u>5.4</u>	-0.8 ss	-1.2 sss	-17.9	—	—
Ritalin	3.5	2.8	2.6	2.5	2.2	2.1	1.7	1.7	1.5	1.4	<u>1.1</u>	-0.3	-3.1 sss	-74.8	—	—
Adderall	—	—	—	4.3	4.5	4.1	4.4	4.4	4.1	4.5	<u>3.9</u>	-0.6 ss	-0.5 s	-10.3	—	—
Methamphetamine	2.0	1.4	1.3	1.3	1.3	1.2	1.0	1.0	0.8	0.6	<u>0.5</u>	-0.1	-3.6 sss	-88.1	—	—
Bath salts (synthetic stimulants)	—	—	—	—	—	—	0.9	0.9	0.8	<u>0.7</u>	0.8	+0.1	-0.1	-13.3	+0.1	+18.8
Tranquilizers	4.6	4.5	4.3	4.5	4.4	3.9	3.7	<u>3.3</u>	3.4	3.4	3.5	+0.1	-2.0 sss	-36.2	+0.2	+5.7
OTC Cough/Cold Medicines	5.4	5.0	4.7	5.2	4.8	4.4	4.4	4.0	3.2	<u>3.1</u>	3.2	+0.1	-2.2 sss	-40.6	+0.1	+2.1
Rohypnol	0.7	0.8	0.7	0.6	0.8	0.9	0.7	0.6	0.5	<u>0.5</u>	0.7	+0.2 s	-0.2	-25.2	+0.2 s	+43.4
GHB ^b	0.9	0.7	0.9	0.9	<u>0.8</u>	<u>0.8</u>	—	—	—	—	—	—	—	—	—	—
Ketamine ^b	1.1	<u>1.0</u>	1.2	1.3	1.2	1.2	—	—	—	—	—	—	—	—	—	—
Alcohol	50.7	50.2	48.7	48.4	47.4	45.3	44.3	42.8	40.7	39.9	<u>36.7</u>	-3.2 sss	-24.7 sss	-40.2	—	—
Been drunk	30.7	29.7	28.1	28.7	27.1	25.9	26.4	25.4	23.6	22.5	<u>20.7</u>	-1.8 ss	-16.2 sss	-44.0	—	—
Flavored alcoholic beverages	42.4	40.8	39.0	37.8	35.9	33.7	32.5	31.3	29.4	28.8	<u>25.3</u>	-3.5 sss	-19.1 sss	-43.0	—	—
Alcoholic beverages containing caffeine	—	—	—	—	—	19.7	18.6	16.6	14.3	13.0	<u>11.2</u>	-1.8	-8.4 sss	-43.0	—	—
Dissolvable tobacco products	—	—	—	—	—	—	1.4	1.4	1.2	1.1	<u>0.9</u>	-0.2	-0.5	-34.1	—	—
Snus	—	—	—	—	—	—	5.6	4.8	4.1	3.8	<u>3.6</u>	-0.2	-2.0 sss	-36.0	—	—
Steroids	1.3	1.1	1.1	1.0	0.9	0.9	0.9	0.9	0.9	1.0	<u>0.8</u>	-0.2 ss	-1.3 sss	-62.5	—	—

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. '‡' indicates a change in the question text. When a question change occurs, peak levels after that change are used to calculate the peak year to current year difference.

Values in bold equal peak levels since 1991. Values in italics equal peak level before wording change. Underlined values equal lowest level since recent peak level.

Level of significance of difference between classes: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aThe proportional change is the percent by which the most recent year deviates from the peak year [or the low year] for the drug in question. So, if a drug was at 20% prevalence in the peak year and declined to 10% prevalence in the most recent year, that would reflect a proportional decline of 50%.

^bQuestion was discontinued among 8th and 10th graders in 2012.

^cIn 2013, for the questions on the use of amphetamines, the text was changed on two of the questionnaire forms for 8th and 10th graders and four of the questionnaire forms for 12th graders. This change also impacted the any illicit drug indices. Data presented here include only the changed forms beginning in 2013.

^dIn 2014, the text was changed on one of the questionnaire forms for 8th, 10th, and 12th graders to include "molly" in the description. The remaining forms were changed in 2015. Data for both versions of the question are presented here.

TABLE 3
Trends in 30-Day Prevalence of Use of Various Drugs for Grades 8, 10, and 12 Combined

(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Any Illicit Drug ^b	10.9	10.5	13.3	16.8	18.6	20.6	20.5	19.5	19.5	19.2	19.4	18.2	17.3	16.2	15.8
Any Illicit Drug other than Marijuana ^b	5.4	5.5	6.5	7.1	8.4	8.4	8.4	8.2	7.9	8.0‡	8.2	7.7	7.1	7.0	6.7
Any Illicit Drug including Inhalants ^b	13.0	12.5	15.4	18.9	20.7	22.4	22.2	21.1	21.1	21.0	20.8	19.5	18.6	17.5	17.5
Marijuana/Hashish	8.3	7.7	10.2	13.9	15.6	17.7	17.9	16.9	16.9	16.3	16.6	15.3	14.8	13.6	13.4
Inhalants	3.2	3.3	3.8	4.0	4.3	3.9	3.7	3.4	3.3	3.2	2.8	2.7	2.7	2.9	2.9
Hallucinogens	1.5	1.6	1.9	2.2	3.1	2.7	3.0	2.8	2.5	2.0‡	2.3	1.7	1.5	1.5	1.5
LSD	1.3	1.5	1.6	1.9	2.8	2.1	2.4	2.3	2.0	1.4	1.5	0.7	0.6	0.6	0.6
Hallucinogens other than LSD	0.5	0.5	0.7	1.0	1.0	1.2	1.2	1.2	1.1	1.1‡	1.4	1.4	1.2	1.3	1.2
Ecstasy (MDMA) ^c , original	—	—	—	—	—	1.5	1.3	1.2	1.6	2.4	2.4	1.8	1.0	0.9	0.9
Revised	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Cocaine	0.8	0.9	0.9	1.2	1.5	1.7	1.8	1.9	1.9	1.7	1.5	1.6	1.4	1.6	1.6
Crack	0.4	0.5	0.5	0.7	0.8	0.9	0.8	1.0	0.9	0.9	0.9	1.0	0.8	0.8	0.8
Other cocaine	0.7	0.7	0.8	1.1	1.2	1.3	1.5	1.6	1.7	1.4	1.3	1.3	1.2	1.4	1.3
Heroin	0.2	0.3	0.3	0.4	0.6	0.6	0.6	0.6	0.6	0.6	0.4	0.5	0.4	0.5	0.5
With a needle	—	—	—	—	0.3	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Without a needle	—	—	—	—	0.4	0.4	0.5	0.4	0.4	0.4	0.3	0.4	0.3	0.3	0.3
Amphetamines ^b	3.0	3.3	3.9	4.0	4.5	4.8	4.5	4.3	4.2	4.5	4.7	4.4	3.9	3.6	3.3
Methamphetamine	—	—	—	—	—	—	—	—	1.5	1.5	1.4	1.5	1.4	1.1	0.9
Tranquilizers	1.1	1.1	1.1	1.3	1.6	1.7	1.7	1.9	1.9	2.1‡	2.3	2.4	2.2	2.1	2.1
Alcohol	39.8	38.4‡	36.3	37.6	37.8	38.8	38.6	37.4	37.2	36.6	35.5	33.3	33.2	32.9	31.4
Been drunk	19.2	17.8	18.2	19.3	20.3	20.4	21.2	20.4	20.6	20.3	19.7	17.4	17.7	18.1	17.0
Flavored alcoholic beverages	—	—	—	—	—	—	—	—	—	—	—	—	—	23.0	21.6
Cigarettes	20.7	21.2	23.4	24.7	26.6	28.3	28.3	27.0	25.2	22.6	20.2	17.7	16.6	16.1	15.3
Smokeless Tobacco	—	9.2	9.1	9.7	9.6	8.5	8.0	7.0	6.3	5.8	6.1	5.2	5.3	5.1	5.3
E-cigarettes	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Large Cigars	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Flavored Little Cigars	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Regular Little Cigars	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids	0.6	0.6	0.6	0.7	0.6	0.5	0.7	0.7	0.9	0.9	0.9	1.0	0.9	0.9	0.7

Table continued on next page.

TABLE 3 (continued)
Trends in 30-Day Prevalence of Use of Various Drugs for Grades 8, 10, and 12 Combined

(Entries are percentages.)

	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015–2016 change	Peak year–2016 change Absolute	Proportional change (%) ^a	Low year–2016 change Absolute	Proportional change
Any Illicit Drug ^b	14.9	14.8	14.6	15.8	16.7	17.0	16.8	17.3†	16.5	15.9	<u>15.5</u>	-0.4	-1.0	-6.0	—	—
Any Illicit Drug other than Marijuana ^b	6.4	6.4	5.9	5.7	5.7	5.7	5.2	5.4†	5.4	5.1	<u>4.6</u>	-0.5 s	-0.9 ss	-15.7	—	—
Any Illicit Drug including Inhalants ^b	16.5	16.5	16.1	17.3	18.0	18.3	17.6	18.4†	17.3	16.8	<u>16.0</u>	-0.7	-1.3 s	-7.4	—	—
Marijuana/Hashish	12.5	<u>12.4</u>	12.5	13.8	14.8	15.2	15.1	15.6	14.4	14.0	13.7	-0.3	-4.2 sss	-23.4	+1.4 ss	+10.9
Inhalants	2.7	2.6	2.6	2.5	2.4	2.1	1.7	1.5	1.4	1.3	<u>1.2</u>	-0.1	-3.1 sss	-72.8	—	—
Hallucinogens	1.3	1.4	1.4	1.3	1.4	1.3	1.1	1.1	1.0	1.0	<u>1.0</u>	0.0	-1.3 sss	-57.3	—	—
LSD	0.6	0.6	0.7	<u>0.5</u>	0.7	0.7	<u>0.5</u>	0.6	0.6	0.7	0.7	0.0	-2.1 sss	-74.9	+0.1	+26.7
Hallucinogens other than LSD	1.1	1.1	1.1	1.0	1.2	1.0	0.9	0.8	0.7	0.6	<u>0.5</u>	-0.1	-0.9 sss	-62.9	—	—
Ecstasy (MDMA) ^c , original	1.0	1.1	1.2	1.2	1.5	1.4	0.8	1.0	0.8	—	—	—	—	—	—	—
Revised	—	—	—	—	—	—	—	—	1.1	0.8	<u>0.6</u>	-0.3 ss	-0.5 s	-49.1	—	—
Cocaine	1.6	1.4	1.3	1.0	0.9	0.8	0.8	0.8	0.7	0.8	<u>0.5</u>	-0.3 ss	-1.3 sss	-71.8	—	—
Crack	0.7	0.7	0.6	0.5	0.5	0.5	0.4	0.4	0.4	0.4	<u>0.3</u>	-0.1	-0.7 sss	-73.9	—	—
Other cocaine	1.4	1.1	1.1	0.8	0.8	0.7	0.7	0.6	0.6	0.7	<u>0.4</u>	-0.3 sss	-1.3 sss	-75.9	—	—
Heroin	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.2	<u>0.2</u>	0.0	-0.4 sss	-64.4	—	—
With a needle	0.3	0.3	0.3	0.2	0.3	0.3	0.2	0.2	0.3	<u>0.1</u>	0.2	0.0	-0.2 sss	-57.6	0.0	+15.6
Without a needle	0.3	0.3	<u>0.2</u>	<u>0.2</u>	0.3	0.3	<u>0.2</u>	<u>0.2</u>	<u>0.2</u>	0.2	<u>0.1</u>	-0.1	-0.4 sss	-74.8	—	—
Amphetamines ^b	3.0	3.2	2.6	2.7	2.7	2.8	2.5	3.2†	3.2	2.7	<u>2.5</u>	-0.3	-0.7 sss	-22.3	—	—
Methamphetamine	0.7	0.5	0.7	0.5	0.6	0.5	0.5	0.4	<u>0.3</u>	<u>0.3</u>	<u>0.3</u>	-0.1	-1.3 sss	-83.1	—	—
Tranquilizers	2.1	2.0	1.9	1.9	1.9	1.7	1.5	1.5	1.5	1.5	<u>1.4</u>	-0.1	-1.0 sss	-43.0	—	—
Alcohol	31.0	30.1	28.1	28.4	26.8	25.5	25.9	24.3	22.6	21.8	<u>19.8</u>	-2.0 sss	-19.0 sss	-49.0	—	—
Been drunk	17.4	16.5	14.9	15.2	14.6	13.5	14.7	13.5	11.9	11.0	<u>10.1</u>	-0.9 s	-11.1 sss	-52.4	—	—
Flavored alcoholic beverages	21.7	20.4	18.6	17.9	17.0	15.2	14.9	14.0	12.9	12.8	<u>10.9</u>	-1.9 sss	-12.2 sss	-52.7	—	—
Cigarettes	14.4	13.6	12.6	12.7	12.8	11.7	10.6	9.6	8.0	7.0	<u>5.9</u>	-1.1 sss	-22.4 sss	-79.3	—	—
Smokeless Tobacco	5.1	5.2	4.9	6.0	6.5	5.9	5.6	5.7	5.4	4.7	<u>4.1</u>	-0.5	-5.5 sss	-57.3	—	—
E-Vaporizers	—	—	—	—	—	—	—	—	—	12.8	<u>9.9</u>	-2.9 sss	-2.9 sss	-22.8	—	—
Large Cigars	—	—	—	—	—	—	—	—	3.9	4.2	<u>3.3</u>	-0.9 sss	-0.9 sss	-20.9	—	—
Flavored Little Cigars	—	—	—	—	—	—	—	—	7.4	7.1	<u>5.6</u>	-1.5 sss	-1.8 sss	-24.4	—	—
Regular Little Cigars	—	—	—	—	—	—	—	—	4.5	4.9	<u>3.6</u>	-1.3 sss	-1.3 sss	-26.9	—	—
Steroids	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.6	0.5	0.5	<u>0.4</u>	-0.1	-0.6 sss	-57.6	—	—

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. '†' indicates a change in the question text. When a question change occurs, peak levels after that change are used to calculate the peak year to current year difference.

Values in bold equal peak levels since 1991. Values in italics equal peak level before wording change. Underlined values equal lowest level since recent peak level.

Level of significance of difference between classes: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aThe proportional change is the percent by which the most recent year deviates from the peak year [or the low year] for the drug in question. So, if a drug was at 20% prevalence in the peak year and declined to 10% prevalence in the most recent year, that would reflect a proportional decline of 50%.

^bIn 2013, for the questions on the use of amphetamines, the text was changed on two of the questionnaire forms for 8th and 10th graders and four of the questionnaire forms for 12th graders. This change also impacted the any illicit drug indices. Data presented here include only the changed forms beginning in 2013.

^cIn 2014, the text was changed on one of the questionnaire forms for 8th, 10th, and 12th graders to include "molly" in the description. The remaining forms were changed in 2015. Data for both versions of the question are presented here.

TABLE 4
Trends in Daily Prevalence of Use of Selected Drugs and Heavy Use of Alcohol and Tobacco
for Grades 8, 10, and 12 Combined

(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Marijuana	0.9	0.9	1.2	2.1	2.7	3.2	3.4	3.4	3.5	3.5	3.7	3.5	3.4	3.0	2.9
Alcohol	1.7	1.6†	2.0	1.8	1.9	2.0	2.1	2.2	2.0	1.7	2.0	1.9	1.7	1.5	1.5
5+ drinks in a row in last 2 weeks	20.0	19.0	19.5	20.3	21.1	21.9	21.9	21.5	21.7	21.2	20.4	18.9	18.6	18.8	17.5
Been drunk	0.4	0.4	0.5	0.6	0.7	0.7	0.9	0.8	0.9	0.8	0.7	0.6	0.7	0.7	0.6
Cigarettes	12.4	11.9	13.5	14.0	15.5	16.8	16.9	15.4	15.0	13.4	11.6	10.2	9.3	9.0	8.0
1/2 pack+/day	6.5	6.1	6.9	7.2	7.9	8.7	8.6	7.9	7.6	6.4	5.7	4.9	4.5	4.1	3.7
Smokeless tobacco	—	3.0	2.7	2.9	2.5	2.3	2.5	2.1	1.7	1.9	2.0	1.4	1.6	1.7	1.6

Table continued on next page.

TABLE 4 (continued)
Trends in Daily Prevalence of Use of Selected Drugs and Heavy Use of Alcohol and Tobacco
for Grades 8, 10, and 12 Combined

(Entries are percentages.)

	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015–2016 change	Peak year–2016 change		Low year–2016 change	
													Absolute change	Proportional change (%) ^a	Absolute change	Proportional change
Marijuana	2.8	<u>2.7</u>	2.8	2.8	3.4	3.6	3.6	3.7	3.3	3.3	3.0	-0.3	-0.7 sss	-18.6	+0.3	+9.9
Alcohol	1.5	1.6	1.4	1.3	1.4	1.0	1.2	1.1	1.0	0.8	<u>0.7</u>	-0.2 s	-1.5 sss	-69.4	—	—
5+ drinks in a row in last 2 weeks	17.4	17.2	15.5	16.1	14.9	13.6	14.3	13.2	11.7	10.7	<u>9.4</u>	-1.4 sss	-12.6 sss	-57.3	—	—
Been drunk	0.7	0.6	0.6	0.5	0.6	0.5	0.6	0.5	0.5	0.3	<u>0.3</u>	0.0	-0.6 sss	-66.7	—	—
Cigarettes	7.6	7.1	6.4	6.4	6.4	5.7	5.2	4.7	3.6	3.2	<u>2.5</u>	-0.7 sss	-14.5 sss	-85.5	—	—
1/2 pack+/day	3.4	3.0	2.7	2.6	2.5	2.1	1.9	1.8	1.4	1.1	<u>0.9</u>	-0.3 s	-7.9 sss	-89.9	—	—
Smokeless tobacco	1.5	1.6	1.6	1.8	2.1	1.8	1.9	1.7	1.8	1.7	<u>1.4</u>	-0.3	-1.6 sss	-52.9	—	—

Source. The Monitoring the Future study, the University of Michigan.

Notes. '—' indicates data not available. '‡' indicates a change in the question text. When a question change occurs, peak levels after that change are used to calculate the peak year to current year difference.

Values in bold equal peak levels since 1991. Values in italics equal peak level before wording change. Underlined values equal lowest level since recent peak level.

Level of significance of difference between classes: s = .05, ss = .01, sss = .001.

Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aThe proportional change is the percent by which the most recent year deviates from the peak year [or the low year] for the drug in question. So, if a drug was at 20% prevalence in the peak year and declined to 10% prevalence in the most recent year, that would reflect a proportional decline of 50%.

TABLE 5
Trends in Lifetime Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
Any Illicit Drug ^a																											
8th Grade	18.7	20.6	22.5	25.7	28.5	31.2	29.4	29.0	28.3	26.8	26.8	24.5	22.8	21.5	21.4	20.9	19.0	19.6	19.9	21.4	20.1	18.5†	21.1	20.3	20.5	17.2	-3.3 sss
10th Grade	30.6	29.8	32.8	37.4	40.9	45.4	47.3	44.9	46.2	45.6	45.6	44.6	41.4	39.8	38.2	36.1	35.6	34.1	36.0	37.0	37.7	36.8†	39.1	37.4	34.7	33.7	-1.1
12th Grade	44.1	40.7	42.9	45.6	48.4	50.8	54.3	54.1	54.7	54.0	53.9	53.0	51.1	51.1	50.4	48.2	46.8	47.4	46.7	48.2	49.9	49.1†	49.8	49.1	48.9	48.3	-0.6
Any Illicit Drug other than Marijuana ^{a,b}																											
8th Grade	14.3	15.6	16.8	17.5	18.8	19.2	17.7	16.9	16.3	15.8†	17.0	13.7	13.6	12.2	12.1	12.2	11.1	11.2	10.4	10.6	9.8	8.7†	10.4	10.0	10.3	8.9	-1.4 s
10th Grade	19.1	19.2	20.9	21.7	24.3	25.5	25.0	23.6	24.0	23.1†	23.6	22.1	19.7	18.8	18.0	17.5	18.2	15.9	16.7	16.8	15.6	14.9†	16.4	15.9	14.6	14.0	-0.6
12th Grade	26.9	25.1	26.7	27.6	28.1	28.5	30.0	29.4	29.4	29.0†	30.7	29.5	27.7	28.7	27.4	26.9	25.5	24.9	24.0	24.7	24.9	24.1†	24.8	22.6	21.1	20.7	-0.4
Any Illicit Drug including Inhalants ^{a,c}																											
8th Grade	28.5	29.6	32.3	35.1	38.1	39.4	38.1	37.8	37.2	35.1	34.5	31.6	30.3	30.2	30.0	29.2	27.7	28.3	27.9	28.6	26.4	25.1†	25.9	25.2	24.9	20.6	-4.2 sss
10th Grade	36.1	36.2	38.7	42.7	45.9	49.8	50.9	49.3	49.9	49.3	48.8	47.7	44.9	43.1	42.1	40.1	39.8	38.7	40.0	40.6	40.8	40.0†	41.6	40.4	37.2	35.9	-1.3
12th Grade	47.6	44.4	46.6	49.1	51.5	53.5	56.3	56.1	56.3	57.0	56.0	54.6	52.8	53.0	53.5	51.2	49.1	49.3	48.4	49.9	51.8	50.3†	52.3	49.9	51.4	49.3	-2.2
Marijuana/Hashish																											
8th Grade	10.2	11.2	12.6	16.7	19.9	23.1	22.6	22.2	22.0	20.3	20.4	19.2	17.5	16.3	16.5	15.7	14.2	14.6	15.7	17.3	16.4	15.2	16.5	15.6	15.5	12.8	-2.6 ss
10th Grade	23.4	21.4	24.4	30.4	34.1	39.8	42.3	39.6	40.9	40.3	40.1	38.7	36.4	35.1	34.1	31.8	31.0	29.9	32.3	33.4	34.5	33.8	35.8	33.7	31.1	29.7	-1.3
12th Grade	36.7	32.6	35.3	38.2	41.7	44.9	49.6	49.1	49.7	48.8	49.0	47.8	46.1	45.7	44.8	42.3	41.8	42.6	42.0	43.8	45.5	45.2	45.5	44.4	44.7	44.5	-0.1
Inhalants ^{c,d}																											
8th Grade	17.6	17.4	19.4	19.9	21.6	21.2	21.0	20.5	19.7	17.9	17.1	15.2	15.8	17.3	17.1	16.1	15.6	15.7	14.9	14.5	13.1	11.8	10.8	10.8	9.4	7.7	-1.7 ss
10th Grade	15.7	16.6	17.5	18.0	19.0	19.3	18.3	18.3	17.0	16.6	15.2	13.5	12.7	12.4	13.1	13.3	13.6	12.8	12.3	12.0	10.1	9.9	8.7	8.7	7.2	6.6	-0.6
12th Grade	17.6	16.6	17.4	17.7	17.4	16.6	16.1	15.2	15.4	14.2	13.0	11.7	11.2	10.9	11.4	11.1	10.5	9.9	9.5	9.0	8.1	7.9	6.9	6.5	5.7	5.0	-0.7
Hallucinogens ^{b,f}																											
8th Grade	3.2	3.8	3.9	4.3	5.2	5.9	5.4	4.9	4.8	4.6†	5.2	4.1	4.0	3.5	3.8	3.4	3.1	3.3	3.0	3.4	3.3	2.8	2.5	2.0	2.0	1.9	-0.1
10th Grade	6.1	6.4	6.8	8.1	9.3	10.5	10.5	9.8	9.7	8.9†	8.9	7.8	6.9	6.4	5.8	6.1	6.4	5.5	6.1	6.1	6.0	5.2	5.4	5.0	4.6	4.4	-0.2
12th Grade	9.6	9.2	10.9	11.4	12.7	14.0	15.1	14.1	13.7	13.0†	14.7	12.0	10.6	9.7	8.8	8.3	8.4	8.7	7.4	8.6	8.3	7.5	7.6	6.3	6.4	6.7	+0.4

(Table continued on next page.)

TABLE 5 (cont.)
Trends in Lifetime Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
LSD ^b																											
8th Grade	2.7	3.2	3.5	3.7	4.4	5.1	4.7	4.1	4.1	3.9	3.4	2.5	2.1	1.8	1.9	1.6	1.6	1.9	1.7	1.8	1.7	1.3	1.4	1.1	1.3	1.2	0.0
10th Grade	5.6	5.8	6.2	7.2	8.4	9.4	9.5	8.5	8.5	7.6	6.3	5.0	3.5	2.8	2.5	2.7	3.0	2.6	3.0	3.0	2.8	2.6	2.7	2.6	3.0	3.2	+0.2
12th Grade	8.8	8.6	10.3	10.5	11.7	12.6	13.6	12.6	12.2	11.1	10.9	8.4	5.9	4.6	3.5	3.3	3.4	4.0	3.1	4.0	4.0	3.8	3.9	3.7	4.3	4.9	+0.6
Hallucinogens other than LSD ^b																											
8th Grade	1.4	1.7	1.7	2.2	2.5	3.0	2.6	2.5	2.4	2.3‡	3.9	3.3	3.2	3.0	3.3	2.8	2.6	2.5	2.4	2.7	2.8	2.3	1.9	1.5	1.2	1.3	0.0
10th Grade	2.2	2.5	2.8	3.8	3.9	4.7	4.8	5.0	4.7	4.8‡	6.6	6.3	5.9	5.8	5.2	5.5	5.7	4.8	5.4	5.3	5.2	4.5	4.4	4.1	3.3	3.1	-0.3
12th Grade	3.7	3.3	3.9	4.9	5.4	6.8	7.5	7.1	6.7	6.9‡	10.4	9.2	9.0	8.7	8.1	7.8	7.7	7.8	6.8	7.7	7.3	6.6	6.4	5.1	4.8	4.7	-0.1
Ecstasy (MDMA) ^g																											
8th Grade, original	—	—	—	—	—	3.4	3.2	2.7	2.7	4.3	5.2	4.3	3.2	2.8	2.8	2.5	2.3	2.4	2.2	3.3	2.6	2.0	1.8	1.4	—	—	—
Revised	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.4	2.3	1.7	-0.6 s
10th Grade, original	—	—	—	—	—	5.6	5.7	5.1	6.0	7.3	8.0	6.6	5.4	4.3	4.0	4.5	5.2	4.3	5.5	6.4	6.6	5.0	5.7	3.7	—	—	—
Revised	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.2	3.8	2.8	-1.1 sss
12th Grade, original	—	—	—	—	—	6.1	6.9	5.8	8.0	11.0	11.7	10.5	8.3	7.5	5.4	6.5	6.5	6.2	6.5	7.3	8.0	7.2	7.1	5.6	—	—	—
Revised	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	7.9	5.9	4.9	-1.0 s
Cocaine																											
8th Grade	2.3	2.9	2.9	3.6	4.2	4.5	4.4	4.6	4.7	4.5	4.3	3.6	3.6	3.4	3.7	3.4	3.1	3.0	2.6	2.6	2.2	1.9	1.7	1.8	1.6	1.4	-0.2
10th Grade	4.1	3.3	3.6	4.3	5.0	6.5	7.1	7.2	7.7	6.9	5.7	6.1	5.1	5.4	5.2	4.8	5.3	4.5	4.6	3.7	3.3	3.3	3.3	2.6	2.7	2.1	-0.6 s
12th Grade	7.8	6.1	6.1	5.9	6.0	7.1	8.7	9.3	9.8	8.6	8.2	7.8	7.7	8.1	8.0	8.5	7.8	7.2	6.0	5.5	5.2	4.9	4.5	4.6	4.0	3.7	-0.4
Crack																											
8th Grade	1.3	1.6	1.7	2.4	2.7	2.9	2.7	3.2	3.1	3.1	3.0	2.5	2.5	2.4	2.4	2.3	2.1	2.0	1.7	1.5	1.5	1.0	1.2	1.2	1.0	0.9	-0.1
10th Grade	1.7	1.5	1.8	2.1	2.8	3.3	3.6	3.9	4.0	3.7	3.1	3.6	2.7	2.6	2.5	2.2	2.3	2.0	2.1	1.8	1.6	1.4	1.5	1.0	1.1	0.8	-0.3
12th Grade	3.1	2.6	2.6	3.0	3.0	3.3	3.9	4.4	4.6	3.9	3.7	3.8	3.6	3.9	3.5	3.5	3.2	2.8	2.4	2.4	1.9	2.1	1.8	1.8	1.7	1.4	-0.4
Other Cocaine ^h																											
8th Grade	2.0	2.4	2.4	3.0	3.4	3.8	3.5	3.7	3.8	3.5	3.3	2.8	2.7	2.6	2.9	2.7	2.6	2.4	2.1	2.1	1.8	1.6	1.4	1.4	1.3	1.1	-0.2
10th Grade	3.8	3.0	3.3	3.8	4.4	5.5	6.1	6.4	6.8	6.0	5.0	5.2	4.5	4.8	4.6	4.3	4.8	4.0	4.1	3.4	3.0	3.0	2.9	2.2	2.3	1.9	-0.4
12th Grade	7.0	5.3	5.4	5.2	5.1	6.4	8.2	8.4	8.8	7.7	7.4	7.0	6.7	7.3	7.1	7.9	6.8	6.5	5.3	5.1	4.9	4.4	4.2	4.1	3.4	3.3	-0.1

(Table continued on next page.)

TABLE 5 (cont.)
Trends in Lifetime Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
Heroin ^{l,j}																											
8th Grade	1.2	1.4	1.4	2.0	2.3	2.4	2.1	2.3	2.3	1.9	1.7	1.6	1.6	1.6	1.5	1.4	1.3	1.4	1.3	1.3	1.2	0.8	1.0	0.9	0.5	0.5	0.0
10th Grade	1.2	1.2	1.3	1.5	1.7	2.1	2.1	2.3	2.3	2.2	1.7	1.8	1.5	1.5	1.5	1.4	1.5	1.2	1.5	1.3	1.2	1.1	1.0	0.9	0.7	0.6	-0.1
12th Grade	0.9	1.2	1.1	1.2	1.6	1.8	2.1	2.0	2.0	2.4	1.8	1.7	1.5	1.5	1.5	1.4	1.5	1.3	1.2	1.6	1.4	1.1	1.0	1.0	0.8	0.7	-0.1
With a Needle ^j																											
8th Grade	—	—	—	—	1.5	1.6	1.3	1.4	1.6	1.1	1.2	1.0	1.0	1.1	1.0	1.0	0.9	0.9	0.9	0.9	0.8	0.6	0.6	0.8	0.3	0.3	0.0
10th Grade	—	—	—	—	1.0	1.1	1.1	1.2	1.3	1.0	0.8	1.0	0.9	0.8	0.8	0.9	0.9	0.7	0.9	0.8	0.8	0.7	0.7	0.6	0.5	0.5	0.0
12th Grade	—	—	—	—	0.7	0.8	0.9	0.8	0.9	0.8	0.7	0.8	0.7	0.7	0.9	0.8	0.7	0.7	0.6	1.1	0.9	0.7	0.7	0.8	0.6	0.5	-0.2
Without a Needle ^j																											
8th Grade	—	—	—	—	1.5	1.6	1.4	1.5	1.4	1.3	1.1	1.0	1.1	1.0	0.9	0.9	0.7	0.9	0.8	0.7	0.7	0.5	0.5	0.4	0.3	0.4	+0.1
10th Grade	—	—	—	—	1.1	1.7	1.7	1.7	1.6	1.7	1.3	1.3	1.0	1.1	1.1	1.0	1.1	0.8	1.0	0.9	0.8	0.8	0.7	0.5	0.4	0.3	-0.1
12th Grade	—	—	—	—	1.4	1.7	2.1	1.6	1.8	2.4	1.5	1.6	1.8	1.4	1.3	1.1	1.4	1.1	0.9	1.4	1.3	0.8	0.9	0.7	0.7	0.6	-0.1
Narcotics other than Heroin ^{k,l}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	6.6	6.1	6.4	6.6	7.2	8.2	9.7	9.8	10.2	10.6	9.9‡	13.5	13.2	13.5	12.8	13.4	13.1	13.2	13.2	13.0	13.0	12.2	11.1	9.5	8.4	7.8	-0.6
Amphetamines ^{k,m}																											
8th Grade	10.5	10.8	11.8	12.3	13.1	13.5	12.3	11.3	10.7	9.9	10.2	8.7	8.4	7.5	7.4	7.3	6.5	6.8	6.0	5.7	5.2	4.5‡	6.9	6.7	6.8	5.7	-1.1 ss
10th Grade	13.2	13.1	14.9	15.1	17.4	17.7	17.0	16.0	15.7	15.7	16.0	14.9	13.1	11.9	11.1	11.2	11.1	9.0	10.3	10.6	9.0	8.9‡	11.2	10.6	9.7	8.8	-0.9
12th Grade	15.4	13.9	15.1	15.7	15.3	15.3	16.5	16.4	16.3	15.6	16.2	16.8	14.4	15.0	13.1	12.4	11.4	10.5	9.9	11.1	12.2	12.0‡	13.8	12.1	10.8	10.0	-0.8
Methamphetamine ^{n,o}																											
8th Grade	—	—	—	—	—	—	—	—	4.5	4.2	4.4	3.5	3.9	2.5	3.1	2.7	1.8	2.3	1.6	1.8	1.3	1.3	1.4	1.0	0.8	0.6	-0.2
10th Grade	—	—	—	—	—	—	—	—	7.3	6.9	6.4	6.1	5.2	5.3	4.1	3.2	2.8	2.4	2.8	2.5	2.1	1.8	1.6	1.4	1.3	0.7	-0.6 ss
12th Grade	—	—	—	—	—	—	—	—	8.2	7.9	6.9	6.7	6.2	6.2	4.5	4.4	3.0	2.8	2.4	2.3	2.1	1.7	1.5	1.9	1.0	1.2	+0.1

(Table continued on next page.)

TABLE 5 (cont.)
Trends in Lifetime Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
Crystal Methamphetamine (Ice) ^o																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	3.3	2.9	3.1	3.4	3.9	4.4	4.4	5.3	4.8	4.0	4.1	4.7	3.9	4.0	4.0	3.4	3.4	2.8	2.1	1.8	2.1	1.7	2.0	1.3	1.2	1.4	+0.2
Sedatives (Barbiturates) ^{k,p}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	6.2	5.5	6.3	7.0	7.4	7.6	8.1	8.7	8.9	9.2	8.7	9.5	8.8	9.9	10.5	10.2	9.3	8.5	8.2	7.5	7.0	6.9	7.5	6.8	5.9	5.2	-0.7
Tranquilizers ^{b,k}																											
8th Grade	3.8	4.1	4.4	4.6	4.5	5.3	4.8	4.6	4.4	4.4±	5.0	4.3	4.4	4.0	4.1	4.3	3.9	3.9	3.9	4.4	3.4	3.0	2.9	2.9	3.0	3.0	0.0
10th Grade	5.8	5.9	5.7	5.4	6.0	7.1	7.3	7.8	7.9	8.0±	9.2	8.8	7.8	7.3	7.1	7.2	7.4	6.8	7.0	7.3	6.8	6.3	5.5	5.8	5.8	6.1	+0.3
12th Grade	7.2	6.0	6.4	6.6	7.1	7.2	7.8	8.5	9.3	8.9±	10.3	11.4	10.2	10.6	9.9	10.3	9.5	8.9	9.3	8.5	8.7	8.5	7.7	7.4	6.9	7.6	+0.7
Any Prescription Drug ^q																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	24.0	23.9	22.2	21.5	20.9	21.6	21.7	21.2±	22.2	19.9	18.3	18.0	-0.3
Rohypnol ^r																											
8th Grade	—	—	—	—	—	1.5	1.1	1.4	1.3	1.0	1.1	0.8	1.0	1.0	1.1	1.0	1.0	0.7	0.7	0.9	2.0	1.0	0.7	0.6	0.8	0.9	+0.1
10th Grade	—	—	—	—	—	1.5	1.7	2.0	1.8	1.3	1.5	1.3	1.0	1.2	1.0	0.8	1.3	0.9	0.7	1.4	1.2	0.8	1.1	1.0	0.5	1.0	+0.5
12th Grade	—	—	—	—	—	1.2	1.8	3.0	2.0	1.5	1.7	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Alcohol ^s																											
Any Use																											
8th Grade	70.1	69.3±	55.7	55.8	54.5	55.3	53.8	52.5	52.1	51.7	50.5	47.0	45.6	43.9	41.0	40.5	38.9	38.9	36.6	35.8	33.1	29.5	27.8	26.8	26.1	22.8	-3.3 ss
10th Grade	83.8	82.3±	71.6	71.1	70.5	71.8	72.0	69.8	70.6	71.4	70.1	66.9	66.0	64.2	63.2	61.5	61.7	58.3	59.1	58.2	56.0	54.0	52.1	49.3	47.1	43.4	-3.7 ss
12th Grade	88.0	87.5±	80.0	80.4	80.7	79.2	81.7	81.4	80.0	80.3	79.7	78.4	76.6	76.8	75.1	72.7	72.2	71.9	72.3	71.0	70.0	69.4	68.2	66.0	64.0	61.2	-2.8

(Table continued on next page.)

TABLE 5 (cont.)
Trends in Lifetime Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
Been Drunk ^o																											
8th Grade	26.7	26.8	26.4	25.9	25.3	26.8	25.2	24.8	24.8	25.1	23.4	21.3	20.3	19.9	19.5	19.5	17.9	18.0	17.4	16.3	14.8	12.8	12.2	10.8	10.9	8.6	-2.3 sss
10th Grade	50.0	47.7	47.9	47.2	46.9	48.5	49.4	46.7	48.9	49.3	48.2	44.0	42.4	42.3	42.1	41.4	41.2	37.2	38.6	36.9	35.9	34.6	33.5	30.2	28.6	26.0	-2.5 s
12th Grade	65.4	63.4	62.5	62.9	63.2	61.8	64.2	62.4	62.3	62.3	63.9	61.6	58.1	60.3	57.5	56.4	55.1	54.7	56.5	54.1	51.0	54.2	52.3	49.8	46.7	46.3	-0.5
Flavored Alcoholic Beverages ^{e,n}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	37.9	35.5	35.5	34.0	32.8	29.4	30.0	27.0	23.5	21.9	19.2	19.3	16.3	-3.0 s
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	58.6	58.8	58.1	55.7	53.5	51.4	51.3	48.4	46.7	44.9	42.3	38.7	33.3	-5.4 ss
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	71.0	73.6	69.9	68.4	65.5	67.4	62.6	62.4	60.5	58.9	57.5	55.6	53.6	-2.0
Cigarettes																											
Any Use																											
8th Grade	44.0	45.2	45.3	46.1	46.4	49.2	47.3	45.7	44.1	40.5	36.6	31.4	28.4	27.9	25.9	24.6	22.1	20.5	20.1	20.0	18.4	15.5	14.8	13.5	13.3	9.8	-3.5 sss
10th Grade	55.1	53.5	56.3	56.9	57.6	61.2	60.2	57.7	57.6	55.1	52.8	47.4	43.0	40.7	38.9	36.1	34.6	31.7	32.7	33.0	30.4	27.7	25.7	22.6	19.9	17.5	-2.4 s
12th Grade	63.1	61.8	61.9	62.0	64.2	63.5	65.4	65.3	64.6	62.5	61.0	57.2	53.7	52.8	50.0	47.1	46.2	44.7	43.6	42.2	40.0	39.5	38.1	34.4	31.1	28.3	-2.8 s
Smokeless Tobacco ^t																											
8th Grade	22.2	20.7	18.7	19.9	20.0	20.4	16.8	15.0	14.4	12.8	11.7	11.2	11.3	11.0	10.1	10.2	9.1	9.8	9.6	9.9	9.7	8.1	7.9	8.0	8.6	6.9	-1.7 s
10th Grade	28.2	26.6	28.1	29.2	27.6	27.4	26.3	22.7	20.4	19.1	19.5	16.9	14.6	13.8	14.5	15.0	15.1	12.2	15.2	16.8	15.6	15.4	14.0	13.6	12.3	10.2	-2.2 s
12th Grade	—	32.4	31.0	30.7	30.9	29.8	25.3	26.2	23.4	23.1	19.7	18.3	17.0	16.7	17.5	15.2	15.1	15.6	16.3	17.6	16.9	17.4	17.2	15.1	13.2	14.2	+1.0
Electronic Vaporizers ^{bb}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	21.7	17.5	-4.2 sss
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	32.8	29.0	-3.7 ss
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	35.5	33.8	-1.7
Steroids ^{k,u}																											
8th Grade	1.9	1.7	1.6	2.0	2.0	1.8	1.8	2.3	2.7	3.0	2.8	2.5	2.5	1.9	1.7	1.6	1.5	1.4	1.3	1.1	1.2	1.2	1.1	1.0	1.0	0.9	-0.1
10th Grade	1.8	1.7	1.7	1.8	2.0	1.8	2.0	2.0	2.7	3.5	3.5	3.5	3.0	2.4	2.0	1.8	1.8	1.4	1.3	1.6	1.4	1.3	1.3	1.4	1.2	1.3	+0.1
12th Grade	2.1	2.1	2.0	2.4	2.3	1.9	2.4	2.7	2.9	2.5	3.7	4.0	3.5	3.4	2.6	2.7	2.2	2.2	2.2	2.0	1.8	1.8	2.1	1.9	2.3	1.6	-0.7 s

(Table continued on next page.)

TABLE 5 (cont.)
Trends in Lifetime Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
Previously surveyed drugs that have been dropped.																											
Nitrites ^e																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.6	1.5	1.4	1.7	1.5	1.8	2.0	2.7	1.7	0.8	1.9	1.5	1.6	1.3	1.1	1.2	1.2	0.6	1.1	—	—	—	—	—	—	—	—
PCP ^e																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	2.9	2.4	2.9	2.8	2.7	4.0	3.9	3.9	3.4	3.4	3.5	3.1	2.5	1.6	2.4	2.2	2.1	1.8	1.7	1.8	2.3	1.6	1.3	—	—	—	—
Methaqualone ^{e,k}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.3	1.6	0.8	1.4	1.2	2.0	1.7	1.6	1.8	0.8	1.1	1.5	1.0	1.3	1.3	1.2	1.0	0.8	0.7	0.4	0.6	0.8	—	—	—	—	—

Source. The Monitoring the Future study, the University of Michigan.

Note: See footnotes following Table 8.

TABLE 6
Trends in Annual Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
Any Illicit Drug ^a																											
8th Grade	11.3	12.9	15.1	18.5	21.4	23.6	22.1	21.0	20.5	19.5	19.5	17.7	16.1	15.2	15.5	14.8	13.2	14.1	14.5	16.0	14.7	13.4†	15.2	14.6	14.8	12.0	-2.9 sss
10th Grade	21.4	20.4	24.7	30.0	33.3	37.5	38.5	35.0	35.9	36.4	37.2	34.8	32.0	31.1	29.8	28.7	28.1	26.9	29.4	30.2	31.1	30.1†	32.1	29.9	27.9	26.8	-1.1
12th Grade	29.4	27.1	31.0	35.8	39.0	40.2	42.4	41.4	42.1	40.9	41.4	41.0	39.3	38.8	38.4	36.5	35.9	36.6	36.5	38.3	40.0	39.7†	40.1	38.7	38.6	38.3	-0.3
Any Illicit Drug other than Marijuana ^{a,b}																											
8th Grade	8.4	9.3	10.4	11.3	12.6	13.1	11.8	11.0	10.5	10.2†	10.8	8.8	8.8	7.9	8.1	7.7	7.0	7.4	7.0	7.1	6.4	5.5†	6.3	6.4	6.3	5.4	-0.9 s
10th Grade	12.2	12.3	13.9	15.2	17.5	18.4	18.2	16.6	16.7	16.7†	17.9	15.7	13.8	13.5	12.9	12.7	13.1	11.3	12.2	12.1	11.2	10.8†	11.2	11.2	10.5	9.8	-0.7
12th Grade	16.2	14.9	17.1	18.0	19.4	19.8	20.7	20.2	20.7	20.4†	21.6	20.9	19.8	20.5	19.7	19.2	18.5	18.3	17.0	17.3	17.6	17.0†	17.8	15.9	15.2	14.3	-0.9
Any Illicit Drug including Inhalants ^{a,c}																											
8th Grade	16.7	18.2	21.1	24.2	27.1	28.7	27.2	26.2	25.3	24.0	23.9	21.4	20.4	20.2	20.4	19.7	18.0	19.0	18.8	20.3	18.2	17.0†	17.6	16.8	17.0	13.5	-3.5 sss
10th Grade	23.9	23.5	27.4	32.5	35.6	39.6	40.3	37.1	37.7	38.0	38.7	36.1	33.5	32.9	31.7	30.7	30.2	28.8	31.2	31.8	32.5	31.5†	33.2	31.0	28.9	27.7	-1.2
12th Grade	31.2	28.8	32.5	37.6	40.2	41.9	43.3	42.4	42.8	42.5	42.6	42.1	40.5	39.1	40.3	38.0	37.0	37.3	37.6	39.2	41.5	40.2†	42.3	39.2	40.2	38.7	-1.5
Marijuana/Hashish																											
8th Grade	6.2	7.2	9.2	13.0	15.8	18.3	17.7	16.9	16.5	15.6	15.4	14.6	12.8	11.8	12.2	11.7	10.3	10.9	11.8	13.7	12.5	11.4	12.7	11.7	11.8	9.4	-2.4 sss
10th Grade	16.5	15.2	19.2	25.2	28.7	33.6	34.8	31.1	32.1	32.2	32.7	30.3	28.2	27.5	26.6	25.2	24.6	23.9	26.7	27.5	28.8	28.0	29.8	27.3	25.4	23.9	-1.5
12th Grade	23.9	21.9	26.0	30.7	34.7	35.8	38.5	37.5	37.8	36.5	37.0	36.2	34.9	34.3	33.6	31.5	31.7	32.4	32.8	34.8	36.4	36.4	36.4	35.1	34.9	35.6	+0.7
Synthetic Marijuana ^{n,o}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	4.4	4.0	3.3	3.1	2.7	-0.5
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	8.8	7.4	5.4	4.3	3.3	-1.0 s
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	11.4	11.3	7.9	5.8	5.2	3.5	-1.7 ss
Inhalants ^{c,d}																											
8th Grade	9.0	9.5	11.0	11.7	12.8	12.2	11.8	11.1	10.3	9.4	9.1	7.7	8.7	9.6	9.5	9.1	8.3	8.9	8.1	8.1	7.0	6.2	5.2	5.3	4.6	3.8	-0.8 s
10th Grade	7.1	7.5	8.4	9.1	9.6	9.5	8.7	8.0	7.2	7.3	6.6	5.8	5.4	5.9	6.0	6.5	6.6	5.9	6.1	5.7	4.5	4.1	3.5	3.3	2.9	2.4	-0.4
12th Grade	6.6	6.2	7.0	7.7	8.0	7.6	6.7	6.2	5.6	5.9	4.5	4.5	3.9	4.2	5.0	4.5	3.7	3.8	3.4	3.6	3.2	2.9	2.5	1.9	1.9	1.7	-0.2

(Table continued on next page.)

TABLE 6 (cont.)
Trends in Annual Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
Hallucinogens ^{b,f}																											
8th Grade	1.9	2.5	2.6	2.7	3.6	4.1	3.7	3.4	2.9	2.8†	3.4	2.6	2.6	2.2	2.4	2.1	1.9	2.1	1.9	2.2	2.2	1.6	1.6	1.3	1.3	1.2	-0.1
10th Grade	4.0	4.3	4.7	5.8	7.2	7.8	7.6	6.9	6.9	6.1†	6.2	4.7	4.1	4.1	4.0	4.1	4.4	3.9	4.1	4.2	4.1	3.5	3.4	3.3	3.1	2.9	-0.1
12th Grade	5.8	5.9	7.4	7.6	9.3	10.1	9.8	9.0	9.4	8.1†	9.1	6.6	5.9	6.2	5.5	4.9	5.4	5.9	4.7	5.5	5.2	4.8	4.5	4.0	4.2	4.3	+0.2
LSD ^b																											
8th Grade	1.7	2.1	2.3	2.4	3.2	3.5	3.2	2.8	2.4	2.4	2.2	1.5	1.3	1.1	1.2	0.9	1.1	1.3	1.1	1.2	1.1	0.8	1.0	0.7	0.9	0.8	-0.1
10th Grade	3.7	4.0	4.2	5.2	6.5	6.9	6.7	5.9	6.0	5.1	4.1	2.6	1.7	1.6	1.5	1.7	1.9	1.8	1.9	1.9	1.8	1.7	1.7	1.9	2.0	2.1	+0.2
12th Grade	5.2	5.6	6.8	6.9	8.4	8.8	8.4	7.6	8.1	6.6	6.6	3.5	1.9	2.2	1.8	1.7	2.1	2.7	1.9	2.6	2.7	2.4	2.2	2.5	2.9	3.0	+0.1
Hallucinogens other than LSD ^b																											
8th Grade	0.7	1.1	1.0	1.3	1.7	2.0	1.8	1.6	1.5	1.4†	2.4	2.1	2.1	1.9	2.0	1.8	1.6	1.6	1.5	1.8	1.8	1.3	1.2	1.0	0.8	0.8	0.0
10th Grade	1.3	1.4	1.9	2.4	2.8	3.3	3.3	3.4	3.2	3.1†	4.3	4.0	3.6	3.7	3.5	3.7	3.8	3.3	3.5	3.5	3.5	3.0	2.7	2.6	1.9	2.0	0.0
12th Grade	2.0	1.7	2.2	3.1	3.8	4.4	4.6	4.6	4.3	4.4†	5.9	5.4	5.4	5.6	5.0	4.6	4.8	5.0	4.2	4.8	4.3	4.0	3.7	3.0	2.9	2.7	-0.2
PCP ^e																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.4	1.4	1.4	1.6	1.8	2.6	2.3	2.1	1.8	2.3	1.8	1.1	1.3	0.7	1.3	0.7	0.9	1.1	1.0	1.0	1.3	0.9	0.7	0.8	1.4	1.3	-0.1
Ecstasy (MDMA) ^g																											
8th Grade, original	—	—	—	—	2.3	2.3	1.8	1.7	3.1	3.5	2.9	2.1	1.7	1.7	1.4	1.5	1.7	1.3	2.4	1.7	1.1	1.1	0.9	—	—	—	—
Revised	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.5	1.4	1.0	-0.4	s
10th Grade, original	—	—	—	—	4.6	3.9	3.3	4.4	5.4	6.2	4.9	3.0	2.4	2.6	2.8	3.5	2.9	3.7	4.7	4.5	3.0	3.6	2.3	—	—	—	—
Revised	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.8	2.4	1.8	-0.6	ss
12th Grade, original	—	—	—	—	4.6	4.0	3.6	5.6	8.2	9.2	7.4	4.5	4.0	3.0	4.1	4.5	4.3	4.3	4.5	5.3	3.8	4.0	3.6	—	—	—	—
Revised	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.0	3.6	2.7	-0.9	s
Salvia ^{n,o}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.7	1.6	1.4	1.2	0.6	0.7	0.9	+0.3
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.7	3.9	2.5	2.3	1.8	1.2	0.9	-0.3
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.7	5.5	5.9	4.4	3.4	1.8	1.9	1.8	-0.1

(Table continued on next page.)

TABLE 6 (cont.)
Trends in Annual Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
Cocaine																											
8th Grade	1.1	1.5	1.7	2.1	2.6	3.0	2.8	3.1	2.7	2.6	2.5	2.3	2.2	2.0	2.2	2.0	2.0	1.8	1.6	1.6	1.4	1.2	1.0	1.0	0.9	0.8	-0.1
10th Grade	2.2	1.9	2.1	2.8	3.5	4.2	4.7	4.7	4.9	4.4	3.6	4.0	3.3	3.7	3.5	3.2	3.4	3.0	2.7	2.2	1.9	2.0	1.9	1.5	1.8	1.3	-0.5 ss
12th Grade	3.5	3.1	3.3	3.6	4.0	4.9	5.5	5.7	6.2	5.0	4.8	5.0	4.8	5.3	5.1	5.7	5.2	4.4	3.4	2.9	2.9	2.7	2.6	2.6	2.5	2.3	-0.3
Crack																											
8th Grade	0.7	0.9	1.0	1.3	1.6	1.8	1.7	2.1	1.8	1.8	1.7	1.6	1.6	1.3	1.4	1.3	1.3	1.1	1.1	1.0	0.9	0.6	0.6	0.7	0.5	0.5	0.0
10th Grade	0.9	0.9	1.1	1.4	1.8	2.1	2.2	2.5	2.4	2.2	1.8	2.3	1.6	1.7	1.7	1.3	1.3	1.3	1.2	1.0	0.9	0.8	0.8	0.5	0.7	0.4	-0.3 s
12th Grade	1.5	1.5	1.5	1.9	2.1	2.1	2.4	2.5	2.7	2.2	2.1	2.3	2.2	2.3	1.9	2.1	1.9	1.6	1.3	1.4	1.0	1.2	1.1	1.1	1.1	0.8	-0.3
Other Cocaine ^h																											
8th Grade	1.0	1.2	1.3	1.7	2.1	2.5	2.2	2.4	2.3	1.9	1.9	1.8	1.6	1.6	1.7	1.6	1.5	1.4	1.3	1.3	1.1	1.0	0.8	0.8	0.8	0.6	-0.2
10th Grade	2.1	1.7	1.8	2.4	3.0	3.5	4.1	4.0	4.4	3.8	3.0	3.4	2.8	3.3	3.0	2.9	3.1	2.6	2.3	1.9	1.7	1.8	1.6	1.3	1.5	1.1	-0.4 s
12th Grade	3.2	2.6	2.9	3.0	3.4	4.2	5.0	4.9	5.8	4.5	4.4	4.4	4.2	4.7	4.5	5.2	4.5	4.0	3.0	2.6	2.6	2.4	2.4	2.4	2.1	2.0	-0.1
Heroin ^{ij}																											
8th Grade	0.7	0.7	0.7	1.2	1.4	1.6	1.3	1.3	1.4	1.1	1.0	0.9	0.9	1.0	0.8	0.8	0.8	0.9	0.7	0.8	0.7	0.5	0.5	0.5	0.3	0.3	0.0
10th Grade	0.5	0.6	0.7	0.9	1.1	1.2	1.4	1.4	1.4	1.4	0.9	1.1	0.7	0.9	0.9	0.9	0.8	0.8	0.9	0.8	0.8	0.6	0.6	0.5	0.5	0.3	-0.1
12th Grade	0.4	0.6	0.5	0.6	1.1	1.0	1.2	1.0	1.1	1.5	0.9	1.0	0.8	0.9	0.8	0.8	0.9	0.7	0.7	0.9	0.8	0.6	0.6	0.6	0.5	0.3	-0.2
With a Needle ^j																											
8th Grade	—	—	—	—	0.9	1.0	0.8	0.8	0.9	0.6	0.7	0.6	0.6	0.7	0.6	0.5	0.6	0.5	0.5	0.6	0.5	0.4	0.3	0.4	0.2	0.2	0.0
10th Grade	—	—	—	—	0.6	0.7	0.7	0.8	0.6	0.5	0.4	0.6	0.5	0.5	0.5	0.5	0.5	0.5	0.6	0.5	0.5	0.4	0.5	0.4	0.2	0.3	+0.1
12th Grade	—	—	—	—	0.5	0.5	0.5	0.4	0.4	0.4	0.3	0.4	0.4	0.4	0.5	0.5	0.4	0.4	0.3	0.7	0.6	0.4	0.4	0.5	0.3	0.3	-0.1
Without a Needle ^j																											
8th Grade	—	—	—	—	0.8	1.0	0.8	0.8	0.9	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.5	0.6	0.4	0.5	0.4	0.3	0.3	0.2	0.2	0.2	0.0
10th Grade	—	—	—	—	0.8	0.9	1.1	1.0	1.1	1.1	0.7	0.8	0.5	0.7	0.7	0.6	0.6	0.6	0.6	0.6	0.5	0.4	0.4	0.3	0.3	0.2	-0.2
12th Grade	—	—	—	—	1.0	1.0	1.2	0.8	1.0	1.6	0.8	0.8	0.8	0.7	0.8	0.6	1.0	0.5	0.6	0.8	0.7	0.4	0.4	0.5	0.4	0.3	-0.1

(Table continued on next page.)

TABLE 6 (cont.)
Trends in Annual Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
Narcotics other than Heroin ^{k,l}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	3.5	3.3	3.6	3.8	4.7	5.4	6.2	6.3	6.7	7.0	6.7‡	9.4	9.3	9.5	9.0	9.0	9.2	9.1	9.2	8.7	8.7	7.9	7.1	6.1	5.4	4.8	-0.6
OxyContin ^{k,n,v}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	1.3	1.7	1.7	1.8	2.6	1.8	2.1	2.0	2.1	1.8	1.6	2.0	1.0	0.8	0.9	+0.1
10th Grade	—	—	—	—	—	—	—	—	—	—	—	3.0	3.6	3.5	3.2	3.8	3.9	3.6	5.1	4.6	3.9	3.0	3.4	3.0	2.6	2.1	-0.5
12th Grade	—	—	—	—	—	—	—	—	—	—	—	4.0	4.5	5.0	5.5	4.3	5.2	4.7	4.9	5.1	4.9	4.3	3.6	3.3	3.7	3.4	-0.2
Vicodin ^{k,n,v}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	2.5	2.8	2.5	2.6	3.0	2.7	2.9	2.5	2.7	2.1	1.3	1.4	1.0	0.9	0.8	0.0
10th Grade	—	—	—	—	—	—	—	—	—	—	—	6.9	7.2	6.2	5.9	7.0	7.2	6.7	8.1	7.7	5.9	4.4	4.6	3.4	2.5	1.7	-0.8
12th Grade	—	—	—	—	—	—	—	—	—	—	—	9.6	10.5	9.3	9.5	9.7	9.6	9.7	9.7	8.0	8.1	7.5	5.3	4.8	4.4	2.9	-1.5 ss
Amphetamines ^{k,m}																											
8th Grade	6.2	6.5	7.2	7.9	8.7	9.1	8.1	7.2	6.9	6.5	6.7	5.5	5.5	4.9	4.9	4.7	4.2	4.5	4.1	3.9	3.5	2.9‡	4.2	4.3	4.1	3.5	-0.6
10th Grade	8.2	8.2	9.6	10.2	11.9	12.4	12.1	10.7	10.4	11.1	11.7	10.7	9.0	8.5	7.8	7.9	8.0	6.4	7.1	7.6	6.6	6.5‡	7.9	7.6	6.8	6.1	-0.7
12th Grade	8.2	7.1	8.4	9.4	9.3	9.5	10.2	10.1	10.2	10.5	10.9	11.1	9.9	10.0	8.6	8.1	7.5	6.8	6.6	7.4	8.2	7.9‡	9.2	8.1	7.7	6.7	-1.0
Ritalin ^{k,n,o}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	2.9	2.8	2.6	2.5	2.4	2.6	2.1	1.6	1.8	1.5	1.3	0.7	1.1	0.9	0.6	0.8	+0.2
10th Grade	—	—	—	—	—	—	—	—	—	—	4.8	4.8	4.1	3.4	3.4	3.6	2.8	2.9	3.6	2.7	2.6	1.9	1.8	1.8	1.6	1.2	-0.5
12th Grade	—	—	—	—	—	—	—	—	—	—	5.1	4.0	4.0	5.1	4.4	4.4	3.8	3.4	2.1	2.7	2.6	2.6	2.3	1.8	2.0	1.2	-0.8 s
Adderall ^{k,n,o}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.0	2.3	1.7	1.7	1.8	1.3	1.0	1.5	+0.5
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.7	5.3	4.6	4.5	4.4	4.6	5.2	4.2	-1.1
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.4	6.5	6.5	7.6	7.4	6.8	7.5	6.2	-1.3

(Table continued on next page.)

TABLE 6 (cont.)
Trends in Annual Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
Methamphetamine ^{n,o}																											
8th Grade	—	—	—	—	—	—	—	—	3.2	2.5	2.8	2.2	2.5	1.5	1.8	1.8	1.1	1.2	1.0	1.2	0.8	1.0	1.0	0.6	0.5	0.4	-0.1
10th Grade	—	—	—	—	—	—	—	—	4.6	4.0	3.7	3.9	3.3	3.0	2.9	1.8	1.6	1.5	1.6	1.6	1.4	1.0	1.0	0.8	0.8	0.4	-0.4 s
12th Grade	—	—	—	—	—	—	—	—	4.7	4.3	3.9	3.6	3.2	3.4	2.5	2.5	1.7	1.2	1.2	1.0	1.4	1.1	0.9	1.0	0.6	0.6	0.0
Crystal Methamphetamine (Ice) ^o																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.4	1.3	1.7	1.8	2.4	2.8	2.3	3.0	1.9	2.2	2.5	3.0	2.0	2.1	2.3	1.9	1.6	1.1	0.9	0.9	1.2	0.8	1.1	0.8	0.5	0.8	+0.3
Bath salts (synthetic stimulants) ^{n,o}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.8	1.0	0.5	0.4	0.9	+0.5 ss
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.6	0.9	0.9	0.7	0.8	+0.1
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.3	0.9	0.9	1.0	0.8	-0.2
Sedatives (Barbiturates) ^{k,p}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	3.4	2.8	3.4	4.1	4.7	4.9	5.1	5.5	5.8	6.2	5.7	6.7	6.0	6.5	7.2	6.6	6.2	5.8	5.2	4.8	4.3	4.5	4.8	4.3	3.6	3.0	-0.6
Tranquilizers ^{b,k}																											
8th Grade	1.8	2.0	2.1	2.4	2.7	3.3	2.9	2.6	2.5	2.6‡	2.8	2.6	2.7	2.5	2.8	2.6	2.4	2.4	2.6	2.8	2.0	1.8	1.8	1.7	1.7	1.7	0.0
10th Grade	3.2	3.5	3.3	3.3	4.0	4.6	4.9	5.1	5.4	5.6‡	7.3	6.3	5.3	5.1	4.8	5.2	5.3	4.6	5.0	5.1	4.5	4.3	3.7	3.9	3.9	4.1	+0.2
12th Grade	3.6	2.8	3.5	3.7	4.4	4.6	4.7	5.5	5.8	5.7‡	6.9	7.7	6.7	7.3	6.8	6.6	6.2	6.2	6.3	5.6	5.6	5.3	4.6	4.7	4.7	4.9	+0.2
Any Prescription Drug ^q																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	17.1	16.8	15.8	15.4	14.4	15.0	15.2	14.8‡	15.9	13.9	12.9	12.0	-1.0

(Table continued on next page.)

TABLE 6 (cont.)
Trends in Annual Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
OTC Cough/Cold Medicines ^{n,o}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	4.2	4.0	3.6	3.8	3.2	2.7	3.0	2.9	2.0	1.6	2.6	+1.0 ss
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	5.3	5.4	5.3	6.0	5.1	5.5	4.7	4.3	3.7	3.3	3.0	-0.3
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6.9	5.8	5.5	5.9	6.6	5.3	5.6	5.0	4.1	4.6	4.0	-0.5
Rohypnol ^r																											
8th Grade	—	—	—	—	—	1.0	0.8	0.8	0.5	0.5	0.7	0.3	0.5	0.6	0.7	0.5	0.7	0.5	0.4	0.5	0.8	0.4	0.4	0.3	0.3	0.5	+0.2
10th Grade	—	—	—	—	—	1.1	1.3	1.2	1.0	0.8	1.0	0.7	0.6	0.7	0.5	0.5	0.7	0.4	0.4	0.6	0.6	0.5	0.6	0.5	0.2	0.5	+0.3
12th Grade	—	—	—	—	—	1.1	1.2	1.4	1.0	0.8	0.9†	1.6	1.3	1.6	1.2	1.1	1.0	1.3	1.0	1.5	1.3	1.5	0.9	0.7	1.0	1.1	+0.1
GHB ^{n,w}																											
8th Grade	—	—	—	—	—	—	—	—	—	1.2	1.1	0.8	0.9	0.7	0.5	0.8	0.7	1.1	0.7	0.6	0.6	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	1.1	1.0	1.4	1.4	0.8	0.8	0.7	0.6	0.5	1.0	0.6	0.5	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	1.9	1.6	1.5	1.4	2.0	1.1	1.1	0.9	1.2	1.1	1.4	1.4	1.4	1.0	1.0	0.7	0.9	+0.2
Ketamine ^{n,x}																											
8th Grade	—	—	—	—	—	—	—	—	—	1.6	1.3	1.3	1.1	0.9	0.6	0.9	1.0	1.2	1.0	1.0	0.8	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	2.1	2.1	2.2	1.9	1.3	1.0	1.0	0.8	1.0	1.3	1.1	1.2	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	2.5	2.5	2.6	2.1	1.9	1.6	1.4	1.3	1.5	1.7	1.6	1.7	1.5	1.4	1.5	1.4	1.2	-0.2
Alcohol ^s																											
Any Use																											
8th Grade	54.0	53.7†	45.4	46.8	45.3	46.5	45.5	43.7	43.5	43.1	41.9	38.7	37.2	36.7	33.9	33.6	31.8	32.1	30.3	29.3	26.9	23.6	22.1	20.8	21.0	17.6	-3.4 sss
10th Grade	72.3	70.2†	63.4	63.9	63.5	65.0	65.2	62.7	63.7	65.3	63.5	60.0	59.3	58.2	56.7	55.8	56.3	52.5	52.8	52.1	49.8	48.5	47.1	44.0	41.9	38.3	-3.5 ss
12th Grade	77.7	76.8†	72.7	73.0	73.7	72.5	74.8	74.3	73.8	73.2	73.3	71.5	70.1	70.6	68.6	66.5	66.4	65.5	66.2	65.2	63.5	63.5	62.0	60.2	58.2	55.6	-2.6
Been Drunk ^o																											
8th Grade	17.5	18.3	18.2	18.2	18.4	19.8	18.4	17.9	18.5	18.5	16.6	15.0	14.5	14.5	14.1	13.9	12.6	12.7	12.2	11.5	10.5	8.6	8.4	7.3	7.7	5.7	-2.0 sss
10th Grade	40.1	37.0	37.8	38.0	38.5	40.1	40.7	38.3	40.9	41.6	39.9	35.4	34.7	35.1	34.2	34.5	34.4	30.0	31.2	29.9	28.8	28.2	27.1	24.6	23.4	20.5	-2.9 ss
12th Grade	52.7	50.3	49.6	51.7	52.5	51.9	53.2	52.0	53.2	51.8	53.2	50.4	48.0	51.8	47.7	47.9	46.1	45.6	47.0	44.0	42.2	45.0	43.5	41.4	37.7	37.3	-0.3

(Table continued on next page.)

TABLE 6 (cont.)
Trends in Annual Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change	
Flavored Alcoholic Beverages ^{e,n,y}																												
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	30.4	27.9	26.8	26.0	25.0	22.2	21.9	19.2	17.0	15.7	13.4	13.4	11.2	-2.2 s	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	49.7	48.5	48.8	45.9	43.4	41.5	41.0	38.3	37.8	35.6	33.2	31.4	26.1	-5.3 sss	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	55.2	55.8	58.4	54.7	53.6	51.8	53.4	47.9	47.0	44.4	44.2	43.6	42.8	40.0	-2.8	
Alcoholic Beverages containing Caffeine ^{n,o,z}																												
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	11.8	10.9	10.2	9.5	8.4	6.5	-1.9 ss	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	22.5	19.7	16.9	14.3	12.8	10.6	-2.2 ss	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	26.4	26.4	23.5	20.0	18.3	17.0	-1.3	
Tobacco using a Hookah ^e																												
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	17.1	18.5	18.3	21.4	22.9	19.8	13.0	-6.9 sss
Small cigars ^e																												
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	23.1	19.5	19.9	20.4	18.9	15.9	15.6	-0.3
Dissolvable Tobacco Products ^{e,n}																												
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.0	1.1	1.1	0.9	0.7	-0.2	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.6	1.2	1.3	1.1	0.9	-0.2	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.5	1.6	1.9	1.1	1.4	1.1	-0.3
Snus ^{e,n}																												
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.4	2.0	2.2	1.9	2.2	+0.3	
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6.9	5.2	4.5	4.0	3.0	-1.0	
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	7.9	7.9	7.7	5.8	5.8	5.8	+0.1

(Table continued on next page.)

TABLE 6 (cont.)
Trends in Annual Prevalence of Use of Various Drugs
in Grades 8, 10, and 12
(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
Steroids ^{k,u}																											
8th Grade	1.0	1.1	0.9	1.2	1.0	0.9	1.0	1.2	1.7	1.7	1.6	1.5	1.4	1.1	1.1	0.9	0.8	0.9	0.8	0.5	0.7	0.6	0.6	0.6	0.5	0.5	0.0
10th Grade	1.1	1.1	1.0	1.1	1.2	1.2	1.2	1.2	1.7	2.2	2.1	2.2	1.7	1.5	1.3	1.2	1.1	0.9	0.8	1.0	0.9	0.8	0.8	0.8	0.7	0.7	0.0
12th Grade	1.4	1.1	1.2	1.3	1.5	1.4	1.4	1.7	1.8	1.7	2.4	2.5	2.1	2.5	1.5	1.8	1.4	1.5	1.5	1.5	1.2	1.3	1.5	1.5	1.7	1.0	-0.7 ss
Previously surveyed drugs that have been dropped.																											
Nitrites ^e																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.9	0.5	0.9	1.1	1.1	1.6	1.2	1.4	0.9	0.6	0.6	1.1	0.9	0.8	0.6	0.5	0.8	0.6	0.9	—	—	—	—	—	—	—	—
Provigil ^{k,o}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.8	1.3	1.5	—	—	—	—	—	—
Methaqualone ^{e,k}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.5	0.6	0.2	0.8	0.7	1.1	1.0	1.1	1.1	0.3	0.8	0.9	0.6	0.8	0.9	0.8	0.5	0.5	0.6	0.3	0.3	0.4	—	—	—	—	—
Bidis ^{n,o}																											
8th Grade	—	—	—	—	—	—	—	—	—	3.9	2.7	2.7	2.0	1.7	1.6	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	6.4	4.9	3.1	2.8	2.1	1.6	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	9.2	7.0	5.9	4.0	3.6	3.3	2.3	1.7	1.9	1.5	1.4	—	—	—	—	—	—	—
Kreteks ^{n,o}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	2.6	2.6	2.0	1.9	1.4	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	6.0	4.9	3.8	3.7	2.8	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	—	10.1	8.4	6.7	6.5	7.1	6.2	6.8	6.8	5.5	4.6	2.9	3.0	1.6	1.6	—	—	—

Source. The Monitoring the Future study, the University of Michigan.

Note: See footnotes following Table 8.

TABLE 7
Trends in 30-Day Prevalence of Use of Various Drugs
in Grades 8, 10, and 12

		Percentage who used in last 30 days																										2015– 2016 change
		1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Any Illicit Drug ^a																												
8th Grade		5.7	6.8	8.4	10.9	12.4	14.6	12.9	12.1	12.2	11.9	11.7	10.4	9.7	8.4	8.5	8.1	7.4	7.6	8.1	9.5	8.5	7.7‡	8.7	8.3	8.1	6.9	-1.2 s
10th Grade		11.6	11.0	14.0	18.5	20.2	23.2	23.0	21.5	22.1	22.5	22.7	20.8	19.5	18.3	17.3	16.8	16.9	15.8	17.8	18.5	19.2	18.6‡	19.2	18.5	16.5	15.9	-0.5
12th Grade		16.4	14.4	18.3	21.9	23.8	24.6	26.2	25.6	25.9	24.9	25.7	25.4	24.1	23.4	23.1	21.5	21.9	22.3	23.3	23.8	25.2	25.2‡	25.2	23.7	23.6	24.4	+0.8
Any Illicit Drug other than Marijuana ^{a,b}																												
8th Grade		3.8	4.7	5.3	5.6	6.5	6.9	6.0	5.5	5.5	5.6‡	5.5	4.7	4.7	4.1	4.1	3.8	3.6	3.8	3.5	3.5	3.4	2.6‡	3.6	3.3	3.1	2.7	-0.5
10th Grade		5.5	5.7	6.5	7.1	8.9	8.9	8.8	8.6	8.6	8.5‡	8.7	8.1	6.9	6.9	6.4	6.3	6.9	5.3	5.7	5.8	5.4	5.0‡	4.9	5.6	4.9	4.4	-0.5
12th Grade		7.1	6.3	7.9	8.8	10.0	9.5	10.7	10.7	10.4	10.4‡	11.0	11.3	10.4	10.8	10.3	9.8	9.5	9.3	8.6	8.6	8.9	8.4‡	8.2	7.7	7.6	6.9	-0.7
Any Illicit Drug including Inhalants ^{a,c}																												
8th Grade		8.8	10.0	12.0	14.3	16.1	17.5	16.0	14.9	15.1	14.4	14.0	12.6	12.1	11.2	11.2	10.9	10.1	10.4	10.6	11.7	10.5	9.5‡	10.0	9.5	9.3	7.9	-1.4 s
10th Grade		13.1	12.6	15.5	20.0	21.6	24.5	24.1	22.5	23.1	23.6	23.6	21.7	20.5	19.3	18.4	17.7	18.1	16.8	18.8	19.4	20.1	19.3‡	20.0	19.1	17.1	16.4	-0.6
12th Grade		17.8	15.5	19.3	23.0	24.8	25.5	26.9	26.6	26.4	26.4	26.5	25.9	24.6	23.3	24.2	22.1	22.8	22.8	24.1	24.5	26.2	25.2‡	26.5	24.3	24.7	24.6	-0.1
Marijuana/Hashish																												
8th Grade		3.2	3.7	5.1	7.8	9.1	11.3	10.2	9.7	9.7	9.1	9.2	8.3	7.5	6.4	6.6	6.5	5.7	5.8	6.5	8.0	7.2	6.5	7.0	6.5	6.5	5.4	-1.1 s
10th Grade		8.7	8.1	10.9	15.8	17.2	20.4	20.5	18.7	19.4	19.7	19.8	17.8	17.0	15.9	15.2	14.2	14.2	13.8	15.9	16.7	17.6	17.0	18.0	16.6	14.8	14.0	-0.8
12th Grade		13.8	11.9	15.5	19.0	21.2	21.9	23.7	22.8	23.1	21.6	22.4	21.5	21.2	19.9	19.8	18.3	18.8	19.4	20.6	21.4	22.6	22.9	22.7	21.2	21.3	22.5	+1.2
Inhalants ^{c,d}																												
8th Grade		4.4	4.7	5.4	5.6	6.1	5.8	5.6	4.8	5.0	4.5	4.0	3.8	4.1	4.5	4.2	4.1	3.9	4.1	3.8	3.6	3.2	2.7	2.3	2.2	2.0	1.8	-0.2
10th Grade		2.7	2.7	3.3	3.6	3.5	3.3	3.0	2.9	2.6	2.6	2.4	2.4	2.2	2.4	2.2	2.3	2.5	2.1	2.2	2.0	1.7	1.4	1.3	1.1	1.2	1.0	-0.2
12th Grade		2.4	2.3	2.5	2.7	3.2	2.5	2.5	2.3	2.0	2.2	1.7	1.5	1.5	1.5	2.0	1.5	1.2	1.4	1.2	1.4	1.0	0.9	1.0	0.7	0.7	0.8	0.0
Hallucinogens ^{b,f}																												
8th Grade		0.8	1.1	1.2	1.3	1.7	1.9	1.8	1.4	1.3	1.2‡	1.6	1.2	1.2	1.0	1.1	0.9	1.0	0.9	0.9	1.0	1.0	0.6	0.8	0.5	0.6	0.6	0.0
10th Grade		1.6	1.8	1.9	2.4	3.3	2.8	3.3	3.2	2.9	2.3‡	2.1	1.6	1.5	1.6	1.5	1.5	1.7	1.3	1.4	1.6	1.4	1.2	1.1	1.2	0.9	0.9	0.0
12th Grade		2.2	2.1	2.7	3.1	4.4	3.5	3.9	3.8	3.5	2.6‡	3.3	2.3	1.8	1.9	1.9	1.5	1.7	2.2	1.6	1.9	1.6	1.6	1.4	1.5	1.6	1.4	-0.1

(Table continued on next page.)

TABLE 7 (cont.)
Trends in 30-Day Prevalence of Use of Various Drugs
in Grades 8, 10, and 12

	Percentage who used in last 30 days																											2015– 2016 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016		
LSD ^b																												
8th Grade	0.6	0.9	1.0	1.1	1.4	1.5	1.5	1.1	1.1	1.0	1.0	0.7	0.6	0.5	0.5	0.4	0.5	0.5	0.5	0.6	0.5	0.3	0.5	0.3	0.4	0.4	0.0	
10th Grade	1.5	1.6	1.6	2.0	3.0	2.4	2.8	2.7	2.3	1.6	1.5	0.7	0.6	0.6	0.6	0.7	0.7	0.7	0.5	0.7	0.7	0.5	0.6	0.6	0.6	0.7	+0.1	
12th Grade	1.9	2.0	2.4	2.6	4.0	2.5	3.1	3.2	2.7	1.6	2.3	0.7	0.6	0.7	0.7	0.6	0.6	1.1	0.5	0.8	0.8	0.8	0.8	1.0	1.1	1.0	-0.1	
Hallucinogens other than LSD ^b																												
8th Grade	0.3	0.4	0.5	0.7	0.8	0.9	0.7	0.7	0.6	0.6	1.1	1.0	1.0	0.8	0.9	0.7	0.7	0.7	0.7	0.8	0.7	0.5	0.5	0.4	0.3	0.3	0.0	
10th Grade	0.4	0.5	0.7	1.0	1.0	1.0	1.2	1.4	1.2	1.2	1.4	1.4	1.2	1.4	1.3	1.3	1.4	1.0	1.1	1.2	1.1	0.9	0.8	0.8	0.6	0.5	0.0	
12th Grade	0.7	0.5	0.8	1.2	1.3	1.6	1.7	1.6	1.6	1.7	1.9	2.0	1.5	1.7	1.6	1.3	1.4	1.6	1.4	1.5	1.2	1.3	1.0	1.0	0.9	0.7	-0.2	
Ecstasy (MDMA) ^g																												
8th Grade, original	—	—	—	—	1.0	1.0	0.9	0.8	1.4	1.8	1.4	0.7	0.8	0.6	0.7	0.6	0.8	0.6	1.1	0.6	0.5	0.5	0.4	—	—	—		
Revised	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	0.7	0.5	0.3	-0.2	
10th Grade, original	—	—	—	—	1.8	1.3	1.3	1.8	2.6	2.6	1.8	1.1	0.8	1.0	1.2	1.2	1.1	1.3	1.9	1.6	1.0	1.2	0.8	—	—	—		
Revised	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.1	0.9	0.5	-0.4 sss	
12th Grade, original	—	—	—	—	2.0	1.6	1.5	2.5	3.6	2.8	2.4	1.3	1.2	1.0	1.3	1.6	1.8	1.8	1.4	2.3	0.9	1.5	1.4	—	—	—		
Revised	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.5	1.1	0.9	-0.2	
Cocaine																												
8th Grade	0.5	0.7	0.7	1.0	1.2	1.3	1.1	1.4	1.3	1.2	1.2	1.1	0.9	0.9	1.0	1.0	0.9	0.8	0.8	0.6	0.8	0.5	0.5	0.5	0.5	0.3	-0.1	
10th Grade	0.7	0.7	0.9	1.2	1.7	1.7	2.0	2.1	1.8	1.8	1.3	1.6	1.3	1.7	1.5	1.5	1.3	1.2	0.9	0.9	0.7	0.8	0.8	0.6	0.8	0.4	-0.4 ss	
12th Grade	1.4	1.3	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.1	2.1	2.3	2.1	2.3	2.3	2.5	2.0	1.9	1.3	1.3	1.1	1.1	1.1	1.0	1.1	0.9	-0.3	
Crack																												
8th Grade	0.3	0.5	0.4	0.7	0.7	0.8	0.7	0.9	0.8	0.8	0.8	0.8	0.7	0.6	0.6	0.6	0.6	0.5	0.5	0.4	0.5	0.3	0.3	0.3	0.3	0.2	-0.1	
10th Grade	0.3	0.4	0.5	0.6	0.9	0.8	0.9	1.1	0.8	0.9	0.7	1.0	0.7	0.8	0.7	0.7	0.5	0.5	0.4	0.5	0.4	0.4	0.4	0.3	0.3	0.2	-0.1	
12th Grade	0.7	0.6	0.7	0.8	1.0	1.0	0.9	1.0	1.1	1.0	1.1	1.2	0.9	1.0	1.0	0.9	0.9	0.8	0.6	0.7	0.5	0.6	0.6	0.7	0.6	0.5	0.0	
Other Cocaine ^h																												
8th Grade	0.5	0.5	0.6	0.9	1.0	1.0	0.8	1.0	1.1	0.9	0.9	0.8	0.7	0.7	0.7	0.7	0.6	0.6	0.7	0.5	0.6	0.3	0.3	0.4	0.4	0.3	-0.1	
10th Grade	0.6	0.6	0.7	1.0	1.4	1.3	1.6	1.8	1.6	1.6	1.2	1.3	1.1	1.5	1.3	1.3	1.1	1.0	0.8	0.7	0.6	0.7	0.7	0.5	0.7	0.3	-0.4 ss	
12th Grade	1.2	1.0	1.2	1.3	1.3	1.6	2.0	2.0	2.5	1.7	1.8	1.9	1.8	2.2	2.0	2.4	1.7	1.7	1.1	1.1	1.0	1.0	0.9	0.9	1.1	0.6	-0.5 ss	

(Table continued on next page.)

TABLE 7 (cont.)
Trends in 30-Day Prevalence of Use of Various Drugs
in Grades 8, 10, and 12

		Percentage who used in last 30 days																										2015– 2016 change
		1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Heroin ^{l,j}																												
8th Grade		0.3	0.4	0.4	0.6	0.6	0.7	0.6	0.6	0.6	0.5	0.6	0.5	0.4	0.5	0.5	0.3	0.4	0.4	0.4	0.4	0.4	0.2	0.3	0.3	0.1	0.2	0.0
10th Grade		0.2	0.2	0.3	0.4	0.6	0.5	0.6	0.7	0.7	0.5	0.3	0.5	0.3	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.4	0.2	0.2	0.0
12th Grade		0.2	0.3	0.2	0.3	0.6	0.5	0.5	0.5	0.5	0.7	0.4	0.5	0.4	0.5	0.5	0.4	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.4	0.3	0.2	-0.1
With a Needle ^j																												
8th Grade		—	—	—	—	0.4	0.5	0.4	0.5	0.4	0.3	0.4	0.3	0.3	0.3	0.3	0.2	0.3	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.1	0.1	0.0
10th Grade		—	—	—	—	0.3	0.3	0.3	0.4	0.3	0.3	0.2	0.3	0.2	0.3	0.3	0.3	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.1	0.2	+0.1 s
12th Grade		—	—	—	—	0.3	0.4	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.2	0.3	0.3	0.2	0.2	0.1	0.4	0.4	0.3	0.2	0.3	0.2	0.2	-0.1
Without a Needle ^j																												
8th Grade		—	—	—	—	0.3	0.4	0.4	0.3	0.4	0.3	0.4	0.3	0.3	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.1	0.1	0.1	0.0
10th Grade		—	—	—	—	0.3	0.3	0.4	0.5	0.5	0.4	0.2	0.4	0.2	0.3	0.3	0.3	0.2	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.1	-0.1
12th Grade		—	—	—	—	0.6	0.4	0.6	0.4	0.4	0.7	0.3	0.5	0.4	0.3	0.5	0.3	0.4	0.2	0.3	0.4	0.4	0.2	0.2	0.4	0.3	0.1	-0.1
Narcotics other than Heroin ^{k,l}																												
8th Grade		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade		1.1	1.2	1.3	1.5	1.8	2.0	2.3	2.4	2.6	2.9	3.0‡	4.0	4.1	4.3	3.9	3.8	3.8	3.8	4.1	3.6	3.6	3.0	2.8	2.2	2.1	1.7	-0.4
Amphetamines ^{k,m}																												
8th Grade		2.6	3.3	3.6	3.6	4.2	4.6	3.8	3.3	3.4	3.4	3.2	2.8	2.7	2.3	2.3	2.1	2.0	2.2	1.9	1.8	1.8	1.3‡	2.3	2.1	1.9	1.7	-0.2
10th Grade		3.3	3.6	4.3	4.5	5.3	5.5	5.1	5.1	5.0	5.4	5.6	5.2	4.3	4.0	3.7	3.5	4.0	2.8	3.3	3.3	3.1	2.8‡	3.3	3.7	3.1	2.7	-0.4
12th Grade		3.2	2.8	3.7	4.0	4.0	4.1	4.8	4.6	4.5	5.0	5.6	5.5	5.0	4.6	3.9	3.7	3.7	2.9	3.0	3.3	3.7	3.3‡	4.2	3.8	3.2	3.0	-0.2
Methamphetamine ^{n,o}																												
8th Grade		—	—	—	—	—	—	—	—	1.1	0.8	1.3	1.1	1.2	0.6	0.7	0.6	0.6	0.7	0.5	0.7	0.4	0.5	0.4	0.2	0.3	0.3	0.0
10th Grade		—	—	—	—	—	—	—	—	1.8	2.0	1.5	1.8	1.4	1.3	1.1	0.7	0.4	0.7	0.6	0.7	0.5	0.6	0.4	0.3	0.3	0.2	-0.1
12th Grade		—	—	—	—	—	—	—	—	1.7	1.9	1.5	1.7	1.7	1.4	0.9	0.9	0.6	0.6	0.5	0.5	0.6	0.5	0.4	0.5	0.4	0.3	-0.1

(Table continued on next page.)

TABLE 7 (cont.)
Trends in 30-Day Prevalence of Use of Various Drugs
in Grades 8, 10, and 12

	Percentage who used in last 30 days																										2015– 2016 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Crystal Methamphetamine (Ice) ^o																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.6	0.5	0.6	0.7	1.1	1.1	0.8	1.2	0.8	1.0	1.1	1.2	0.8	0.8	0.9	0.7	0.6	0.6	0.5	0.6	0.6	0.4	0.8	0.4	0.3	0.4	+0.1
Sedatives (Barbiturates) ^{k,p}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	1.4	1.1	1.3	1.7	2.2	2.1	2.1	2.6	2.6	3.0	2.8	3.2	2.9†	2.9	3.3	3.0	2.7	2.8	2.5	2.2	1.8	2.0	2.2	2.0	1.7	1.5	-0.2
Tranquilizers ^{b,k}																											
8th Grade	0.8	0.8	0.9	1.1	1.2	1.5	1.2	1.2	1.1	1.4†	1.2	1.2	1.4	1.2	1.3	1.3	1.1	1.2	1.2	1.2	1.0	0.8	0.9	0.8	0.8	0.8	0.0
10th Grade	1.2	1.5	1.1	1.5	1.7	1.7	2.2	2.2	2.2	2.5†	2.9	2.9	2.4	2.3	2.3	2.4	2.6	1.9	2.0	2.2	1.9	1.7	1.6	1.6	1.7	1.5	-0.2
12th Grade	1.4	1.0	1.2	1.4	1.8	2.0	1.8	2.4	2.5	2.6†	2.9	3.3	2.8	3.1	2.9	2.7	2.6	2.6	2.7	2.5	2.3	2.1	2.0	2.1	2.0	1.9	-0.1
Any Prescription Drug ^q																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	8.6	8.1	7.8	7.2	7.3	6.9	7.2	7.0†	7.1	6.4	5.9	5.4	-0.5
Rohypnol ^r																											
8th Grade	—	—	—	—	—	0.5	0.3	0.4	0.3	0.3	0.4	0.2	0.1	0.2	0.2	0.4	0.3	0.1	0.2	0.2	0.6	0.1	0.1	0.2	0.1	0.2	+0.1
10th Grade	—	—	—	—	—	0.5	0.5	0.4	0.5	0.4	0.2	0.4	0.2	0.3	0.2	0.2	0.2	0.2	0.3	0.3	0.3	0.2	0.1	0.4	0.1	0.3	+0.2
12th Grade	—	—	—	—	—	0.5	0.3	0.3	0.3	0.4	0.3	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Alcohol ^s																											
Any Use																											
8th Grade	25.1	26.1†	24.3	25.5	24.6	26.2	24.5	23.0	24.0	22.4	21.5	19.6	19.7	18.6	17.1	17.2	15.9	15.9	14.9	13.8	12.7	11.0	10.2	9.0	9.7	7.3	-2.4 sss
10th Grade	42.8	39.9†	38.2	39.2	38.8	40.4	40.1	38.8	40.0	41.0	39.0	35.4	35.4	35.2	33.2	33.8	33.4	28.8	30.4	28.9	27.2	27.6	25.7	23.5	21.5	19.9	-1.6
12th Grade	54.0	51.3†	48.6	50.1	51.3	50.8	52.7	52.0	51.0	50.0	49.8	48.6	47.5	48.0	47.0	45.3	44.4	43.1	43.5	41.2	40.0	41.5	39.2	37.4	35.3	33.2	-2.1

(Table continued on next page.)

TABLE 7 (cont.)
Trends in 30-Day Prevalence of Use of Various Drugs
in Grades 8, 10, and 12

		Percentage who used in last 30 days																												2015– 2016 change
		1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016			
Been Drunk ^o																														
8th Grade		7.6	7.5	7.8	8.7	8.3	9.6	8.2	8.4	9.4	8.3	7.7	6.7	6.7	6.2	6.0	6.2	5.5	5.4	5.4	5.0	4.4	3.6	3.5	2.7	3.1	1.8	-1.3 sss		
10th Grade		20.5	18.1	19.8	20.3	20.8	21.3	22.4	21.1	22.5	23.5	21.9	18.3	18.2	18.5	17.6	18.8	18.1	14.4	15.5	14.7	13.7	14.5	12.8	11.2	10.3	9.0	-1.2		
12th Grade		31.6	29.9	28.9	30.8	33.2	31.3	34.2	32.9	32.9	32.3	32.7	30.3	30.9	32.5	30.2	30.0	28.7	27.6	27.4	26.8	25.0	28.1	26.0	23.5	20.6	20.4	-0.2		
Flavored Alcoholic Beverages ^{e,n}																														
8th Grade		—	—	—	—	—	—	—	—	—	—	—	—	—	14.6	12.9	13.1	12.2	10.2	9.5	9.4	8.6	7.6	6.3	5.7	5.5	4.0	-1.5 ss		
10th Grade		—	—	—	—	—	—	—	—	—	—	—	—	—	25.1	23.1	24.7	21.8	20.2	19.0	19.4	15.8	16.3	15.5	14.0	12.8	11.0	-1.8		
12th Grade		—	—	—	—	—	—	—	—	—	—	—	—	—	31.1	30.5	29.3	29.1	27.4	27.4	24.1	23.1	21.8	21.0	19.9	20.8	18.3	-2.5		
Cigarettes Any Use																														
8th Grade		14.3	15.5	16.7	18.6	19.1	21.0	19.4	19.1	17.5	14.6	12.2	10.7	10.2	9.2	9.3	8.7	7.1	6.8	6.5	7.1	6.1	4.9	4.5	4.0	3.6	2.6	-1.0 ss		
10th Grade		20.8	21.5	24.7	25.4	27.9	30.4	29.8	27.6	25.7	23.9	21.3	17.7	16.7	16.0	14.9	14.5	14.0	12.3	13.1	13.6	11.8	10.8	9.1	7.2	6.3	4.9	-1.5 ss		
12th Grade		28.3	27.8	29.9	31.2	33.5	34.0	36.5	35.1	34.6	31.4	29.5	26.7	24.4	25.0	23.2	21.6	21.6	20.4	20.1	19.2	18.7	17.1	16.3	13.6	11.4	10.5	-0.9		
Smokeless Tobacco ^t																														
8th Grade		6.9	7.0	6.6	7.7	7.1	7.1	5.5	4.8	4.5	4.2	4.0	3.3	4.1	4.1	3.3	3.7	3.2	3.5	3.7	4.1	3.5	2.8	2.8	3.0	3.2	2.5	-0.7		
10th Grade		10.0	9.6	10.4	10.5	9.7	8.6	8.9	7.5	6.5	6.1	6.9	6.1	5.3	4.9	5.6	5.7	6.1	5.0	6.5	7.5	6.6	6.4	6.4	5.3	4.9	3.5	-1.3 s		
12th Grade		—	11.4	10.7	11.1	12.2	9.8	9.7	8.8	8.4	7.6	7.8	6.5	6.7	6.7	7.6	6.1	6.6	6.5	8.4	8.5	8.3	7.9	8.1	8.4	6.1	6.6	+0.5		
Electronic Vaporizers ^{bb}																														
8th Grade		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	8.0	6.2	-1.8 s		
10th Grade		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	14.2	11.0	-3.3 ss		
12th Grade		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	16.3	12.5	-3.8 ss		
Large Cigars ^{bb}																														
8th Grade		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	1.9	2.4	1.5	-0.9 s		
10th Grade		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	3.9	3.4	2.3	-1.1 s		
12th Grade		—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6.4	7.0	6.5	-0.6		

(Table continued on next page.)

TABLE 7 (cont.)
Trends in 30-Day Prevalence of Use of Various Drugs
in Grades 8, 10, and 12

	Percentage who used in last 30 days																										2015– 2016 change
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Flavored Little Cigars^{bb}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	4.1	4.1	2.8	-1.3 ss
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6.9	6.1	4.9	-1.2
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	11.9	11.4	9.5	-1.9 s
Regular Little Cigars^{bb}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2.5	3.3	1.9	-1.5 ss
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	4.4	3.8	3.0	-0.8
12th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	7.0	7.8	6.1	-1.7 s
Steroids^{k,u}																											
8th Grade	0.4	0.5	0.5	0.5	0.6	0.4	0.5	0.5	0.7	0.8	0.7	0.8	0.7	0.5	0.5	0.5	0.4	0.5	0.4	0.3	0.4	0.3	0.3	0.2	0.3	0.3	0.0
10th Grade	0.6	0.6	0.5	0.6	0.6	0.5	0.7	0.6	0.9	1.0	0.9	1.0	0.8	0.8	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.4	0.4	0.4	0.4	0.3	0.0
12th Grade	0.8	0.6	0.7	0.9	0.7	0.7	1.0	1.1	0.9	0.8	1.3	1.4	1.3	1.6	0.9	1.1	1.0	1.0	1.0	1.1	0.7	0.9	1.0	0.9	1.0	0.7	-0.3
Previously surveyed drugs that have been dropped.																											
Nitrites^e																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.4	0.3	0.6	0.4	0.4	0.7	0.7	1.0	0.4	0.3	0.5	0.6	0.7	0.7	0.5	0.3	0.5	0.3	0.6	—	—	—	—	—	—	—	—
PCP^e																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.5	0.6	1.0	0.7	0.6	1.3	0.7	1.0	0.8	0.9	0.5	0.4	0.6	0.4	0.7	0.4	0.5	0.6	0.5	0.8	0.8	0.5	0.4	—	—	—	—
Methaqualone^{e,k}																											
8th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
10th Grade	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
12th Grade	0.2	0.4	0.1	0.4	0.4	0.6	0.3	0.6	0.4	0.2	0.5	0.3	0.4	0.5	0.5	0.4	0.4	0.2	0.3	0.2	0.2	0.3	—	—	—	—	—

Source: The Monitoring the Future study, the University of Michigan.

Note: See footnotes following Table 8.

TABLE 8
Trends in 30-Day Prevalence of Daily Use of Various Drugs
in Grades 8, 10, and 12

(Entries are percentages.)

	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015– 2016 change
Marijuana/Hashish																											
Daily ^{aa}																											
8th Grade	0.2	0.2	0.4	0.7	0.8	1.5	1.1	1.1	1.4	1.3	1.3	1.2	1.0	0.8	1.0	1.0	0.8	0.9	1.0	1.2	1.3	1.1	1.1	1.0	1.1	0.7	-0.3 s
10th Grade	0.8	0.8	1.0	2.2	2.8	3.5	3.7	3.6	3.8	3.8	4.5	3.9	3.6	3.2	3.1	2.8	2.8	2.7	2.8	3.3	3.6	3.5	4.0	3.4	3.0	2.5	-0.5
12th Grade	2.0	1.9	2.4	3.6	4.6	4.9	5.8	5.6	6.0	6.0	5.8	6.0	6.0	5.6	5.0	5.0	5.1	5.4	5.2	6.1	6.6	6.5	6.5	5.8	6.0	6.0	0.0
Alcohol ^{s,aa}																											
Any Daily Use																											
8th Grade	0.5	0.6‡	1.0	1.0	0.7	1.0	0.8	0.9	1.0	0.8	0.9	0.7	0.8	0.6	0.5	0.5	0.6	0.7	0.5	0.5	0.4	0.3	0.3	0.3	0.2	0.2	0.0
10th Grade	1.3	1.2‡	1.8	1.7	1.7	1.6	1.7	1.9	1.9	1.8	1.9	1.8	1.5	1.3	1.3	1.4	1.4	1.0	1.1	1.1	0.8	1.0	0.9	0.8	0.5	0.5	+0.1
12th Grade	3.6	3.4‡	3.4	2.9	3.5	3.7	3.9	3.9	3.4	2.9	3.6	3.5	3.2	2.8	3.1	3.0	3.1	2.8	2.5	2.7	2.1	2.5	2.2	1.9	1.9	1.3	-0.5 s
Been Drunk																											
Daily ^{o,aa}																											
8th Grade	0.1	0.1	0.2	0.3	0.2	0.2	0.2	0.3	0.4	0.3	0.2	0.3	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.0	0.0
10th Grade	0.2	0.3	0.4	0.4	0.6	0.4	0.6	0.6	0.7	0.5	0.6	0.5	0.5	0.4	0.4	0.5	0.5	0.3	0.4	0.3	0.2	0.4	0.3	0.3	0.1	0.1	0.0
12th Grade	0.9	0.8	0.9	1.2	1.3	1.6	2.0	1.5	1.9	1.7	1.4	1.2	1.6	1.8	1.5	1.6	1.3	1.4	1.1	1.6	1.3	1.5	1.3	1.1	0.8	0.8	+0.1
5+ Drinks in a Row																											
in Last 2 Weeks																											
8th Grade	10.9	11.3	11.3	12.1	12.3	13.3	12.3	11.5	13.1	11.7	11.0	10.3	9.8	9.4	8.4	8.7	8.3	8.1	7.8	7.2	6.4	5.1	5.1	4.1	4.6	3.4	-1.2 sss
10th Grade	21.0	19.1	21.0	21.9	22.0	22.8	23.1	22.4	23.5	24.1	22.8	20.3	20.0	19.9	19.0	19.9	19.6	16.0	17.5	16.3	14.7	15.6	13.7	12.6	10.9	9.7	-1.2
12th Grade	29.8	27.9	27.5	28.2	29.8	30.2	31.3	31.5	30.8	30.0	29.7	28.6	27.9	29.2	27.1	25.4	25.9	24.6	25.2	23.2	21.6	23.7	22.1	19.4	17.2	15.5	-1.7
Cigarettes																											
Any Daily Use																											
8th Grade	7.2	7.0	8.3	8.8	9.3	10.4	9.0	8.8	8.1	7.4	5.5	5.1	4.5	4.4	4.0	4.0	3.0	3.1	2.7	2.9	2.4	1.9	1.8	1.4	1.3	0.9	-0.4 s
10th Grade	12.6	12.3	14.2	14.6	16.3	18.3	18.0	15.8	15.9	14.0	12.2	10.1	8.9	8.3	7.5	7.6	7.2	5.9	6.3	6.6	5.5	5.0	4.4	3.2	3.0	1.9	-1.1 sss
12th Grade	18.5	17.2	19.0	19.4	21.6	22.2	24.6	22.4	23.1	20.6	19.0	16.9	15.8	15.6	13.6	12.2	12.3	11.4	11.2	10.7	10.3	9.3	8.5	6.7	5.5	4.8	-0.7

(Table continued on next page.)

TABLE 8 (cont.)
Trends in 30-Day Prevalence of Daily Use of Various Drugs
in Grades 8, 10, and 12

(Entries are percentages.)

	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>	<u>2006</u>	<u>2007</u>	<u>2008</u>	<u>2009</u>	<u>2010</u>	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2014</u>	<u>2015</u>	<u>2016</u>	<u>2015– 2016 change</u>
1/2 Pack+ /Day																											
8th Grade	3.1	2.9	3.5	3.6	3.4	4.3	3.5	3.6	3.3	2.8	2.3	2.1	1.8	1.7	1.7	1.5	1.1	1.2	1.0	0.9	0.7	0.6	0.7	0.5	0.4	0.3	-0.1
10th Grade	6.5	6.0	7.0	7.6	8.3	9.4	8.6	7.9	7.6	6.2	5.5	4.4	4.1	3.3	3.1	3.3	2.7	2.0	2.4	2.4	1.9	1.5	1.5	1.2	1.0	0.6	-0.4 s
12th Grade	10.7	10.0	10.9	11.2	12.4	13.0	14.3	12.6	13.2	11.3	10.3	9.1	8.4	8.0	6.9	5.9	5.7	5.4	5.0	4.7	4.3	4.0	3.4	2.6	2.1	1.8	-0.3
Smokeless Tobacco																											
Daily ^t																											
8th Grade	1.6	1.8	1.5	1.9	1.2	1.5	1.0	1.0	0.9	0.9	1.2	0.8	0.8	1.0	0.7	0.7	0.8	0.8	0.8	0.9	0.8	0.5	0.5	0.5	0.8	0.6	-0.1
10th Grade	3.3	3.0	3.3	3.0	2.7	2.2	2.2	2.2	1.5	1.9	2.2	1.7	1.8	1.6	1.9	1.7	1.6	1.4	1.9	2.5	1.7	2.0	1.9	1.8	1.6	1.0	-0.6 s
12th Grade	—	4.3	3.3	3.9	3.6	3.3	4.4	3.2	2.9	3.2	2.8	2.0	2.2	2.8	2.5	2.2	2.8	2.7	2.9	3.1	3.1	3.2	3.0	3.4	2.9	2.7	-0.2

Source. The Monitoring the Future study, the University of Michigan.

Note. See footnotes following Table 8.

Footnotes for Tables 5 through 8

Approximate Weighted <i>N</i> s	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
8th Graders	17,500	18,600	18,300	17,300	17,500	17,800	18,600	18,100	16,700	16,700	16,200	15,100	16,500
10th Graders	14,800	14,800	15,300	15,800	17,000	15,600	15,500	15,000	13,600	14,300	14,000	14,300	15,800
12th Graders	15,000	15,800	16,300	15,400	15,400	14,300	15,400	15,200	13,600	12,800	12,800	12,900	14,600

Approximate Weighted <i>N</i> s	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016
8th Graders	17,000	16,800	16,500	16,100	15,700	15,000	15,300	16,000	15,100	14,600	14,600	14,400	16,900
10th Graders	16,400	16,200	16,200	16,100	15,100	15,900	15,200	14,900	15,000	12,900	13,000	15,600	14,700
12th Graders	14,600	14,700	14,200	14,500	14,000	13,700	14,400	14,100	13,700	12,600	12,400	12,900	11,800

Notes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$. '—' indicates data not available. '‡' indicates that the question changed in the following year. See relevant footnote for that drug. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aFor 12th graders only: Use of any illicit drug includes any use of marijuana, LSD, other hallucinogens, crack, other cocaine, or heroin; or any use of narcotics other than heroin, amphetamines, sedatives (barbiturates), or tranquilizers not under a doctor's orders. For 8th and 10th graders only: The use of narcotics other than heroin and sedatives (barbiturates) has been excluded because these younger respondents appear to overreport use (perhaps because they include the use of nonprescription drugs in their answers). Due to changes in the amphetamines questions 2013 data for all grades for any illicit drug use, any illicit drug use other than marijuana and 8th and 10th grade any illicit drug use including inhalants are based on one half of the *N* indicated. 12th grade any illicit drug use including inhalants data are based on one form; *N* is one sixth of *N* indicated. 2014 data are based on all forms. See the amphetamine note for details.

^bIn 2001 the question text was changed on half of the questionnaire forms for each age group. Other psychedelics was changed to other hallucinogens and shrooms was added to the list of examples. For the tranquilizer list of examples, Miltown was replaced with Xanax. For 8th, 10th, and 12th graders: The 2001 data presented here are based on the changed forms only; *N* is one half of *N* indicated. In 2002 the remaining forms were changed to the new wording. The data are based on all forms beginning in 2002. Data for any illicit drug other than marijuana and data for hallucinogens are also affected by these changes and have been handled in a parallel manner. Hallucinogens, LSD, and hallucinogens other than LSD are based on five of six forms beginning in 2014; *N* is five sixths of *N* indicated.

^cFor 12th graders only: Data based on five of six forms in 1991–1998; *N* is five sixths of *N* indicated. Data based on three of six forms beginning in 1999; *N* is three sixths of *N* indicated. For 8th and 10th graders only, beginning in 2014 data based on two thirds of *N* indicated.

^dInhalants are unadjusted for underreporting of amyl and butyl nitrites.

^eFor 12th graders only: Data based on one of six forms; *N* is one sixth of *N* indicated. In 2011 for flavored alcoholic beverages Skyy Blue and Zima were dropped from the list of examples. An examination of the data did not show any effect from the wording change. In 2014 the PCP use questions were dropped; annual PCP use was moved to another form. In 2016 a question on use of tobacco using a hookah was added to two additional forms; *N* is three sixths of *N* indicated.

^fHallucinogens are unadjusted for underreporting of PCP.

^gFor 8th and 10th graders only: Data based on one of two forms in 1996; *N* is one half of *N* indicated. Data based on one third of *N* indicated in 1997–2001 due to changes in the questionnaire forms. Data based on two of four forms beginning in 2002; *N* is one half of *N* indicated. In 2014 a revised question on use of ecstasy (MDMA) including "Molly" was added to one form. The 2013 and 2014 "Original wording" data reported here are for only the questionnaires using the original question wording; *N* is one half of *N* indicated. Beginning in 2014 data

(Footnote continued on next page.)

Footnotes for Tables 5 through 8 (cont.)

reported here for the "Revised wording" are for only the questionnaires which include "Molly;" N is two sixths of N indicated in 2014 and five sixths of the N indicated in 2015. For 12th graders only: Data based on one of six forms in 1996–2001; N is one sixth of N indicated. Data based on two of six forms beginning in 2002; N is two sixths of N indicated. In 2014 a revised question on use of ecstasy (MDMA) including "Molly" was added to one form. The 2013 and 2014 "Original wording" data reported here are for only the questionnaires using the original question wording; N is two sixths of N indicated. Beginning in 2014 data reported for the "Revised wording" are for only the questionnaires which include "Molly."; N is one sixth of the N indicated in 2014 and three sixths of the N indicated in 2015.

^hFor 12th graders only: Data based on four of six forms; N is four sixths of N indicated.

ⁱIn 1995 the heroin question was changed in one of two forms for 8th and 10th graders and in three of six forms for 12th graders. Separate questions were asked for use with and without injection. In 1996, the heroin question was changed in the remaining 8th- and 10th-grade forms. Data presented here represent the combined data from all forms.

^jFor 8th and 10th graders only: Data based on one of two forms in 1995; N is one half of N indicated. Data based on all forms in 1996 through 2014. In 2015 the question was dropped from 1 form; N is four sixths of N indicated. For 12th graders only: Data based on three of six forms; N is three sixths of N indicated.

^kOnly drug use not under a doctor's orders is included here.

^lIn 2002 the question text was changed in half of the questionnaire forms. The list of examples of narcotics other than heroin was updated: Talwin, laudanum, and paregoric—all of which had negligible rates of use by 2001—were replaced with Vicodin, OxyContin, and Percocet. The 2002 data presented here are based on the changed forms only; N is one half of N indicated. In 2003, the remaining forms were changed to the new wording. The data are based on all forms beginning in 2003. In 2013 the list of examples was changed on one form: MS Contin, Roxycodone, Hydrocodone (Lortab, Lorcet, Norco), Suboxone, Tylox, and Tramadol were added to the list. An examination of the data did not show any effect from the wording change.

^mFor 8th, 10th, and 12th graders: In 2009, the question text was changed slightly in half of the forms. An examination of the data did not show any effect from the wording change. In 2010 the remaining forms were changed in a like manner. In 2011 the question text was changed slightly in one form; bennies, Benzedrine and Methadrine were dropped from the list of examples. An examination of the data did not show any effect from the wording change. In 2013 the question wording was changed slightly in two of the 8th and 10th grade questionnaires and in three of the 12th grade questionnaires. The new wording in 2013 asked "On how many occasions (if any) have taken amphetamines or other prescription stimulant drugs..." In contrast, the old wording did not include the text highlighted in red. Results in 2013 indicated higher prevalence in questionnaires with the new wording as compared to the old wording; it was proportionally 61% higher in 8th grade, 34% higher in 10th grade, and 21% higher in 12th grade. 2013 data are based on the changed forms only; for 8th, 10th, and 12th graders N is one half of N indicated. Beginning in 2014 all questionnaires included the new, updated wording.

ⁿFor 8th and 10th graders only: Data based on one of four forms; N is one third of N indicated. See text for detailed explanation. In 2011 for flavored alcoholic beverages: Skyy Blue and Zima were dropped from the list of examples. An examination of the data did not show any effect from the wording change. Annual synthetic marijuana use questions asked of one third of N indicated.

^oFor 12th graders only: Data based on two of six forms; N is two sixths of N indicated. Bidis and kreteks based on one of six forms beginning in 2009; N is one sixth N indicated.

^pFor 12th graders only: In 2004 the barbiturate question text was changed on half of the questionnaire forms. Barbiturates was changed to sedatives including barbiturates, and "have you taken barbiturates . . ." was changed to "have you taken sedatives . . ." In the list of examples downs, downers, goofballs, yellow, reds, blues, rainbows were changed to downs, or downers, and include Phenobarbital, Tuinal, Nembutal, and Seconal. An examination of the data did not show any effect from the wording change. In 2005 the remaining forms were changed in a like manner. In 2013 the question text was changed in all forms: Tuinal, Nembutal, and Seconal were replaced with Ambien, Lunesta, and Sonata. In one form the list of examples was also changed: Tuinal was dropped from the list and Dalmane, Restoril, Halcion, Intermezzo, and Zolpimist were added. An examination of the data did not show any effect from the wording change.

(Footnote continued on next page.)

Footnotes for Tables 5 through 8 (cont.)

^qThe use of any prescription drug includes use of any of the following: amphetamines, sedatives (barbiturates), narcotics other than heroin, or tranquilizers "...without a doctor telling you to use them."

^rFor 8th and 10th graders only: Data based on one of two forms in 1996; *N* is one half of *N* indicated. Data based on three of four forms in 1997–1998; *N* is two thirds of *N* indicated. Data based on two of four forms in 1999–2001; *N* is one third of *N* indicated. Data based on one of four forms beginning in 2002; *N* is one sixth of *N* indicated. See text for detailed explanation. For 12th graders only: Data based on one of six forms in 1996–2001; *N* is one sixth of *N* indicated. Data based on two of six forms in 2002–2009; *N* is two sixths of *N* indicated. Data for 2001 and 2002 are not comparable due to changes in the questionnaire forms. Data based on one of six forms beginning in 2010; *N* is one sixth of *N* indicated.

^sFor 8th, 10th, and 12th graders: In 1993, the question text was changed slightly in half of the forms to indicate that a drink meant more than just a few sips. The 1993 data are based on the changed forms only; *N* is one half of *N* indicated for these groups. In 1994 the remaining forms were changed to the new wording. The data are based on all forms beginning in 1994. In 2004, the question text was changed slightly in half of the forms. An examination of the data did not show any effect from the wording change. The remaining forms were changed in 2005.

^tFor 8th and 10th graders only: Data based on one of two forms for 1991–1996 and on two of four forms beginning in 1997; *N* is one half of *N* indicated. For 12th graders only: Data based on one of six forms; *N* is one sixth of *N* indicated. For all grades in 2011: snus and dissolvable tobacco were added to the list of examples. An examination of the data did not show any effect from the wording change.

^uFor 8th and 10th graders only: In 2006, the question text was changed slightly in half of the questionnaire forms. An examination of the data did not show any effect from the wording change. In 2007 the remaining forms were changed in a like manner. In 2008 the question text was changed slightly in half of the questionnaire forms. An examination of the data did not show any effect from the wording change. In 2009 the remaining forms were changed in a like manner. For 12th graders only: Data based on two of six forms in 1991–2005; *N* is two sixths of *N* indicated. Data based on three of six forms beginning in 2006; *N* is three sixths of *N* indicated. In 2006 a slightly altered version of the question was added to a third form. An examination of the data did not show any effect from the wording change. In 2007 the remaining forms were changed in a like manner. In 2008 the question text was changed slightly in two of the questionnaire forms. An examination of the data did not show any effect from the wording change. In 2009 the remaining form was changed in a like manner.

^vFor 12th graders only: Data based on two of six forms in 2002–2005; *N* is two sixths of *N* indicated. Data based on three of six forms beginning in 2006; *N* is three sixths of *N* indicated.

^wFor 12th graders only: Data based on two of six forms in 2000; *N* is two sixths of *N* indicated. Data based on three of six forms in 2001; *N* is three sixths of *N* indicated. Data based on one of six forms beginning in 2002; *N* is one sixth of *N* indicated.

^xFor 12th graders only: Data based on two of six forms in 2000; *N* is two sixths of *N* indicated. Data based on three of six forms in 2001–2009; *N* is three sixths of *N* indicated. Data based on two of six forms beginning in 2010; *N* is two sixths of *N* indicated.

^yThe 2003 flavored alcoholic beverage data were created by adjusting the 2004 data to reflect the change in the 2003 and 2004 alcopops data.

^zFor 8th and 10th graders only: Data based on one of four forms; *N* is one third of *N* indicated. See text for detailed explanation.

For 12th graders only: Data based on two of six forms; *N* is two sixths of *N* indicated. For all grades: In 2011 the question text was "...had an alcoholic beverage containing caffeine (like Four Loko or Joose)." In 2012 the question text was changed to "...had an alcoholic beverage mixed with an energy drink (like Red Bull)." An examination of the data did not show any effect from the wording changes.

^{aa}Daily use is defined as use on 20 or more occasions in the past 30 days except for cigarettes and smokeless tobacco, for which actual daily use is measured, and for 5+ drinks, for which the prevalence of having five or more drinks in a row in the last two weeks is measured.

^{bb}8th and 10th grade data based on one third of *N* indicated. 12th grade data based on two of six forms; *N* is two sixths of *N* indicated.

TABLE 9
Trends in Harmfulness of Drugs as Perceived by 8th Graders

How much do you think people risk harming themselves (physically or in other ways), if they . . .	Percentage saying great risk ^a														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Try marijuana once or twice ^b	40.4	39.1	36.2	31.6	28.9	27.9	25.3	28.1	28.0	29.0	27.7	28.2	30.2	31.9	31.4
Smoke marijuana occasionally ^b	57.9	56.3	53.8	48.6	45.9	44.3	43.1	45.0	45.7	47.4	46.3	46.0	48.6	50.5	48.9
Smoke marijuana regularly ^b	83.8	82.0	79.6	74.3	73.0	70.9	72.7	73.0	73.3	74.8	72.2	71.7	74.2	76.2	73.9
Try synthetic marijuana once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take synthetic marijuana occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try inhalants once or twice ^d	35.9	37.0	36.5	37.9	36.4	40.8	40.1	38.9	40.8	41.2	45.6	42.8	40.3	38.7	37.5
Take inhalants regularly ^d	65.6	64.4	64.6	65.5	64.8	68.2	68.7	67.2	68.8	69.9	71.6	69.9	67.4	66.4	64.1
Take LSD once or twice ^e	—	—	42.1	38.3	36.7	36.5	37.0	34.9	34.1	34.0	31.6	29.6	27.9	26.8	25.8
Take LSD regularly ^e	—	—	68.3	65.8	64.4	63.6	64.1	59.6	58.8	57.5	52.9	49.3	48.2	45.2	44.0
Try ecstasy (MDMA) once or twice ^f	—	—	—	—	—	—	—	—	—	—	35.8	38.9	41.9	42.5	40.0
Take ecstasy (MDMA) occasionally ^f	—	—	—	—	—	—	—	—	—	—	55.5	61.8	65.8	65.1	60.8
Try salvia once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take salvia occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try crack once or twice ^d	62.8	61.2	57.2	54.4	50.8	51.0	49.9	49.3	48.7	48.5	48.6	47.4	48.7	49.0	49.6
Take crack occasionally ^d	82.2	79.6	76.8	74.4	72.1	71.6	71.2	70.6	70.6	70.1	70.0	69.7	70.3	70.4	69.4
Try cocaine powder once or twice ^d	55.5	54.1	50.7	48.4	44.9	45.2	45.0	44.0	43.3	43.9	43.2	43.7	44.4	44.2	44.2
Take cocaine powder occasionally ^d	77.0	74.3	71.8	69.1	66.4	65.7	65.8	65.2	65.4	65.5	65.8	64.9	65.8	66.0	65.3
Try heroin once or twice without using a needle ^e	—	—	—	—	60.1	61.3	63.0	62.8	63.0	62.0	61.1	62.6	62.7	61.6	61.4
Take heroin occasionally without using a needle ^e	—	—	—	—	76.8	76.6	79.2	79.0	78.9	78.6	78.5	78.5	77.8	77.5	76.8
Try OxyContin once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take OxyContin occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try Vicodin once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take Vicodin occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try Adderall once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take Adderall occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try bath salts (synthetic stimulants) once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take bath salts (synthetic stimulants) occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try cough/cold medicine once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take cough/cold medicine occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try one or two drinks of an alcoholic beverage (beer, wine, liquor) ^b	11.0	12.1	12.4	11.6	11.6	11.8	10.4	12.1	11.6	11.9	12.2	12.5	12.6	13.7	13.9
Take one or two drinks nearly every day ^b	31.8	32.4	32.6	29.9	30.5	28.6	29.1	30.3	29.7	30.4	30.0	29.6	29.9	31.0	31.4
Have five or more drinks once or twice each weekend ^b	59.1	58.0	57.7	54.7	54.1	51.8	55.6	56.0	55.3	55.9	56.1	56.4	56.5	56.9	57.2
Smoke one to five cigarettes per day ^c	—	—	—	—	—	—	—	—	26.9	28.9	30.5	32.8	33.4	37.0	37.5
Smoke one or more packs of cigarettes per day ^a	51.6	50.8	52.7	50.8	49.8	50.4	52.6	54.3	54.8	58.8	57.1	57.5	57.7	62.4	61.5
Use electronic cigarettes (e-cigarettes) regularly ^h	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Smoke little cigars or cigarillos regularly ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Use smokeless tobacco regularly	35.1	35.1	36.9	35.5	33.5	34.0	35.2	36.5	37.1	39.0	38.2	39.4	39.7	41.3	40.8
Take dissolvable tobacco regularly ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take snus regularly ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take steroids ⁱ	64.2	69.5	70.2	67.6	—	—	—	—	—	—	—	—	—	—	—
Approximate weighted N = 17,400 18,700 18,400 17,400 17,500 17,900 18,800 18,100 16,700 16,700 16,200 15,100 16,500 17,000 16,800															

Table continued on next page.

TABLE 9 (cont.)
Trends in Harmfulness of Drugs as Perceived by 8th Graders

How much do you think people risk harming themselves (physically or in other ways), if they . . .	Percentage saying great risk ^a											2015– 2016 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Try marijuana once or twice ^b	32.2	32.8	31.1	29.5	29.5	28.2	26.0	24.1	23.0	23.0	22.8	-0.2
Smoke marijuana occasionally ^b	48.9	50.2	48.1	44.8	44.1	43.4	41.7	37.2	36.7	36.8	36.8	+0.1
Smoke marijuana regularly ^b	73.2	74.3	72.0	69.8	68.0	68.3	66.9	61.0	58.9	58.0	57.5	-0.4
Try synthetic marijuana once or twice ^c	—	—	—	—	—	—	24.4	24.2	23.9	26.0	27.5	+1.5
Take synthetic marijuana occasionally ^c	—	—	—	—	—	—	36.8	36.2	32.4	33.5	35.4	+1.9
Try inhalants once or twice ^d	35.8	35.9	33.9	34.1	35.5	34.7	34.2	33.7	34.5	33.7	32.0	-1.8
Take inhalants regularly ^d	62.1	61.9	59.2	58.1	60.6	59.0	59.0	56.7	55.3	54.1	52.1	-1.9
Take LSD once or twice ^e	23.8	22.8	21.9	21.4	23.6	21.7	19.9	19.6	20.0	22.2	22.6	+0.4
Take LSD regularly ^e	40.0	38.5	36.9	37.0	38.6	37.8	35.0	34.5	33.7	37.0	36.8	-0.2
Try ecstasy (MDMA) once or twice ^f	32.8	30.4	28.6	26.0	27.0	25.4	23.6	24.1†	46.1	45.5	42.5	-3.0
Take ecstasy (MDMA) occasionally ^f	52.0	48.6	46.8	43.9	45.0	43.7	41.0	42.1†	59.7	58.5	54.0	-4.5 ss
Try salvia once or twice ^c	—	—	—	—	—	—	9.5	8.5	—	—	—	—
Take salvia occasionally ^c	—	—	—	—	—	—	16.1	14.6	—	—	—	—
Try crack once or twice ^d	47.6	47.3	47.1	46.6	49.6	48.1	47.0	47.1	48.3	49.6	48.9	-0.7
Take crack occasionally ^d	68.7	68.3	67.9	66.6	68.4	67.7	67.8	66.5	65.5	65.7	65.7	+0.1
Try cocaine powder once or twice ^d	43.5	43.5	42.7	42.3	45.7	43.3	42.8	43.5	43.9	44.3	44.3	-0.1
Take cocaine powder occasionally ^d	64.0	64.2	62.7	62.3	64.2	63.5	63.3	62.7	61.8	61.6	62.4	+0.8
Try heroin once or twice without using a needle ^e	60.4	60.3	60.8	60.0	62.3	61.7	59.1	59.8	60.9	61.4	59.2	-2.2
Take heroin occasionally without using a needle ^e	75.3	76.4	75.5	74.0	76.7	75.9	75.1	73.4	73.2	72.7	70.3	-2.4
Try OxyContin once or twice ^c	—	—	—	—	—	—	21.9	19.9	22.1	20.2	21.3	+1.1
Take OxyContin occasionally ^c	—	—	—	—	—	—	35.3	32.6	34.4	32.5	33.5	+0.9
Try Vicodin once or twice ^c	—	—	—	—	—	—	17.5	15.0	18.4	16.9	18.3	+1.4
Take Vicodin occasionally ^c	—	—	—	—	—	—	29.4	26.2	28.2	26.7	28.8	+2.1
Try Adderall once or twice ^c	—	—	—	—	—	—	17.6	16.5	20.7	19.2	21.4	+2.2 s
Take Adderall occasionally ^c	—	—	—	—	—	—	29.9	28.3	32.5	32.0	35.9	+3.9 ss
Try bath salts (synthetic stimulants) once or twice ^c	—	—	—	—	—	—	24.9	39.3	36.8	33.9	31.8	-2.1
Take bath salts (synthetic stimulants) occasionally ^c	—	—	—	—	—	—	38.8	51.9	49.1	45.5	42.5	-3.0 s
Try cough/cold medicine once or twice ^c	—	—	—	—	—	—	21.2	20.1	22.9	20.9	23.5	+2.6 ss
Take cough/cold medicine occasionally ^c	—	—	—	—	—	—	38.8	37.3	37.9	37.3	38.6	+1.3
Try one or two drinks of an alcoholic beverage (beer, wine, liquor) ^b	14.2	14.9	13.5	14.4	14.9	14.5	13.9	13.7	14.8	15.3	14.7	-0.6
Take one or two drinks nearly every day ^b	31.3	32.6	31.5	31.5	32.3	31.8	31.4	30.6	31.0	30.9	30.7	-0.2
Have five or more drinks once or twice each weekend ^b	56.4	57.9	57.0	55.8	57.2	58.4	58.2	55.7	54.3	53.9	53.4	-0.5
Smoke one to five cigarettes per day ^c	37.0	38.6	38.6	38.6	38.2	37.4	40.4	42.8	41.9	41.7	43.2	+1.5
Smoke one or more packs of cigarettes per day ^a	59.4	61.1	59.8	59.1	60.9	62.5	62.6	62.4	62.1	63.0	61.2	-1.7
Use electronic cigarettes (e-cigarettes) regularly ^h	—	—	—	—	—	—	—	—	14.5	18.5	21.3	+2.8 ss
Smoke little cigars or cigarillos regularly ^c	—	—	—	—	—	—	—	—	28.8	31.0	32.5	+1.5
Use smokeless tobacco regularly	39.5	41.8	41.0	40.8	41.8	40.8	37.8	36.2	34.5	36.6	35.1	-1.5
Take dissolvable tobacco regularly ^c	—	—	—	—	—	—	34.8	32.2	33.5	33.0	34.3	+1.3
Take snus regularly ^c	—	—	—	—	—	—	42.2	38.9	38.3	37.7	37.9	+0.2
Take steroids ⁱ	—	—	—	—	—	—	—	—	—	—	—	—

Approximate weighted N = 16,500 16,100 15,700 15,000 15,300 16,000 15,100 14,600 14,600 14,400 16,900

Table continued on next page.

TABLE 9 (cont.)
Trends in Harmfulness of Drugs as Perceived by 8th Graders

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. "‡" indicates that the question changed the following year.

^aAnswer alternatives were: (1) No risk, (2) Slight risk, (3) Moderate risk, (4) Great risk, and (5) Can't say, drug unfamiliar.

^bBeginning in 2012 data based on two thirds of N indicated.

^cData based on one third of N indicated.

^dBeginning in 1997, data based on two thirds of N indicated due to changes in questionnaire forms.

^eData based on one of two forms in 1993–1996; N is one half of N indicated. Beginning in 1997, data based on one third of N indicated due to changes in questionnaire forms.

^fBeginning in 2014 data are based on the revised question which included "Molly," N is one third of N indicated in 2014 and two thirds of N indicated in 2015. 2014 and 2015 data are not comparable to earlier years due to the revision of the question text.

^gBeginning in 1999, data based on two thirds of N indicated due to changes in questionnaire forms.

^hE-cigarette data based on two thirds of N indicated. Little cigars or cigarillos data based on one third N indicated.

ⁱData based on two forms in 1991 and 1992. Data based on one of two forms in 1993 and 1994; N is one half of N indicated.

TABLE 10
Trends in Harmfulness of Drugs as Perceived by 10th Graders

<i>How much do you think people risk harming themselves (physically or in other ways), if they . . .</i>	<i>Percentage saying great risk ^a</i>														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
Try marijuana once or twice ^b	30.0	31.9	29.7	24.4	21.5	20.0	18.8	19.6	19.2	18.5	17.9	19.9	21.1	22.0	22.3
Smoke marijuana occasionally ^b	48.6	48.9	46.1	38.9	35.4	32.8	31.9	32.5	33.5	32.4	31.2	32.0	34.9	36.2	36.6
Smoke marijuana regularly ^b	82.1	81.1	78.5	71.3	67.9	65.9	65.9	65.8	65.9	64.7	62.8	60.8	63.9	65.6	65.5
Try synthetic marijuana once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take synthetic marijuana occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try inhalants once or twice ^d	37.8	38.7	40.9	42.7	41.6	47.2	47.5	45.8	48.2	46.6	49.9	48.7	47.7	46.7	45.7
Take inhalants regularly ^d	69.8	67.9	69.6	71.5	71.8	75.8	74.5	73.3	76.3	75.0	76.4	73.4	72.2	73.0	71.2
Take LSD once or twice ^e	—	—	48.7	46.5	44.7	45.1	44.5	43.5	45.0	43.0	41.3	40.1	40.8	40.6	40.3
Take LSD regularly ^e	—	—	78.9	75.9	75.5	75.3	73.8	72.3	73.9	72.0	68.8	64.9	63.0	63.1	60.8
Try ecstasy (MDMA) once or twice ^f	—	—	—	—	—	—	—	—	—	—	39.4	43.5	49.7	52.0	51.4
Take ecstasy (MDMA) occasionally ^f	—	—	—	—	—	—	—	—	—	—	64.8	67.3	71.7	74.6	72.8
Try salvia once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take salvia occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try crack once or twice ^d	70.4	69.6	66.6	64.7	60.9	60.9	59.2	58.0	57.8	56.1	57.1	57.4	57.6	56.7	57.0
Take crack occasionally ^d	87.4	86.4	84.4	83.1	81.2	80.3	78.7	77.5	79.1	76.9	77.3	75.7	76.4	76.7	76.9
Try cocaine powder once or twice ^d	59.1	59.2	57.5	56.4	53.5	53.6	52.2	50.9	51.6	48.8	50.6	51.3	51.8	50.7	51.3
Take cocaine powder occasionally ^d	82.2	80.1	79.1	77.8	75.6	75.0	73.9	71.8	73.6	70.9	72.3	71.0	71.4	72.2	72.4
Try heroin once or twice without using a needle ^e	—	—	—	—	70.7	72.1	73.1	71.7	73.7	71.7	72.0	72.2	70.6	72.0	72.4
Take heroin occasionally without using a needle ^e	—	—	—	—	85.1	85.8	86.5	84.9	86.5	85.2	85.4	83.4	83.5	85.4	85.2
Try OxyContin once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take OxyContin occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try Vicodin once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take Vicodin occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try Adderall once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take Adderall occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try bath salts (synthetic stimulants) once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take bath salts (synthetic stimulants) occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try cough/cold medicine once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take cough/cold medicine occasionally ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try one or two drinks of an alcoholic beverage (beer, wine, liquor) ^b	9.0	10.1	10.9	9.4	9.3	8.9	9.0	10.1	10.5	9.6	9.8	11.5	11.5	10.8	11.5
Take one or two drinks nearly every day ^b	36.1	36.8	35.9	32.5	31.7	31.2	31.8	31.9	32.9	32.3	31.5	31.0	30.9	31.3	32.6
Have five or more drinks once or twice each weekend ^u	54.7	55.9	54.9	52.9	52.0	50.9	51.8	52.5	51.9	51.0	50.7	51.7	51.6	51.7	53.3
Smoke one to five cigarettes per day ^c	—	—	—	—	—	—	—	—	28.4	30.2	32.4	35.1	38.1	39.7	41.0
Smoke one or more packs of cigarettes per day ^g	60.3	59.3	60.7	59.0	57.0	57.9	59.9	61.9	62.7	65.9	64.7	64.3	65.7	68.4	68.1
Use electronic cigarettes (e-cigarettes) regularly ^h	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Smoke little cigars or cigarillos regularly ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Use smokeless tobacco regularly	40.3	39.6	44.2	42.2	38.2	41.0	42.2	42.8	44.2	46.7	46.2	46.9	48.0	47.8	46.1
Take dissolvable tobacco regularly ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take snus regularly ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take steroids ⁱ	67.1	72.7	73.4	72.5	—	—	—	—	—	—	—	—	—	—	—
<i>Approximate weighted N =</i>	<i>14,700</i>	<i>14,800</i>	<i>15,300</i>	<i>15,900</i>	<i>17,000</i>	<i>15,700</i>	<i>15,600</i>	<i>15,000</i>	<i>13,600</i>	<i>14,300</i>	<i>14,000</i>	<i>14,300</i>	<i>15,800</i>	<i>16,400</i>	<i>16,200</i>

Table continued on next page.

TABLE 10 (cont.)
Trends in Harmfulness of Drugs as Perceived by 10th Graders

<i>How much do you think people risk harming themselves (physically or in other ways), if they . . .</i>	Percentage saying great risk ^a											2015– 2016 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Try marijuana once or twice ^b	22.2	22.2	23.1	20.5	19.9	19.3	17.2	15.7	15.2	15.8	16.4	+0.6
Smoke marijuana occasionally ^b	35.6	36.0	37.0	32.9	30.9	30.1	26.8	25.1	23.9	24.7	24.4	-0.3
Smoke marijuana regularly ^b	64.9	64.5	64.8	59.5	57.2	55.2	50.9	46.5	45.4	43.2	44.0	+0.8
Try synthetic marijuana once or twice ^c	—	—	—	—	—	—	24.6	24.1	25.0	26.3	26.8	+0.5
Take synthetic marijuana occasionally ^c	—	—	—	—	—	—	34.9	32.8	30.7	31.7	31.8	+0.1
Try inhalants once or twice ^d	43.9	43.0	41.2	42.0	42.5	42.4	42.4	43.0	43.1	43.1	40.7	-2.4 s
Take inhalants regularly ^d	70.2	68.6	66.8	66.8	67.1	66.2	66.1	65.9	64.7	63.1	59.7	-3.4 ss
Take LSD once or twice ^e	38.8	35.4	34.6	34.9	33.9	34.2	34.7	34.7	34.5	36.4	34.4	-2.0
Take LSD regularly ^e	60.7	56.8	55.7	56.7	56.1	54.9	56.4	55.9	54.8	58.3	55.2	-3.0
Try ecstasy (MDMA) once or twice ^f	48.4	45.3	43.2	38.9	36.3	37.2	36.2	36.0†	53.2	54.8	54.2	-0.6
Take ecstasy (MDMA) occasionally ^f	71.3	68.2	66.4	62.1	59.2	60.8	59.8	58.6†	69.0	70.1	69.3	-0.8
Try salvia once or twice ^c	—	—	—	—	—	—	12.2	10.7	—	—	—	—
Take salvia occasionally ^c	—	—	—	—	—	—	20.3	17.1	—	—	—	—
Try crack once or twice ^d	56.6	56.4	56.5	57.7	58.1	59.5	59.0	60.2	61.4	62.5	61.3	-1.2
Take crack occasionally ^d	76.2	76.0	76.5	75.9	76.2	76.5	76.7	77.8	76.4	77.5	75.2	-2.3 s
Try cocaine powder once or twice ^d	50.2	49.5	49.8	50.8	52.9	53.0	53.4	54.5	54.1	54.8	54.6	-0.2
Take cocaine powder occasionally ^d	71.3	70.9	71.1	71.0	72.2	72.0	72.6	72.8	71.7	72.6	70.9	-1.7
Try heroin once or twice without using a needle ^e	70.0	70.5	70.8	72.2	73.0	72.9	72.6	73.2	72.6	74.1	73.3	-0.9
Take heroin occasionally without using a needle ^e	83.6	84.2	83.1	83.3	84.8	83.4	84.4	84.0	82.5	83.3	82.2	-1.1
Try OxyContin once or twice ^c	—	—	—	—	—	—	30.9	29.4	29.7	29.9	28.7	-1.2
Take OxyContin occasionally ^c	—	—	—	—	—	—	48.3	44.7	44.4	43.7	41.4	-2.3
Try Vicodin once or twice ^c	—	—	—	—	—	—	23.2	21.0	22.5	24.1	21.8	-2.3 s
Take Vicodin occasionally ^c	—	—	—	—	—	—	40.3	36.0	36.4	35.4	32.6	-2.8 s
Try Adderall once or twice ^c	—	—	—	—	—	—	19.7	17.6	22.2	22.9	22.5	-0.3
Take Adderall occasionally ^c	—	—	—	—	—	—	34.3	30.5	37.0	37.0	35.8	-1.2
Try bath salts (synthetic stimulants) once or twice ^c	—	—	—	—	—	—	32.3	50.1	49.6	49.1	42.7	-6.4 sss
Take bath salts (synthetic stimulants) occasionally ^c	—	—	—	—	—	—	44.9	61.8	61.1	60.4	53.0	-7.3 sss
Try cough/cold medicine once or twice ^c	—	—	—	—	—	—	23.6	21.6	22.9	24.0	24.0	+0.1
Take cough/cold medicine occasionally ^c	—	—	—	—	—	—	40.4	37.3	38.3	38.2	37.6	-0.7
Try one or two drinks of an alcoholic beverage (beer, wine, liquor) ^b	11.1	11.6	12.6	11.9	11.9	12.3	11.3	11.3	11.6	12.4	13.3	+0.8
Take one or two drinks nearly every day ^b	31.7	33.3	35.0	33.8	33.1	32.9	31.8	30.6	31.3	31.2	32.2	+1.0
Have five or more drinks once or twice each weekend ^b	52.4	54.1	56.6	54.2	54.6	55.5	52.8	52.3	54.0	54.5	54.5	0.0
Smoke one to five cigarettes per day ^c	41.3	41.7	43.5	42.8	41.4	44.8	49.1	47.7	52.0	52.9	53.0	+0.2
Smoke one or more packs of cigarettes per day ^d	67.7	68.2	69.1	67.3	67.2	69.8	71.6	70.8	72.0	72.9	71.5	-1.4
Use electronic cigarettes (e-cigarettes) regularly ^h	—	—	—	—	—	—	—	—	14.1	17.0	19.1	+2.1 s
Smoke little cigars or cigarillos regularly ^c	—	—	—	—	—	—	—	—	31.0	34.9	35.3	+0.4
Use smokeless tobacco regularly	45.9	46.7	48.0	44.7	43.7	45.7	42.9	40.0	39.9	42.5	43.0	+0.4
Take dissolvable tobacco regularly ^c	—	—	—	—	—	—	33.3	31.3	32.0	35.6	34.2	-1.4
Take snus regularly ^c	—	—	—	—	—	—	41.0	38.9	38.8	41.8	39.9	-1.9
Take steroids ⁱ	—	—	—	—	—	—	—	—	—	—	—	—
<i>Approximate weighted N = 16,200 16,100 15,100 15,900 15,200 14,900 15,000 12,900 13,000 15,600 14,700</i>												

Table continued on next page.

TABLE 10 (cont.)
Trends in Harmfulness of Drugs as Perceived by 10th Graders

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. ‡ indicates that the question changed the following year.

^aAnswer alternatives were: (1) No risk, (2) Slight risk, (3) Moderate risk, (4) Great risk, and (5) Can't say, drug unfamiliar.

^bBeginning in 2012 data based on two thirds of N indicated.

^cData based on one third of N indicated.

^dBeginning in 1997, data based on two thirds of N indicated due to changes in questionnaire forms.

^eData based on one of two forms in 1993–1996; N is one half of N indicated. Beginning in 1997, data based on one third of N indicated due to changes in questionnaire forms.

^fBeginning in 2014 data are based on the revised question which included "Molly," N is one third of N indicated in 2014 and two thirds of N indicated in 2015. 2014 and 2015 data are not comparable to earlier years due to the revision of the question text.

^gBeginning in 1999, data based on two thirds of N indicated due to changes in questionnaire forms.

^hE-cigarette data based on two thirds of N indicated. Little cigars or cigarillos data based on one third N indicated.

ⁱData based on two forms in 1991 and 1992. Data based on one of two forms in 1993 and 1994; N is one half of N indicated.

TABLE 11
Trends in Harmfulness of Drugs as Perceived by 12th Graders

	Percentage saying great risk ^a																		
<i>How much do you think people risk harming themselves (physically or in other ways), if they . . .</i>	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990			
Try marijuana once or twice	15.1	11.4	9.5	8.1	9.4	10.0	13.0	11.5	12.7	14.7	14.8	15.1	18.4	19.0	23.6	23.1			
Smoke marijuana occasionally	18.1	15.0	13.4	12.4	13.5	14.7	19.1	18.3	20.6	22.6	24.5	25.0	30.4	31.7	36.5	36.9			
Smoke marijuana regularly	43.3	38.6	36.4	34.9	42.0	50.4	57.6	60.4	62.8	66.9	70.4	71.3	73.5	77.0	77.5	77.8			
Try synthetic marijuana once or twice	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Take synthetic marijuana occasionally	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Try LSD once or twice	49.4	45.7	43.2	42.7	41.6	43.9	45.5	44.9	44.7	45.4	43.5	42.0	44.9	45.7	46.0	44.7			
Take LSD regularly	81.4	80.8	79.1	81.1	82.4	83.0	83.5	83.5	83.2	83.8	82.9	82.6	83.8	84.2	84.3	84.5			
Try PCP once or twice	—	—	—	—	—	—	—	—	—	—	—	—	55.6	58.8	56.6	55.2			
Try ecstasy (MDMA) once or twice ^b	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Try salvia once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Take salvia occasionally	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Try cocaine once or twice	42.6	39.1	35.6	33.2	31.5	31.3	32.1	32.8	33.0	35.7	34.0	33.5	47.9	51.2	54.9	59.4			
Take cocaine occasionally	—	—	—	—	—	—	—	—	—	—	—	54.2	66.8	69.2	71.8	73.9			
Take cocaine regularly	73.1	72.3	68.2	68.2	69.5	69.2	71.2	73.0	74.3	78.8	79.0	82.2	88.5	89.2	90.2	91.1			
Try crack once or twice	—	—	—	—	—	—	—	—	—	—	—	—	57.0	62.1	62.9	64.3			
Take crack occasionally	—	—	—	—	—	—	—	—	—	—	—	—	70.4	73.2	75.3	80.4			
Take crack regularly	—	—	—	—	—	—	—	—	—	—	—	—	84.6	84.8	85.6	91.6			
Try cocaine powder once or twice	—	—	—	—	—	—	—	—	—	—	—	—	45.3	51.7	53.8	53.9			
Take cocaine powder occasionally	—	—	—	—	—	—	—	—	—	—	—	—	56.8	61.9	65.8	71.1			
Take cocaine powder regularly	—	—	—	—	—	—	—	—	—	—	—	—	81.4	82.9	83.9	90.2			
Try heroin once or twice	60.1	58.9	55.8	52.9	50.4	52.1	52.9	51.1	50.8	49.8	47.3	45.8	53.6	54.0	53.8	55.4			
Take heroin occasionally	75.6	75.6	71.9	71.4	70.9	70.9	72.2	69.8	71.8	70.7	69.8	68.2	74.6	73.8	75.5	76.6			
Take heroin regularly	87.2	88.6	86.1	86.6	87.5	86.2	87.5	86.0	86.1	87.2	86.0	87.1	88.7	88.8	89.5	90.2			
Try heroin once or twice without using a needle	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Take heroin occasionally without using a needle	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Try any narcotic other than heroin (codeine, Vicodin, OxyContin, Percocet, etc.) once or twice	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Take any narcotic other than heroin occasionally	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Take any narcotic other than heroin regularly	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Try amphetamines once or twice ^d	35.4	33.4	30.8	29.9	29.7	29.7	26.4	25.3	24.7	25.4	25.2	25.1	29.1	29.6	32.8	32.2			
Take amphetamines regularly ^d	69.0	67.3	66.6	67.1	69.9	69.1	66.1	64.7	64.8	67.1	67.2	67.3	69.4	69.8	71.2	71.2			
Try Adderall once or twice ^e	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Try Adderall occasionally ^e	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Try crystal methamphetamine (ice) once or twice	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Try bath salts (synthetic stimulants) once or twice	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Take bath salts (synthetic stimulants) occasionally	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Try sedatives (barbiturates) once or twice ^f	34.8	32.5	31.2	31.3	30.7	30.9	28.4	27.5	27.0	27.4	26.1	25.4	30.9	29.7	32.2	32.4			
Take sedatives (barbiturates) regularly ^f	69.1	67.7	68.6	68.4	71.6	72.2	69.9	67.6	67.7	68.5	68.3	67.2	69.4	69.6	70.5	70.2			
Try one or two drinks of an alcoholic beverage (beer, wine, liquor)	5.3	4.8	4.1	3.4	4.1	3.8	4.6	3.5	4.2	4.6	5.0	4.6	6.2	6.0	6.0	8.3			
Take one or two drinks nearly every day	21.5	21.2	18.5	19.6	22.6	20.3	21.6	21.6	21.6	23.0	24.4	25.1	26.2	27.3	28.5	31.3			
Take four or five drinks nearly every day	63.5	61.0	62.9	63.1	66.2	65.7	64.5	65.5	66.8	68.4	69.8	66.5	69.7	68.5	69.8	70.9			
Have five or more drinks once or twice each weekend	37.8	37.0	34.7	34.5	34.9	35.9	36.3	36.0	38.6	41.7	43.0	39.1	41.9	42.6	44.0	47.1			
Smoke one or more packs of cigarettes per day	51.3	56.4	58.4	59.0	63.0	63.7	63.3	60.5	61.2	63.8	66.5	66.0	68.6	68.0	67.2	68.2			
Use electronic cigarettes (e-cigarettes) regularly ^g	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Smoke little cigars or cigarillos regularly	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—			
Use smokeless tobacco regularly	—	—	—	—	—	—	—	—	—	—	—	25.8	30.0	33.2	32.9	34.2			
Take steroids	—	—	—	—	—	—	—	—	—	—	—	—	—	—	63.8	69.9			
<i>Approximate weighted N =</i>	<i>2,804</i>	<i>2,918</i>	<i>3,052</i>	<i>3,770</i>	<i>3,250</i>	<i>3,234</i>	<i>3,604</i>	<i>3,557</i>	<i>3,305</i>	<i>3,262</i>	<i>3,250</i>	<i>3,020</i>	<i>3,315</i>	<i>3,276</i>	<i>2,796</i>	<i>2,553</i>			

Table continued on n

(Table continued on next page.)

TABLE 11 (cont.)
Trends in Harmfulness of Drugs as Perceived by 12th Graders

	Percentage saying great risk ^a														
<i>How much do you think people risk harming themselves (physically or in other ways), if they . . .</i>	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Try marijuana once or twice	27.1	24.5	21.9	19.5	16.3	15.6	14.9	16.7	15.7	13.7	15.3	16.1	16.1	15.9	16.1
Smoke marijuana occasionally	40.6	39.6	35.6	30.1	25.6	25.9	24.7	24.4	23.9	23.4	23.5	23.2	26.6	25.4	25.8
Smoke marijuana regularly	78.6	76.5	72.5	65.0	60.8	59.9	58.1	58.5	57.4	58.3	57.4	53.0	54.9	54.6	58.0
Try synthetic marijuana once or twice	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take synthetic marijuana occasionally	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try LSD once or twice	46.6	42.3	39.5	38.8	36.4	36.2	34.7	37.4	34.9	34.3	33.2	36.7	36.2	36.2	36.5
Take LSD regularly	84.3	81.8	79.4	79.1	78.1	77.8	76.6	76.5	76.1	75.9	74.1	73.9	72.3	70.2	69.9
Try PCP once or twice	51.7	54.8	50.8	51.5	49.1	51.0	48.8	46.8	44.8	45.0	46.2	48.3	45.2	47.1	46.6
Try ecstasy (MDMA) once or twice ^b	—	—	—	—	—	—	33.8	34.5	35.0	37.9	45.7	52.2	56.3	57.7	60.1
Try salvia once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take salvia occasionally	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try cocaine once or twice	59.4	56.8	57.6	57.2	53.7	54.2	53.6	54.6	52.1	51.1	50.7	51.2	51.0	50.7	50.5
Take cocaine occasionally	75.5	75.1	73.3	73.7	70.8	72.1	72.4	70.1	70.1	69.5	69.9	68.3	69.1	67.2	66.7
Take cocaine regularly	90.4	90.2	90.1	89.3	87.9	88.3	87.1	86.3	85.8	86.2	84.1	84.5	83.0	82.2	82.8
Try crack once or twice	60.6	62.4	57.6	58.4	54.6	56.0	54.0	52.2	48.2	48.4	49.4	50.8	47.3	47.8	48.4
Take crack occasionally	76.5	76.3	73.9	73.8	72.8	71.4	70.3	68.7	67.3	65.8	65.4	65.6	64.0	64.5	63.8
Take crack regularly	90.1	89.3	87.5	89.6	88.6	88.0	86.2	85.3	85.4	85.3	85.8	84.1	83.2	83.5	83.3
Try cocaine powder once or twice	53.6	57.1	53.2	55.4	52.0	53.2	51.4	48.5	46.1	47.0	49.0	49.5	46.2	45.4	46.2
Take cocaine powder occasionally	69.8	70.8	68.6	70.6	69.1	68.8	67.7	65.4	64.2	64.7	63.2	64.4	61.4	61.6	60.8
Take cocaine powder regularly	88.9	88.4	87.0	88.6	87.8	86.8	86.0	84.1	84.6	85.5	84.4	84.2	82.3	81.7	82.7
Try heroin once or twice	55.2	50.9	50.7	52.8	50.9	52.5	56.7	57.8	56.0	54.2	55.6	56.0	58.0	56.6	55.2
Take heroin occasionally	74.9	74.2	72.0	72.1	71.0	74.8	76.3	76.9	77.3	74.6	75.9	76.6	78.5	75.7	76.0
Take heroin regularly	89.6	89.2	88.3	88.0	87.2	89.5	88.9	89.1	89.9	89.2	88.3	88.5	89.3	86.8	87.5
Try heroin once or twice without using a needle	—	—	—	—	55.6	58.6	60.5	59.6	58.5	61.6	60.7	60.6	58.9	61.2	60.5
Take heroin occasionally without using a needle	—	—	—	—	71.2	71.0	74.3	73.4	73.6	74.7	74.4	74.7	73.0	76.1	73.3
Try any narcotic other than heroin (codeine, Vicodin, OxyContin, Percocet, etc.) once or twice	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take any narcotic other than heroin occasionally	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take any narcotic other than heroin regularly	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try amphetamines once or twice ^d	36.3	32.6	31.3	31.4	28.8	30.8	31.0	35.3	32.2	32.6	34.7	34.4	36.8	35.7	37.7
Take amphetamines regularly ^d	74.1	72.4	69.9	67.0	65.9	66.8	66.0	67.7	66.4	66.3	67.1	64.8	65.6	63.9	67.1
Try Adderall once or twice ^e	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try Adderall occasionally ^e	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try crystal methamphetamine (ice) once or twice	61.6	61.9	57.5	58.3	54.4	55.3	54.4	52.7	51.2	51.3	52.7	53.8	51.2	52.4	54.6
Try bath salts (synthetic stimulants) once or twice	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Take bath salts (synthetic stimulants) occasionally	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Try sedatives (barbiturates) once or twice ^f	35.1	32.2	29.2	29.9	26.3	29.1	26.9	29.0	26.1	25.0	25.7	26.2	27.9†	24.9	24.7
Take sedatives (barbiturates) regularly ^f	70.5	70.2	66.1	63.3	61.6	60.4	56.8	56.3	54.1	52.3	50.3	49.3	49.6†	54.0	54.1
Try one or two drinks of an alcoholic beverage (beer, wine, liquor)	9.1	8.6	8.2	7.6	5.9	7.3	6.7	8.0	8.3	6.4	8.7	7.6	8.4	8.6	8.5
Take one or two drinks nearly every day	32.7	30.6	28.2	27.0	24.8	25.1	24.8	24.3	21.8	21.7	23.4	21.0	20.1	23.0	23.7
Take four or five drinks nearly every day	69.5	70.5	67.8	66.2	62.8	65.6	63.0	62.1	61.1	59.9	60.7	58.8	57.8	59.2	61.8
Have five or more drinks once or twice each weekend	48.6	49.0	48.3	46.5	45.2	49.5	43.0	42.8	43.1	42.7	43.6	42.2	43.5	43.6	45.0
Smoke one or more packs of cigarettes per day	69.4	69.2	69.5	67.6	65.6	68.2	68.7	70.8	70.8	73.1	73.3	74.2	72.1	74.0	76.5
Use electronic cigarettes (e-cigarettes) regularly ^g	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Smoke little cigars or cigarillos regularly	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Use smokeless tobacco regularly	37.4	35.5	38.9	36.6	33.2	37.4	38.6	40.9	41.1	42.2	45.4	42.6	43.3	45.0	43.6
Take steroids	65.6	70.7	69.1	66.1	66.4	67.6	67.2	68.1	62.1	57.9	58.9	57.1	55.0	55.7	56.8
Approximate weighted N =	2,549	2,684	2,759	2,591	2,603	2,449	2,579	2,564	2,306	2,130	2,173	2,198	2,466	2,491	2,512

Table continued on next pag

(Table continued on next page.)

TABLE 11 (cont.)
Trends in Harmfulness of Drugs as Perceived by 12th Graders

	Percentage saying great risk ^a											2015– 2016
<i>How much do you think people risk harming themselves (physically or in other ways), if they . . .</i>	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	change
Try marijuana once or twice	17.8	18.6	17.4	18.5	17.1	15.6	14.8	14.5	12.5	12.3	12.9	+0.7
Smoke marijuana occasionally	25.9	27.1	25.8	27.4	24.5	22.7	20.6	19.5	16.4	15.8	17.1	+1.3
Smoke marijuana regularly	57.9	54.8	51.7	52.4	46.8	45.7	44.1	39.5	36.1	31.9	31.1	-0.8
Try synthetic marijuana once or twice	—	—	—	—	—	—	23.5	25.9	32.5	33.0	35.6	+2.6
Take synthetic marijuana occasionally	—	—	—	—	—	—	32.7	36.2	39.4	40.9	43.9	+3.0
Try LSD once or twice	36.1	37.0	33.9	37.1	35.6	34.7	33.1	34.9	35.5	33.2	31.7	-1.5
Take LSD regularly	69.3	67.3	63.6	67.8	65.3	65.5	66.8	66.8	62.7	60.7	58.2	-2.5
Try PCP once or twice	47.0	48.0	47.4	49.7	52.4	53.9	51.6	53.9	53.8	54.4	55.1	+0.7
Try ecstasy (MDMA) once or twice ^b	59.3	58.1	57.0	53.3	50.6	49.0	49.4	47.5†	47.8	49.5	48.8	-0.7
Try salvia once or twice ^c	—	—	—	—	39.8	36.7‡	13.8	12.9	14.1	13.1	13.0	-0.1
Take salvia occasionally	—	—	—	—	—	—	23.1	21.3	20.0	17.6	16.3	-1.3
Try cocaine once or twice	52.5	51.3	50.3	53.1	52.8	54.0	51.6	54.4	53.7	51.1	52.7	+1.5
Take cocaine occasionally	69.8	68.8	67.1	71.4	67.8	69.7	69.0	70.2	68.1	66.3	68.6	+2.3
Take cocaine regularly	84.6	83.3	80.7	84.4	81.7	83.8	82.6	83.3	80.6	79.1	78.3	-0.7
Try crack once or twice	47.8	47.3	47.5	48.4	50.2	51.7	52.0	55.6	54.5	53.6	53.9	+0.3
Take crack occasionally	64.8	63.6	65.2	64.7	64.3	66.2	66.5	69.5	68.5	67.8	66.2	-1.6
Take crack regularly	82.8	82.6	83.4	84.0	83.8	83.9	84.0	85.4	82.0	81.2	81.9	+0.7
Try cocaine powder once or twice	45.8	45.1	45.1	46.5	48.2	48.0	48.1	49.9	49.9	49.0	49.3	+0.2
Take cocaine powder occasionally	61.9	59.9	61.6	62.6	62.6	64.2	62.6	65.4	64.8	62.8	62.9	+0.1
Take cocaine powder regularly	82.1	81.5	82.5	83.4	81.8	83.3	83.3	83.9	81.5	80.1	80.7	+0.6
Try heroin once or twice	59.1	58.4	55.5	59.3	58.3	59.1	59.4	61.7	62.8	64.0	64.5	+0.5
Take heroin occasionally	79.1	76.2	75.3	79.7	74.8	77.2	78.0	78.2	77.9	78.0	78.7	+0.7
Take heroin regularly	89.7	87.8	86.4	89.9	85.5	87.9	88.6	87.6	85.7	84.8	85.4	+0.6
Try heroin once or twice without using a needle	62.6	60.2	60.8	61.5	63.8	61.1	63.3	64.5	65.3	62.5	66.1	+3.5
Take heroin occasionally without using a needle	76.2	73.9	73.2	74.8	76.2	74.7	76.1	76.4	73.6	71.1	74.6	+3.5
Try any narcotic other than heroin (codeine, Vicodin, OxyContin, Percocet, etc.) once or twice	—	—	—	—	40.4	39.9	38.4	43.1	42.7	44.1	43.6	-0.5
Take any narcotic other than heroin occasionally	—	—	—	—	54.3	54.8	53.8	57.3	59.0	58.5	55.7	-2.8
Take any narcotic other than heroin regularly	—	—	—	—	74.9	75.5	73.9	75.8	72.7	73.9	72.4	-1.5
Try amphetamines once or twice ^d	39.5	41.3	39.2	41.9	40.6‡	34.8	34.3	36.3	34.1	34.0	31.1	-2.9
Take amphetamines regularly ^d	68.1	68.1	65.4	69.0	63.6‡	58.7	60.0	59.5	55.1	54.3	51.3	-3.0
Try Adderall once or twice ^e	—	—	—	—	33.3	31.2	27.2	31.8	33.6	34.3	32.5	-1.8
Try Adderall occasionally ^e	—	—	—	—	41.6	40.8	35.3	38.8	41.5	41.6	40.9	-0.8
Try crystal methamphetamine (ice) once or twice	59.1	60.2	62.2	63.4	64.9	66.5	67.8	72.2	70.2	70.0	70.0	0.0
Try bath salts (synthetic stimulants) once or twice	—	—	—	—	—	—	33.2	59.5	59.2	57.5	54.9	-2.6
Take bath salts (synthetic stimulants) occasionally	—	—	—	—	—	—	45.0	69.9	68.8	67.4	64.2	-3.3
Try sedatives (barbiturates) once or twice ^f	28.0	27.9	25.9	29.6	28.0	27.8	27.8	29.4	29.6	28.9	27.4	-1.5
Take sedatives (barbiturates) regularly ^f	56.8	55.1	50.2	54.7	52.1	52.4	53.9	53.3	50.5	50.6	47.0	-3.6
Try one or two drinks of an alcoholic beverage (beer, wine, liquor)	9.3	10.5	10.0	9.4	10.8	9.4	8.7	9.9	8.6	10.3	9.5	-0.8
Take one or two drinks nearly every day	25.3	25.1	24.2	23.7	25.4	24.6	23.7	23.1	21.1	21.5	21.6	+0.1
Take four or five drinks nearly every day	63.4	61.8	60.8	62.4	61.1	62.3	63.6	62.4	61.2	59.1	59.1	0.0
Have five or more drinks once or twice each weekend	47.6	45.8	46.3	48.0	46.3	47.6	48.8	45.8	45.4	46.9	48.4	+1.4
Smoke one or more packs of cigarettes per day	77.6	77.3	74.0	74.9	75.0	77.7	78.2	78.2	78.0	75.9	76.5	+0.6
Use electronic cigarettes (e-cigarettes) regularly ^g	—	—	—	—	—	—	—	—	14.2	16.2	18.2	+2.1
Smoke little cigars or cigarillos regularly	—	—	—	—	—	—	—	—	38.3	39.7	39.5	-0.2
Use smokeless tobacco regularly	45.9	44.0	42.9	40.8	41.2	42.6	44.3	41.6	40.7	38.5	38.1	-0.4
Take steroids	60.2	57.4	60.8	60.2	59.2	61.1	58.6	54.2	54.6	54.4	54.5	+0.1
<i>Approximate weighted N =</i>	<i>2,407</i>	<i>2,450</i>	<i>2,389</i>	<i>2,290</i>	<i>2,440</i>	<i>2,408</i>	<i>2,331</i>	<i>2,098</i>	<i>2,067</i>	<i>2,174</i>	<i>1,988</i>	

Table continued on next page.

(Table continued on next page.)

TABLE 11 (cont.)
Trends in Harmfulness of Drugs as Perceived by 12th Graders

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. '‡' indicates that the question changed the following year. See relevant footnote for that drug. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aAnswer alternatives were: (1) No risk, (2) Slight risk, (3) Moderate risk, (4) Great risk, and (5) Can't say, drug unfamiliar.

^bBeginning in 2014 data are based on the revised question which included "Molly." 2014 and 2015 data are not comparable to earlier years due to the revision of the question text.

^cIn 2011 the question on perceived risk of using salvia once or twice appeared at the end of a form. In 2012 the question was moved to an earlier section of the same form. A question on perceived risk of using salvia occasionally was also added following the question on perceived risk of trying salvia once or twice. These changes likely explain the discontinuity in the 2012 results.

^dIn 2011 the list of examples was changed from uppers, pep pills, bennies, speed to uppers, speed, Adderall, Ritalin, etc. These changes likely explain the discontinuity in the 2011 results.

^eIn 2014 "(without a doctor's orders)" added to the questions on perceived risk of using Adderall.

^fIn 2004 the question text was changed from barbiturates to sedatives/barbiturates and the list of examples was changed from downers, goofballs, reds, yellows, etc. to just downers. These changes likely explain the discontinuity in the 2004 results.

^gBased on two of six forms; N is two times the N indicated.

TABLE 12
Trends in Disapproval of Drug Use in Grade 8

	Percentage who disapprove or strongly disapprove ^a														
<i>Do you disapprove of people who . . .</i>	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
Try marijuana once or twice ^b	84.6	82.1	79.2	72.9	70.7	67.5	67.6	69.0	70.7	72.5	72.4	73.3	73.8	75.9	75.3
Smoke marijuana occasionally ^b	89.5	88.1	85.7	80.9	79.7	76.5	78.1	78.4	79.3	80.6	80.6	80.9	81.5	83.1	82.4
Smoke marijuana regularly ^b	92.1	90.8	88.9	85.3	85.1	82.8	84.6	84.5	84.5	85.3	84.5	85.3	85.7	86.8	86.3
Try inhalants once or twice ^c	84.9	84.0	82.5	81.6	81.8	82.9	84.1	83.0	85.2	85.4	86.6	86.1	85.1	85.1	84.6
Take inhalants regularly ^c	90.6	90.0	88.9	88.1	88.8	89.3	90.3	89.5	90.3	90.2	90.5	90.4	89.8	90.1	89.8
Take LSD once or twice ^d	—	—	77.1	75.2	71.6	70.9	72.1	69.1	69.4	66.7	64.6	62.6	61.0	58.1	58.5
Take LSD regularly ^d	—	—	79.8	78.4	75.8	75.3	76.3	72.5	72.5	69.3	67.0	65.5	63.5	60.5	60.7
Try ecstasy (MDMA) once or twice ^e	—	—	—	—	—	—	—	—	—	—	69.0	74.3	77.7	76.3	75.0
Take ecstasy (MDMA) occasionally ^e	—	—	—	—	—	—	—	—	—	—	73.6	78.6	81.3	79.4	77.9
Try crack once or twice ^c	91.7	90.7	89.1	86.9	85.9	85.0	85.7	85.4	86.0	85.4	86.0	86.2	86.4	87.4	87.6
Take crack occasionally ^c	93.3	92.5	91.7	89.9	89.8	89.3	90.3	89.5	89.9	88.8	89.8	89.6	89.8	90.3	90.5
Try cocaine powder once or twice ^c	91.2	89.6	88.5	86.1	85.3	83.9	85.1	84.5	85.2	84.8	85.6	85.8	85.6	86.8	87.0
Take cocaine powder occasionally ^c	93.1	92.4	91.6	89.7	89.7	88.7	90.1	89.3	89.9	88.8	89.6	89.9	89.8	90.3	90.7
Try heroin once or twice without using a needle ^d	—	—	—	—	85.8	85.0	87.7	87.3	88.0	87.2	87.2	87.8	86.9	86.6	86.9
Take heroin occasionally without using a needle ^d	—	—	—	—	88.5	87.7	90.1	89.7	90.2	88.9	88.9	89.6	89.0	88.6	88.5
Try one or two drinks of an alcoholic beverage (beer, wine, liquor) ^b	51.7	52.2	50.9	47.8	48.0	45.5	45.7	47.5	48.3	48.7	49.8	51.1	49.7	51.1	51.2
Take one or two drinks nearly every day ^b	82.2	81.0	79.6	76.7	75.9	74.1	76.6	76.9	77.0	77.8	77.4	78.3	77.1	78.6	78.7
Have five or more drinks once or twice each weekend ^b	85.2	83.9	83.3	80.7	80.7	79.1	81.3	81.0	80.3	81.2	81.6	81.9	81.9	82.3	82.9
Smoke one to five cigarettes per day ^e	—	—	—	—	—	—	—	—	75.1	79.1	80.4	81.1	81.4	83.1	82.9
Smoke one or more packs of cigarettes per day ^f	82.8	82.3	80.6	78.4	78.6	77.3	80.3	80.0	81.4	81.9	83.5	84.6	84.6	85.7	85.3
Use electronic cigarettes (e-cigarettes) regularly ^e	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Use smokeless tobacco regularly ^b	79.1	77.2	77.1	75.1	74.0	74.1	76.5	76.3	78.0	79.2	79.4	80.6	80.7	81.0	82.0
Take steroids ^g	89.8	90.3	89.9	87.9	—	—	—	—	—	—	—	—	—	—	—
<i>Approximate weighted N =</i>	<i>17,400</i>	<i>18,500</i>	<i>18,400</i>	<i>17,400</i>	<i>17,600</i>	<i>18,000</i>	<i>18,800</i>	<i>18,100</i>	<i>16,700</i>	<i>16,700</i>	<i>16,200</i>	<i>15,100</i>	<i>16,500</i>	<i>17,000</i>	<i>16,800</i>

(Table continued on next page.)

Table continued on next page.

TABLE 12 (cont.)
Trends in Disapproval of Drug Use in Grade 8

Do you disapprove of people who . . .	Percentage who disapprove or strongly disapprove ^a											2015–2016 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Try marijuana once or twice ^b	76.0	78.7	76.6	75.3	73.5	74.4	75.1	72.0	70.5	70.3	70.1	-0.2
Smoke marijuana occasionally ^b	82.2	84.5	82.6	81.9	79.9	81.1	81.6	78.8	77.7	77.5	77.5	0.0
Smoke marijuana regularly ^b	86.1	87.7	86.8	85.9	84.3	85.7	85.6	83.8	82.2	82.2	82.3	+0.1
Try inhalants once or twice ^c	83.4	84.1	82.3	83.1	83.1	82.9	83.1	81.6	80.7	80.6	78.3	-2.4 s
Take inhalants regularly ^c	89.0	89.5	88.5	88.4	88.9	88.5	88.6	86.8	85.5	85.4	83.3	-2.1 s
Take LSD once or twice ^d	53.9	53.5	52.6	53.2	53.7	55.4	51.8	52.0	52.8	56.0	55.2	-0.8
Take LSD regularly ^d	55.8	55.6	54.7	55.7	55.8	57.6	54.1	53.6	54.8	58.1	57.6	-0.5
Try ecstasy (MDMA) once or twice ^e	66.7	65.7	63.5	62.3	62.4	64.2	60.2	60.9	61.0†	68.2	64.8	-3.4 s
Take ecstasy (MDMA) occasionally ^e	69.8	68.3	66.5	65.7	65.9	67.5	63.2	63.4	64.1‡	71.7	67.5	-4.1 ss
Try crack once or twice ^c	87.2	88.6	87.2	88.4	89.1	88.5	89.0	88.1	88.0	87.5	87.0	-0.5
Take crack occasionally ^c	90.0	91.2	90.3	91.0	91.5	91.0	91.2	90.3	89.8	89.8	88.8	-1.0
Try cocaine powder once or twice ^c	86.5	88.2	86.8	88.1	88.4	88.3	88.6	88.0	87.7	87.5	86.8	-0.7
Take cocaine powder occasionally ^c	90.2	91.0	90.1	90.7	91.4	91.3	91.5	90.6	90.1	90.1	89.3	-0.7
Try heroin once or twice without using a needle ^d	87.2	88.4	86.9	88.6	89.5	87.5	86.8	87.2	87.1	87.1	85.6	-1.5
Take heroin occasionally without using a needle ^d	88.5	89.7	88.2	90.1	90.6	89.0	87.7	88.2	88.1	88.0	86.7	-1.3
Try one or two drinks of an alcoholic beverage (beer, wine, liquor) ^b	51.3	54.0	52.5	52.7	54.2	54.0	54.1	53.3	53.3	53.7	52.6	-1.2
Take one or two drinks nearly every day ^b	78.7	80.4	79.2	78.5	79.5	80.7	81.3	80.2	79.6	79.7	79.1	-0.5
Have five or more drinks once or twice each weekend ^b	82.0	83.8	83.2	83.2	83.6	84.8	86.0	85.0	84.9	85.4	84.9	-0.5
Smoke one to five cigarettes per day ^e	83.5	85.3	85.0	83.6	84.7	86.8	—	—	—	—	—	—
Smoke one or more packs of cigarettes per day ^f	85.6	87.0	86.7	87.1	87.0	88.0	88.8	88.0	87.5	88.8	88.1	-0.7
Use electronic cigarettes (e-cigarettes) regularly ^e	—	—	—	—	—	—	—	—	58.4	65.0	66.6	+1.6
Use smokeless tobacco regularly ^b	81.0	82.3	82.1	81.5	81.2	82.6	82.7	81.5	80.2	82.5	81.1	-1.5
Take steroids ^g	—	—	—	—	—	—	—	—	—	—	—	—
Approximate weighted N = 16,500 16,100 15,700 15,000 15,300 16,000 15,100 14,600 14,600 14,400 16,900												

(Table continued on next page.)

Table continued on next page.

TABLE 12 (cont.)
Trends in Disapproval of Drug Use in Grade 8

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$. ' — ' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. ' ‡ ' indicates that the question changed the following year.

^aAnswer alternatives were: (1) Don't disapprove, (2) Disapprove, (3) Strongly disapprove, and (4) Can't say, drug unfamiliar. Percentages are shown for categories (2) and (3) combined.

^bBeginning in 2012, data based on two thirds of N indicated.

^cBeginning in 1997, data based on two thirds of N indicated due to changes in questionnaire forms.

^dData based on one of two forms in 1993–1996; N is one half of N indicated. Beginning in 1997, data based on one third of N indicated due to changes in questionnaire forms.

^eData based on one third of N indicated. For MDMA "Molly" was added to the question text in 2015; 2014 and 2015 data are not comparable due to this change.

^fBeginning in 1999, data based on two thirds of N indicated due to changes in questionnaire forms.

^gData based on two forms in 1991 and 1992. Data based on one of two forms in 1993 and 1994; N is one half of N indicated.

TABLE 13
Trends in Disapproval of Drug Use in Grade 10

Do you disapprove of people who . . .	Percentage who disapprove or strongly disapprove ^a														
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005
Try marijuana once or twice ^b	74.6	74.8	70.3	62.4	59.8	55.5	54.1	56.0	56.2	54.9	54.8	57.8	58.1	60.4	61.3
Smoke marijuana occasionally ^b	83.7	83.6	79.4	72.3	70.0	66.9	66.2	67.3	68.2	67.2	66.2	68.3	68.4	70.8	71.9
Smoke marijuana regularly ^b	90.4	90.0	87.4	82.2	81.1	79.7	79.7	80.1	79.8	79.1	78.0	78.6	78.8	81.3	82.0
Try inhalants once or twice ^c	85.2	85.6	84.8	84.9	84.5	86.0	86.9	85.6	88.4	87.5	87.8	88.6	87.7	88.5	88.1
Take inhalants regularly ^c	91.0	91.5	90.9	91.0	90.9	91.7	91.7	91.1	92.4	91.8	91.3	91.8	91.0	92.3	91.9
Take LSD once or twice ^d	—	—	82.1	79.3	77.9	76.8	76.6	76.7	77.8	77.0	75.4	74.6	74.4	72.4	71.8
Take LSD regularly ^d	—	—	86.8	85.6	84.8	84.5	83.4	82.9	84.3	82.1	80.8	79.4	77.6	75.9	75.0
Try ecstasy (MDMA) once or twice ^e	—	—	—	—	—	—	—	—	—	—	72.6	77.4	81.0	83.7	83.1
Take ecstasy (MDMA) occasionally ^e	—	—	—	—	—	—	—	—	—	—	81.0	84.6	86.3	88.0	87.4
Try crack once or twice ^c	92.5	92.5	91.4	89.9	88.7	88.2	87.4	87.1	87.8	87.1	86.9	88.0	87.6	88.6	88.8
Take crack occasionally ^c	94.3	94.4	93.6	92.5	91.7	91.9	91.0	90.6	91.5	90.9	90.6	91.0	91.0	91.8	91.8
Try cocaine powder once or twice ^c	90.8	91.1	90.0	88.1	86.8	86.1	85.1	84.9	86.0	84.8	85.3	86.4	85.9	86.8	86.9
Take cocaine powder occasionally ^c	94.0	94.0	93.2	92.1	91.4	91.1	90.4	89.7	90.7	89.9	90.2	89.9	90.4	91.2	91.2
Try heroin once or twice without using a needle ^d	—	—	—	—	89.7	89.5	89.1	88.6	90.1	90.1	89.1	89.2	89.3	90.1	90.3
Take heroin occasionally without using a needle ^d	—	—	—	—	91.6	91.7	91.4	90.5	91.8	92.3	90.8	90.7	90.6	91.8	92.0
Try one or two drinks of an alcoholic beverage (beer, wine, liquor) ^b	37.6	39.9	38.5	36.5	36.1	34.2	33.7	34.7	35.1	33.4	34.7	37.7	36.8	37.6	38.5
Take one or two drinks nearly every day ^b	81.7	81.7	78.6	75.2	75.4	73.8	75.4	74.6	75.4	73.8	73.8	74.9	74.2	75.1	76.9
Have five or more drinks once or twice each weekend ^b	76.7	77.6	74.7	72.3	72.2	70.7	70.2	70.5	69.9	68.2	69.2	71.5	71.6	71.8	73.7
Smoke one to five cigarettes per day ^e	—	—	—	—	—	—	—	—	67.8	69.1	71.2	74.3	76.2	77.5	79.3
Smoke one or more packs of cigarettes per day ^f	79.4	77.8	76.5	73.9	73.2	71.6	73.8	75.3	76.1	76.7	78.2	80.6	81.4	82.7	84.3
Use electronic cigarettes (e-cigarettes) regularly ^e	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Use smokeless tobacco regularly ^b	75.4	74.6	73.8	71.2	71.0	71.0	72.3	73.2	75.1	75.8	76.1	78.7	79.4	80.2	80.5
Take steroids ^g	90.0	91.0	91.2	90.8	—	—	—	—	—	—	—	—	—	—	—
Approximate weighted N = 14,800 14,800 15,300 15,900 17,000 15,700 15,600 15,000 13,600 14,300 14,000 14,300 15,800 16,400 16,200															

Table continued on next page.

TABLE 13 (cont.)
Trends in Disapproval of Drug Use in Grade 10

Do you disapprove of people who . . .	Percentage who disapprove or strongly disapprove ^a											2015–2016 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Try marijuana once or twice ^b	62.5	63.9	64.5	60.1	59.2	58.5	56.2	53.2	53.8	52.7	52.6	0.0
Smoke marijuana occasionally ^b	72.6	73.3	73.6	69.2	68.0	67.9	65.7	62.1	62.9	62.6	61.9	-0.7
Smoke marijuana regularly ^b	82.5	82.4	83.0	79.9	78.7	78.8	77.3	73.8	74.6	74.3	73.5	-0.8
Try inhalants once or twice ^c	88.1	87.6	87.1	87.0	86.5	86.9	85.7	86.1	85.9	84.1	83.3	-0.8
Take inhalants regularly ^c	92.2	91.8	91.6	91.1	90.8	90.9	90.0	89.7	89.7	88.3	87.1	-1.2
Take LSD once or twice ^d	71.2	67.7	66.3	67.8	68.2	68.5	68.3	69.1	67.8	70.3	69.5	-0.7
Take LSD regularly ^d	74.9	71.5	69.8	72.2	72.9	72.5	73.0	74.2	73.3	76.5	74.9	-1.7
Try ecstasy (MDMA) once or twice ^e	81.6	80.0	78.1	76.5	75.5	76.1	75.3	75.4	74.4†	78.0	76.8	-1.2
Take ecstasy (MDMA) occasionally ^e	86.0	84.3	83.0	81.3	81.3	82.2	81.2	81.3	80.4†	84.0	81.7	-2.3 s
Try crack once or twice ^c	89.5	89.5	90.8	90.4	90.3	90.9	91.0	90.6	90.6	90.1	89.7	-0.4
Take crack occasionally ^c	92.0	92.7	92.9	92.8	92.4	93.0	93.0	92.4	92.4	92.1	91.1	-1.1
Try cocaine powder once or twice ^c	87.3	87.7	88.6	88.4	89.0	89.4	89.3	88.7	88.9	87.9	87.9	0.0
Take cocaine powder occasionally ^c	91.4	92.0	92.1	92.1	92.2	92.5	92.4	91.8	91.9	91.8	90.8	-1.1
Try heroin once or twice without using a needle ^d	91.1	90.7	91.4	91.6	91.4	91.6	91.9	91.3	91.9	91.7	90.2	-1.5
Take heroin occasionally without using a needle ^d	92.5	92.5	92.5	93.0	92.4	92.4	92.9	92.3	92.7	92.7	90.9	-1.8 s
Try one or two drinks of an alcoholic beverage (beer, wine, liquor) ^b	37.8	39.5	41.8	39.7	40.3	41.5	39.6	38.5	40.7	40.0	41.8	+1.8
Take one or two drinks nearly every day ^b	76.4	77.1	79.1	77.6	77.6	80.0	78.0	77.1	77.9	78.2	78.6	+0.4
Have five or more drinks once or twice each weekend ^b	72.9	74.1	77.2	75.1	75.9	77.3	77.5	77.8	79.5	79.6	80.8	+1.2
Smoke one to five cigarettes per day ^e	80.2	79.7	82.5	80.0	80.6	82.1	—	—	—	—	—	—
Smoke one or more packs of cigarettes per day ^f	83.2	84.7	85.2	84.5	83.9	85.8	86.0	86.1	88.0	88.3	88.5	+0.3
Use electronic cigarettes (e-cigarettes) regularly ^e	—	—	—	—	—	—	—	—	54.6	59.9	65.0	+5.1 sss
Use smokeless tobacco regularly ^b	80.5	80.9	81.8	79.5	78.5	79.5	79.5	77.7	78.7	80.1	81.2	+1.1
Take steroids ^g	—	—	—	—	—	—	—	—	—	—	—	—
Approximate weighted N = 16,200 16,100 15,100 15,900 15,200 14,900 15,000 12,900 13,000 15,600 14,700												

Table continued on next page.

TABLE 13 (cont.)
Trends in Disapproval of Drug Use in Grade 10

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: $s = .05$, $ss = .01$, $sss = .001$. '—' indicates data not available. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding. '‡' indicates that the question changed the following year.

^aAnswer alternatives were: (1) Don't disapprove, (2) Disapprove, (3) Strongly disapprove, and (4) Can't say, drug unfamiliar. Percentages are shown for categories (2) and (3) combined.

^bBeginning in 2012, data based on two thirds of N indicated.

^cBeginning in 1997, data based on two thirds of N indicated due to changes in questionnaire forms.

^dData based on one of two forms in 1993–1996; N is one half of N indicated. Beginning in 1997, data based on one third of N indicated due to changes in questionnaire forms.

^eData based on one third of N indicated. For MDMA "Molly" was added to the question text in 2015; 2014 and 2015 data are not comparable due to this change.

^fBeginning in 1999, data based on two thirds of N indicated due to changes in questionnaire forms.

^gData based on two forms in 1991 and 1992. Data based on one of two forms in 1993 and 1994; N is one half of N indicated.

TABLE 14
Trends in Disapproval of Drug Use in Grade 12

Percentage who disapprove or strongly disapprove ^b

Do you disapprove of people (who are 18 or older) doing each of the following?^a

	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990
Trying marijuana once or twice	47.0	38.4	33.4	33.4	34.2	39.0	40.0	45.5	46.3	49.3	51.4	54.6	56.6	60.8	64.6	67.8
Smoking marijuana occasionally	54.8	47.8	44.3	43.5	45.3	49.7	52.6	59.1	60.7	63.5	65.8	69.0	71.6	74.0	77.2	80.5
Smoking marijuana regularly	71.9	69.5	65.5	67.5	69.2	74.6	77.4	80.6	82.5	84.7	85.5	86.6	89.2	89.3	89.8	91.0
Trying LSD once or twice	82.8	84.6	83.9	85.4	86.6	87.3	86.4	88.8	89.1	88.9	89.5	89.2	91.6	89.8	89.7	89.8
Taking LSD regularly	94.1	95.3	95.8	96.4	96.9	96.7	96.8	96.7	97.0	96.8	97.0	96.6	97.8	96.4	96.4	96.3
Trying ecstasy (MDMA) once or twice ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Trying cocaine once or twice	81.3	82.4	79.1	77.0	74.7	76.3	74.6	76.6	77.0	79.7	79.3	80.2	87.3	89.1	90.5	91.5
Taking cocaine regularly	93.3	93.9	92.1	91.9	90.8	91.1	90.7	91.5	93.2	94.5	93.8	94.3	96.7	96.2	96.4	96.7
Trying crack once or twice	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	92.3
Taking crack occasionally	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	94.3
Taking crack regularly	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	94.9
Trying cocaine powder once or twice	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	87.9
Taking cocaine powder occasionally	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	92.1
Taking cocaine powder regularly	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	93.7
Trying heroin once or twice	91.5	92.6	92.5	92.0	93.4	93.5	93.5	94.6	94.3	94.0	94.0	93.3	96.2	95.0	95.4	95.1
Taking heroin occasionally	94.8	96.0	96.0	96.4	96.8	96.7	97.2	96.9	96.9	97.1	96.8	96.6	97.9	96.9	97.2	96.7
Taking heroin regularly	96.7	97.5	97.2	97.8	97.9	97.6	97.8	97.5	97.7	98.0	97.6	97.6	98.1	97.2	97.4	97.5
Trying heroin once or twice without using a needle	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Taking heroin occasionally without using a needle	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Trying amphetamines once or twice ^d	74.8	75.1	74.2	74.8	75.1	75.4	71.1	72.6	72.3	72.8	74.9	76.5	80.7	82.5	83.3	85.3
Taking amphetamines regularly ^d	92.1	92.8	92.5	93.5	94.4	93.0	91.7	92.0	92.6	93.6	93.3	93.5	95.4	94.2	94.2	95.5
Trying sedatives (barbiturates) once or twice ^e	77.7	81.3	81.1	82.4	84.0	83.9	82.4	84.4	83.1	84.1	84.9	86.8	89.6	89.4	89.3	90.5
Taking sedatives (barbiturates) regularly ^e	93.3	93.6	93.0	94.3	95.2	95.4	94.2	94.4	95.1	95.1	95.5	94.9	96.4	95.3	95.3	96.4
Trying one or two drinks of an alcoholic beverage (beer, wine, liquor)	21.6	18.2	15.6	15.6	15.8	16.0	17.2	18.2	18.4	17.4	20.3	20.9	21.4	22.6	27.3	29.4
Taking one or two drinks nearly every day	67.6	68.9	66.8	67.7	68.3	69.0	69.1	69.9	68.9	72.9	70.9	72.8	74.2	75.0	76.5	77.9
Taking four or five drinks nearly every day	88.7	90.7	88.4	90.2	91.7	90.8	91.8	90.9	90.0	91.0	92.0	91.4	92.2	92.8	91.6	91.9
Having five or more drinks once or twice each weekend	60.3	58.6	57.4	56.2	56.7	55.6	55.5	58.8	56.6	59.6	60.4	62.4	62.0	65.3	66.5	68.9
Smoking one or more packs of cigarettes per day	67.5	65.9	66.4	67.0	70.3	70.8	69.9	69.4	70.8	73.0	72.3	75.4	74.3	73.1	72.4	72.8
Taking steroids	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	90.8
<i>Approximate weighted N =</i>	<i>2,677</i>	<i>2,957</i>	<i>3,085</i>	<i>3,686</i>	<i>3,221</i>	<i>3,261</i>	<i>3,610</i>	<i>3,651</i>	<i>3,341</i>	<i>3,254</i>	<i>3,265</i>	<i>3,113</i>	<i>3,302</i>	<i>3,311</i>	<i>2,799</i>	<i>2,566</i>

Table continued on next page.

TABLE 14 (cont.)
Trends in Disapproval of Drug Use in Grade 12

Percentage who disapprove or strongly disapprove ^b

<i>Do you disapprove of people (who are 18 or older) doing each of the following?^a</i>	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
Trying marijuana once or twice	68.7	69.9	63.3	57.6	56.7	52.5	51.0	51.6	48.8	52.5	49.1	51.6	53.4	52.7	55.0
Smoking marijuana occasionally	79.4	79.7	75.5	68.9	66.7	62.9	63.2	64.4	62.5	65.8	63.2	63.4	64.2	65.4	67.8
Smoking marijuana regularly	89.3	90.1	87.6	82.3	81.9	80.0	78.8	81.2	78.6	79.7	79.3	78.3	78.7	80.7	82.0
Trying LSD once or twice	90.1	88.1	85.9	82.5	81.1	79.6	80.5	82.1	83.0	82.4	81.8	84.6	85.5	87.9	87.9
Taking LSD regularly	96.4	95.5	95.8	94.3	92.5	93.2	92.9	93.5	94.3	94.2	94.0	94.0	94.4	94.6	95.6
Trying ecstasy (MDMA) once or twice ^c	—	—	—	—	—	—	82.2	82.5	82.1	81.0	79.5	83.6	84.7	87.7	88.4
Trying cocaine once or twice	93.6	93.0	92.7	91.6	90.3	90.0	88.0	89.5	89.1	88.2	88.1	89.0	89.3	88.6	88.9
Taking cocaine regularly	97.3	96.9	97.5	96.6	96.1	95.6	96.0	95.6	94.9	95.5	94.9	95.0	95.8	95.4	96.0
Trying crack once or twice	92.1	93.1	89.9	89.5	91.4	87.4	87.0	86.7	87.6	87.5	87.0	87.8	86.6	86.9	86.7
Taking crack occasionally	94.2	95.0	92.8	92.8	94.0	91.2	91.3	90.9	92.3	91.9	91.6	91.5	90.8	92.1	91.9
Taking crack regularly	95.0	95.5	93.4	93.1	94.1	93.0	92.3	91.9	93.2	92.8	92.2	92.4	91.2	93.1	92.1
Trying cocaine powder once or twice	88.0	89.4	86.6	87.1	88.3	83.1	83.0	83.1	84.3	84.1	83.3	83.8	83.6	82.2	83.2
Taking cocaine powder occasionally	93.0	93.4	91.2	91.0	92.7	89.7	89.3	88.7	90.0	90.3	89.8	90.2	88.9	90.0	89.4
Taking cocaine powder regularly	94.4	94.3	93.0	92.5	93.8	92.9	91.5	91.1	92.3	92.6	92.5	92.2	90.7	92.6	92.0
Trying heroin once or twice	96.0	94.9	94.4	93.2	92.8	92.1	92.3	93.7	93.5	93.0	93.1	94.1	94.1	94.2	94.3
Taking heroin occasionally	97.3	96.8	97.0	96.2	95.7	95.0	95.4	96.1	95.7	96.0	95.4	95.6	95.9	96.4	96.3
Taking heroin regularly	97.8	97.2	97.5	97.1	96.4	96.3	96.4	96.6	96.4	96.6	96.2	96.2	97.1	97.1	96.7
Trying heroin once or twice without using a needle	—	—	—	—	92.9	90.8	92.3	93.0	92.6	94.0	91.7	93.1	92.2	93.1	93.2
Taking heroin occasionally without using a needle	—	—	—	—	94.7	93.2	94.4	94.3	93.8	95.2	93.5	94.4	93.5	94.4	95.0
Trying amphetamines once or twice ^d	86.5	86.9	84.2	81.3	82.2	79.9	81.3	82.5	81.9	82.1	82.3	83.8	85.8	84.1	86.1
Taking amphetamines regularly ^d	96.0	95.6	96.0	94.1	94.3	93.5	94.3	94.0	93.7	94.1	93.4	93.5	94.0	93.9	94.8
Trying sedatives (barbiturates) once or twice ^e	90.6	90.3	89.7	87.5	87.3	84.9	86.4	86.0	86.6	85.9	85.9	86.6	87.8†	83.7	85.4
Taking sedatives (barbiturates) regularly ^e	97.1	96.5	97.0	96.1	95.2	94.8	95.3	94.6	94.7	95.2	94.5	94.7	94.4†	94.2	95.2
Trying one or two drinks of an alcoholic beverage (beer, wine, liquor)	29.8	33.0	30.1	28.4	27.3	26.5	26.1	24.5	24.6	25.2	26.6	26.3	27.2	26.0	26.4
Taking one or two drinks nearly every day	76.5	75.9	77.8	73.1	73.3	70.8	70.0	69.4	67.2	70.0	69.2	69.1	68.9	69.5	70.8
Taking four or five drinks nearly every day	90.6	90.8	90.6	89.8	88.8	89.4	88.6	86.7	86.9	88.4	86.4	87.5	86.3	87.8	89.4
Having five or more drinks once or twice each weekend	67.4	70.7	70.1	65.1	66.7	64.7	65.0	63.8	62.7	65.2	62.9	64.7	64.2	65.7	66.5
Smoking one or more packs of cigarettes per day	71.4	73.5	70.6	69.8	68.2	67.2	67.1	68.8	69.5	70.1	71.6	73.6	74.8	76.2	79.8
Taking steroids	90.5	92.1	92.1	91.9	91.0	91.7	91.4	90.8	88.9	88.8	86.4	86.8	86.0	87.9	88.8
<i>Approximate weighted N =</i>	<i>2,547</i>	<i>2,645</i>	<i>2,723</i>	<i>2,588</i>	<i>2,603</i>	<i>2,399</i>	<i>2,601</i>	<i>2,545</i>	<i>2,310</i>	<i>2,150</i>	<i>2,144</i>	<i>2,160</i>	<i>2,442</i>	<i>2,455</i>	<i>2,460</i>

Table continued on next page.

TABLE 14 (cont.)
Trends in Disapproval of Drug Use in Grade 12

Percentage who disapprove or strongly disapprove ^b

<i>Do you disapprove of people (who are 18 or older) doing each of the following?^a</i>	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2015–2016 change
Trying marijuana once or twice	55.6	58.6	55.5	54.8	51.6	51.3	48.8	49.1	48.0	45.5	43.1	-2.4
Smoking marijuana occasionally	69.3	70.2	67.3	65.6	62.0	60.9	59.1	58.9	56.7	52.9	50.5	-2.4
Smoking marijuana regularly	82.2	83.3	79.6	80.3	77.7	77.5	77.8	74.5	73.4	70.7	68.5	-2.2
Trying LSD once or twice	88.0	87.8	85.5	88.2	86.5	86.3	87.2	86.6	85.0	81.7	82.4	+0.7
Taking LSD regularly	95.9	94.9	93.5	95.3	94.3	94.9	95.2	95.3	94.7	92.5	92.4	-0.2
Trying ecstasy (MDMA) once or twice ^c	89.0	87.8	88.2	88.2	86.3	83.9	87.1	84.9†	83.1	84.5	84.0	-0.5
Trying cocaine once or twice	89.1	89.6	89.2	90.8	90.5	91.1	91.0	92.3	90.0	89.0	88.4	-0.6
Taking cocaine regularly	96.1	96.2	94.8	96.5	96.0	96.0	96.8	96.7	96.3	95.2	94.8	-0.3
Trying crack once or twice	88.8	88.8	89.6	90.9	89.8	91.4	92.8	91.4	89.3	90.2	90.1	0.0
Taking crack occasionally	92.9	92.4	93.3	94.0	92.6	93.9	95.0	93.6	91.9	92.5	92.0	-0.6
Taking crack regularly	93.8	93.6	93.5	94.3	93.1	94.4	95.4	94.1	92.4	92.8	92.6	-0.2
Trying cocaine powder once or twice	84.1	83.5	85.7	87.3	87.0	88.1	88.7	88.2	85.5	86.4	86.6	+0.3
Taking cocaine powder occasionally	90.4	90.6	91.7	92.3	91.0	92.2	93.0	91.7	90.4	91.3	90.6	-0.7
Taking cocaine powder regularly	93.2	92.6	92.8	93.9	92.6	93.8	95.0	94.1	91.7	92.4	92.0	-0.4
Trying heroin once or twice	93.8	94.8	93.3	94.7	93.9	94.3	95.8	95.6	94.7	94.2	94.1	-0.1
Taking heroin occasionally	96.2	96.8	95.3	96.9	96.2	96.3	97.0	96.9	96.6	95.3	95.5	+0.2
Taking heroin regularly	96.9	97.1	95.9	97.4	96.4	96.7	97.4	97.4	97.1	96.4	95.7	-0.6
Trying heroin once or twice without using a needle	93.7	93.6	94.2	94.7	93.2	92.6	95.2	93.7	92.5	92.6	93.8	+1.2
Taking heroin occasionally without using a needle	94.5	94.9	95.3	95.5	94.5	94.1	95.9	94.6	93.5	92.8	94.0	+1.1
Trying amphetamines once or twice ^d	86.3	87.3	87.2	88.2	88.1‡	84.1	83.9	84.9	83.1	81.4	82.1	+0.7
Taking amphetamines regularly ^d	95.3	95.4	94.2	95.6	94.9‡	92.9	93.9	93.2	93.0	92.2	92.2	0.0
Trying sedatives (barbiturates) once or twice ^e	85.3	86.5	86.1	87.7	87.6	87.3	88.2	88.9	88.5	87.4	86.5	-0.9
Taking sedatives (barbiturates) regularly ^e	95.1	94.6	94.3	95.8	94.7	95.1	96.1	95.8	95.0	94.7	94.8	+0.1
Trying one or two drinks of an alcoholic beverage (beer, wine, liquor)	29.0	31.0	29.8	30.6	30.7	28.7	25.4	27.3	29.2	28.9	28.8	-0.2
Taking one or two drinks nearly every day	72.8	73.3	74.5	70.5	71.5	72.8	70.8	71.9	71.7	71.1	71.8	+0.7
Taking four or five drinks nearly every day	90.6	90.5	89.8	89.7	88.8	90.8	90.1	90.6	91.9	89.7	91.1	+1.3
Having five or more drinks once or twice each weekend	68.5	68.8	68.9	67.6	68.8	70.0	70.1	71.6	72.6	71.9	74.2	+2.3
Smoking one or more packs of cigarettes per day	81.5	80.7	80.5	81.8	81.0	83.0	83.7	82.6	85.0	84.1	85.3	+1.3
Taking steroids	89.4	89.2	90.9	90.3	89.8	89.7	90.4	88.2	87.5	87.8	86.7	-1.1
<i>Approximate weighted N =</i>	2,377	2,450	2,314	2,233	2,449	2,384	2,301	2,147	2,078	2,193	2,000	

Table continued on next page.

TABLE 14 (cont.)
Trends in Disapproval of Drug Use in Grade 12

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. '—' indicates data not available. '‡' indicates that the question changed the following year. See relevant footnote for that drug. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aThe 1975 question asked about people who are 20 or older.

^bAnswer alternatives were: (1) Don't disapprove, (2) Disapprove, and (3) Strongly disapprove. Percentages are shown for categories (2) and (3) combined.

^cBeginning in 2014 "molly" was added to the question on disapproval of using MDMA once or twice. 2014 and 2015 data are not comparable to earlier years due to this change.

^dIn 2011 the list of examples was changed from upper, pep pill, bennie, speed to upper, speed, Adderall, Ritalin, etc. These changes likely explain the discontinuity in the 2011 results.

^eIn 2004 the question text was changed from barbiturates to sedatives/barbiturates and the list of examples was changed from downers, goofballs, reds, yellows, etc. to just downers. These changes likely explain the discontinuity in the 2004 results.

TABLE 15
Trends in Availability of Drugs as Perceived by 8th Graders

<i>How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?</i>	<i>Percentage saying fairly easy or very easy to get ^a</i>														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
Marijuana	—	42.3	43.8	49.9	52.4	54.8	54.2	50.6	48.4	47.0	48.1	46.6	44.8	41.0	41.1
LSD	—	21.5	21.8	21.8	23.5	23.6	22.7	19.3	18.3	17.0	17.6	15.2	14.0	12.3	11.5
PCP ^b	—	18.0	18.5	17.7	19.0	19.6	19.2	17.5	17.1	16.0	15.4	14.1	13.7	11.4	11.0
Ecstasy (MDMA) ^b	—	—	—	—	—	—	—	—	—	—	23.8	22.8	21.6	16.6	15.6
Crack	—	25.6	25.9	26.9	28.7	27.9	27.5	26.5	25.9	24.9	24.4	23.7	22.5	20.6	20.8
Cocaine powder	—	25.7	25.9	26.4	27.8	27.2	26.9	25.7	25.0	23.9	23.9	22.5	21.6	19.4	19.9
Heroin	—	19.7	19.8	19.4	21.1	20.6	19.8	18.0	17.5	16.5	16.9	16.0	15.6	14.1	13.2
Narcotics other than Heroin ^{b,c}	—	19.8	19.0	18.3	20.3	20.0	20.6	17.1	16.2	15.6	15.0	14.7	15.0	12.4	12.9
Amphetamines ^d	—	32.2	31.4	31.0	33.4	32.6	30.6	27.3	25.9	25.5	26.2	24.4	24.4	21.9	21.0
Crystal methamphetamine (ice) ^b	—	16.0	15.1	14.1	16.0	16.3	15.7	16.0	14.7	14.9	13.9	13.3	14.1	11.9	13.5
Sedatives (barbiturates)	—	27.4	26.1	25.3	26.5	25.6	24.4	21.1	20.8	19.7	20.7	19.4	19.3	18.0	17.6
Tranquilizers	—	22.9	21.4	20.4	21.3	20.4	19.6	18.1	17.3	16.2	17.8	16.9	17.3	15.8	14.8
Alcohol	—	76.2	73.9	74.5	74.9	75.3	74.9	73.1	72.3	70.6	70.6	67.9	67.0	64.9	64.2
Cigarettes	—	77.8	75.5	76.1	76.4	76.9	76.0	73.6	71.5	68.7	67.7	64.3	63.1	60.3	59.1
Steroids	—	24.0	22.7	23.1	23.8	24.1	23.6	22.3	22.6	22.3	23.1	22.0	21.7	19.7	18.1
<i>Approximate weighted N =</i>		8,355	16,775	16,119	15,496	16,318	16,482	16,208	15,397	15,180	14,804	13,972	15,583	15,944	15,730

Table continued on next page.

TABLE 15 (cont.)
Trends in Availability of Drugs as Perceived by 8th Graders

<i>How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?</i>	<i>Percentage saying fairly easy or very easy to get ^a</i>											<i>2015–2016 change</i>
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Marijuana	39.6	37.4	39.3	39.8	41.4	37.9	36.9	39.1	36.9	37.0	34.6	-2.4 s
LSD	10.8	10.5	10.9	10.0	10.0	9.3	7.5	7.4	6.9	6.6	6.9	+0.3
PCP ^b	10.5	9.5	10.1	9.1	8.0	7.9	6.7	5.8	5.5	5.1	4.8	-0.4
Ecstasy (MDMA) ^b	14.5	13.4	14.1	13.1	12.9	12.0	9.6	9.5	10.1	9.6	8.7	-0.9
Crack	20.9	19.7	20.2	18.6	17.9	15.7	14.4	13.7	12.0	11.3	11.1	-0.2
Cocaine powder	20.2	19.0	19.5	17.8	16.6	14.9	14.1	13.5	11.9	11.6	11.0	-0.6
Heroin	13.0	12.6	13.3	12.0	11.6	9.9	9.4	10.0	8.6	7.8	8.9	+1.0 s
Narcotics other than Heroin ^{b,c}	13.0	11.7	12.1	11.8‡	14.6	12.3	10.6	9.7	9.2	8.8	8.9	+0.1
Amphetamines ^d	20.7	19.9	21.3	20.2	19.6‡	15.0	13.4	12.8	12.1	11.8	12.1	+0.3
Crystal methamphetamine (ice) ^b	14.5	12.1	12.8	11.9	10.9	9.6	8.8	8.5	7.7	6.9	6.6	-0.3
Sedatives (barbiturates)	17.3	16.8	17.5	15.9	15.3	12.6	11.1	10.6	10.0	9.0	9.3	+0.3
Tranquilizers	14.4	14.4	15.4	14.1	13.7	12.0	10.5	10.4	9.8	9.8	11.4	+1.6 ss
Alcohol	63.0	62.0	64.1	61.8	61.1	59.0	57.5	56.1	54.4	53.6	52.7	-1.0
Cigarettes	58.0	55.6	57.4	55.3	55.5	51.9	50.7	49.9	47.2	47.0	45.6	-1.4
Steroids	17.1	17.0	16.8	15.2	14.2	13.3	12.5	12.9	11.8	11.6	12.6	+1.0
<i>Approximate weighted N = 15,502 15,043 14,482 13,989 14,485 15,233 14,235 13,605 13,208 13,494 15,628</i>												

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. ' ‡ ' indicates that the question changed the following year. See relevant footnote for that drug. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aAnswer alternatives were: (1) Probably impossible, (2) Very difficult, (3) Fairly difficult, (4) Fairly easy, (5) Very easy, and (6) Can't say, drug unfamiliar.

^bBeginning in 1993, data based on one of two of forms; *N* is one half of *N* indicated. Beginning in 2014 data based on one sixth of *N* indicated. For MDMA only: In 2014 the question text was changed in one form to include "Molly." In 2015 a second form was changed to including "Molly;" data based on one sixth of *N* indicated in 2014 and on one half of *N* indicated in 2015. An examination of the data did not show any effect from this wording change.

^cIn 2010 the list of examples for narcotics other than heroin was changed from methadone, opium to Vicodin, OxyContin, Percocet, etc. This change likely explains the discontinuity in the 2010 results.

^dIn 2011 the list of examples for amphetamines was changed from uppers, pep pills, bennies, speed to uppers, speed, Adderall, Ritalin, etc. These changes likely explain the discontinuity in the 2012 results.

TABLE 16
Trends in Availability of Drugs as Perceived by 10th Graders

<i>How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?</i>	<i>Percentage saying fairly easy or very easy to get ^a</i>														
	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
Marijuana	—	65.2	68.4	75.0	78.1	81.1	80.5	77.9	78.2	77.7	77.4	75.9	73.9	73.3	72.6
LSD	—	33.6	35.8	36.1	39.8	41.0	38.3	34.0	34.3	32.9	31.2	26.8	23.1	21.6	20.7
PCP ^b	—	23.7	23.4	23.8	24.7	26.8	24.8	23.9	24.5	25.0	21.6	20.8	19.4	18.0	18.1
Ecstasy (MDMA) ^b	—	—	—	—	—	—	—	—	—	—	41.4	41.0	36.3	31.2	30.2
Crack	—	33.7	33.0	34.2	34.6	36.4	36.0	36.3	36.5	34.0	30.6	31.3	29.6	30.6	31.0
Cocaine powder	—	35.0	34.1	34.5	35.3	36.9	37.1	36.8	36.7	34.5	31.0	31.8	29.6	31.2	31.5
Heroin	—	24.3	24.3	24.7	24.6	24.8	24.4	23.0	23.7	22.3	20.1	19.9	18.8	18.7	19.3
Narcotics other than Heroin ^{b,c}	—	26.9	24.9	26.9	27.8	29.4	29.0	26.1	26.6	27.2	25.8	25.4	23.5	23.1	23.6
Amphetamines ^d	—	43.4	46.4	46.6	47.7	47.2	44.6	41.0	41.3	40.9	40.6	39.6	36.1	35.7	35.6
Crystal methamphetamine (ice) ^b	—	18.8	16.4	17.8	20.7	22.6	22.9	22.1	21.8	22.8	19.9	20.5	19.0	19.5	21.6
Sedatives (barbiturates)	—	38.0	38.8	38.3	38.8	38.1	35.6	32.7	33.2	32.4	32.8	32.4	28.8	30.0	29.7
Tranquilizers	—	31.6	30.5	29.8	30.6	30.3	28.7	26.5	26.8	27.6	28.5	28.3	25.6	25.6	25.4
Alcohol	—	88.6	88.9	89.8	89.7	90.4	89.0	88.0	88.2	87.7	87.7	84.8	83.4	84.3	83.7
Cigarettes	—	89.1	89.4	90.3	90.7	91.3	89.6	88.1	88.3	86.8	86.3	83.3	80.7	81.4	81.5
Steroids	—	37.6	33.6	33.6	34.8	34.8	34.2	33.0	35.9	35.4	33.1	33.2	30.6	29.6	29.7
<i>Approximate weighted N =</i>		<i>7,014</i>	<i>14,652</i>	<i>15,192</i>	<i>16,209</i>	<i>14,887</i>	<i>14,856</i>	<i>14,423</i>	<i>13,112</i>	<i>13,690</i>	<i>13,518</i>	<i>13,694</i>	<i>15,255</i>	<i>15,806</i>	<i>15,636</i>

Table continued on next page.

TABLE 16 (cont.)
Trends in Availability of Drugs as Perceived by 10th Graders

<i>How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?</i>	Percentage saying fairly easy or very easy to get ^a											2015–2016 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Marijuana	70.7	69.0	67.4	69.3	69.4	68.4	68.8	69.7	66.9	65.6	64.0	-1.6
LSD	19.2	19.0	19.3	17.8	18.3	16.6	14.9	16.3	14.8	15.5	15.2	-0.4
PCP ^b	15.8	15.4	14.4	13.4	12.6	12.0	10.2	9.4	8.3	9.0	7.6	-1.4 s
Ecstasy (MDMA) ^b	27.4	27.7	26.7	25.6	25.7	24.8	21.0	20.7	20.4	19.3	16.3	-3.0 ss
Crack	29.9	29.0	27.2	23.9	22.5	19.7	18.4	17.1	15.1	14.4	13.9	-0.5
Cocaine powder	30.7	30.0	28.2	24.7	22.6	20.6	19.2	18.3	16.4	16.1	14.9	-1.2
Heroin	17.4	17.3	17.2	15.0	14.5	13.2	11.9	11.9	10.9	11.0	10.6	-0.4
Narcotics other than Heroin ^{b,c}	22.2	21.5	20.3	18.8‡	28.7	25.0	24.3	22.5	18.8	19.2	16.8	-2.4 s
Amphetamines ^d	34.7	33.3	32.0	31.8	32.6‡	28.5	27.3	26.5	25.2	27.3	22.9	-4.5 sss
Crystal methamphetamine (ice) ^b	20.8	18.8	15.8	14.0	13.3	11.8	10.7	10.0	9.8	8.9	8.2	-0.8
Sedatives (barbiturates)	29.9	28.2	26.9	25.5	24.9	22.0	20.2	18.3	16.7	16.6	14.2	-2.4 sss
Tranquilizers	25.1	24.9	24.1	22.3	21.6	20.8	19.7	18.3	17.5	19.4	20.5	+1.1
Alcohol	83.1	82.6	81.1	80.9	80.0	77.9	78.2	77.2	75.3	74.9	71.1	-3.8 sss
Cigarettes	79.5	78.2	76.5	76.1	75.6	73.6	72.9	71.4	69.0	66.6	62.9	-3.6 ss
Steroids	30.2	27.7	24.5	20.8	20.3	18.8	18.0	17.2	16.5	17.0	15.3	-1.7 s
<i>Approximate weighted N =</i> 15,804 15,511 14,634 15,451 14,827 14,509 14,628 12,601 12,574 15,186 14,126												

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. ' ‡ ' indicates that the question changed the following year. See relevant footnote for that drug. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aAnswer alternatives were: (1) Probably impossible, (2) Very difficult, (3) Fairly difficult, (4) Fairly easy, (5) Very easy, and (6) Can't say, drug unfamiliar.

^bBeginning in 1993, data based on one of two forms; *N* is one half of *N* indicated. Beginning in 2014 data based on one sixth of *N* indicated.

^cBeginning in 1993, data based on one of two of forms; *N* is one half of *N* indicated. Beginning in 2014 data based on one sixth of *N* indicated. For MDMA only:

In 2014 the question text was changed in one form to include "Molly." In 2015 a second form was changed to including "Molly;" data based on one sixth of *N* indicated in 2014 and on one half of *N* indicated in 2015. An examination of the data did not show any effect from this wording change.

^dIn 2011 the list of examples for amphetamines was changed from uppers, pep pills, bennies, speed to uppers, speed, Adderall, Ritalin, etc. These changes likely explain the discontinuity in the 2011 results.

TABLE 17
Trends in Availability of Drugs as Perceived by 12th Graders

	Percentage saying fairly easy or very easy to get ^a															
<i>How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?</i>	<u>1975</u>	<u>1976</u>	<u>1977</u>	<u>1978</u>	<u>1979</u>	<u>1980</u>	<u>1981</u>	<u>1982</u>	<u>1983</u>	<u>1984</u>	<u>1985</u>	<u>1986</u>	<u>1987</u>	<u>1988</u>	<u>1989</u>	<u>1990</u>
Marijuana	87.8	87.4	87.9	87.8	90.1	89.0	89.2	88.5	86.2	84.6	85.5	85.2	84.8	85.0	84.3	84.4
Amyl/butyl nitrites	—	—	—	—	—	—	—	—	—	—	—	—	23.9	25.9	26.8	24.4
LSD	46.2	37.4	34.5	32.2	34.2	35.3	35.0	34.2	30.9	30.6	30.5	28.5	31.4	33.3	38.3	40.7
Some other hallucinogen ^b	47.8	35.7	33.8	33.8	34.6	35.0	32.7	30.6	26.6	26.6	26.1	24.9	25.0	26.2	28.2	28.3
PCP	—	—	—	—	—	—	—	—	—	—	—	—	22.8	24.9	28.9	27.7
Ecstasy (MDMA) ^c	—	—	—	—	—	—	—	—	—	—	—	—	—	—	21.7	22.0
Cocaine	37.0	34.0	33.0	37.8	45.5	47.9	47.5	47.4	43.1	45.0	48.9	51.5	54.2	55.0	58.7	54.5
Crack	—	—	—	—	—	—	—	—	—	—	—	—	41.1	42.1	47.0	42.4
Cocaine powder	—	—	—	—	—	—	—	—	—	—	—	—	52.9	50.3	53.7	49.0
Heroin	24.2	18.4	17.9	16.4	18.9	21.2	19.2	20.8	19.3	19.9	21.0	22.0	23.7	28.0	31.4	31.9
Some other narcotic (including methadone) ^c	34.5	26.9	27.8	26.1	28.7	29.4	29.6	30.4	30.0	32.1	33.1	32.2	33.0	35.8	38.3	38.1
Amphetamines ^d	67.8	61.8	58.1	58.5	59.9	61.3	69.5	70.8	68.5	68.2	66.4	64.3	64.5	63.9	64.3	59.7
Crystal methamphetamine (ice)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	24.1
Sedatives (barbiturates) ^e	60.0	54.4	52.4	50.6	49.8	49.1	54.9	55.2	52.5	51.9	51.3	48.3	48.2	47.8	48.4	45.9
Tranquilizers	71.8	65.5	64.9	64.3	61.4	59.1	60.8	58.9	55.3	54.5	54.7	51.2	48.6	49.1	45.3	44.7
Alcohol	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Steroids	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
<i>Approximate weighted N =</i>	<i>2,627</i>	<i>2,865</i>	<i>3,065</i>	<i>3,598</i>	<i>3,172</i>	<i>3,240</i>	<i>3,578</i>	<i>3,602</i>	<i>3,385</i>	<i>3,269</i>	<i>3,274</i>	<i>3,077</i>	<i>3,271</i>	<i>3,231</i>	<i>2,806</i>	<i>2,549</i>

Table continued on next page.

TABLE 17 (cont.)
Trends in Availability of Drugs as Perceived by 12th Graders

	Percentage saying fairly easy or very easy to get ^a														
<i>How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?</i>	<u>1991</u>	<u>1992</u>	<u>1993</u>	<u>1994</u>	<u>1995</u>	<u>1996</u>	<u>1997</u>	<u>1998</u>	<u>1999</u>	<u>2000</u>	<u>2001</u>	<u>2002</u>	<u>2003</u>	<u>2004</u>	<u>2005</u>
Marijuana	83.3	82.7	83.0	85.5	88.5	88.7	89.6	90.4	88.9	88.5	88.5	87.2	87.1	85.8	85.6
Amyl/butyl nitrites	22.7	25.9	25.9	26.7	26.0	23.9	23.8	25.1	21.4	23.3	22.5	22.3	19.7	20.0	19.7
LSD	39.5	44.5	49.2	50.8	53.8	51.3	50.7	48.8	44.7	46.9	44.7	39.6	33.6	33.1	28.6
Some other hallucinogen ^b	28.0	29.9	33.5	33.8	35.8	33.9	33.9	35.1	29.5	34.5‡	48.5	47.7	47.2	49.4	45.0
PCP	27.6	31.7	31.7	31.4	31.0	30.5	30.0	30.7	26.7	28.8	27.2	25.8	21.9	24.2	23.2
Ecstasy (MDMA) ^c	22.1	24.2	28.1	31.2	34.2	36.9	38.8	38.2	40.1	51.4	61.5	59.1	57.5	47.9	40.3
Cocaine	51.0	52.7	48.5	46.6	47.7	48.1	48.5	51.3	47.6	47.8	46.2	44.6	43.3	47.8	44.7
Crack	39.9	43.5	43.6	40.5	41.9	40.7	40.6	43.8	41.1	42.6	40.2	38.5	35.3	39.2	39.3
Cocaine powder	46.0	48.0	45.4	43.7	43.8	44.4	43.3	45.7	43.7	44.6	40.7	40.2	37.4	41.7	41.6
Heroin	30.6	34.9	33.7	34.1	35.1	32.2	33.8	35.6	32.1	33.5	32.3	29.0	27.9	29.6	27.3
Some other narcotic (including methadone) ^c	34.6	37.1	37.5	38.0	39.8	40.0	38.9	42.8	40.8	43.9	40.5	44.0	39.3	40.2	39.2
Amphetamines ^d	57.3	58.8	61.5	62.0	62.8	59.4	59.8	60.8	58.1	57.1	57.1	57.4	55.0	55.4	51.2
Crystal methamphetamine (ice)	24.3	26.0	26.6	25.6	27.0	26.9	27.6	29.8	27.6	27.8	28.3	28.3	26.1	26.7	27.2
Sedatives (barbiturates) ^e	42.4	44.0	44.5	43.3	42.3	41.4	40.0	40.7	37.9	37.4	35.7	36.6	35.3‡	46.3	44.4
Tranquilizers	40.8	40.9	41.1	39.2	37.8	36.0	35.4	36.2	32.7	33.8	33.1	32.9	29.8	30.1	25.7
Alcohol	—	—	—	—	—	—	—	—	95.0	94.8	94.3	94.7	94.2	94.2	93.0
Steroids	46.7	46.8	44.8	42.9	45.5	40.3	41.7	44.5	44.6	44.8	44.4	45.5	40.7	42.6	39.7
<i>Approximate weighted N =</i>	<i>2,476</i>	<i>2,586</i>	<i>2,670</i>	<i>2,526</i>	<i>2,552</i>	<i>2,340</i>	<i>2,517</i>	<i>2,520</i>	<i>2,215</i>	<i>2,095</i>	<i>2,120</i>	<i>2,138</i>	<i>2,391</i>	<i>2,169</i>	<i>2,161</i>

Table continued on next page.

TABLE 17 (cont.)
Trends in Availability of Drugs as Perceived by 12th Graders

<i>How difficult do you think it would be for you to get each of the following types of drugs, if you wanted some?</i>	Percentage saying "fairly easy" or "very easy" to get ^a											2015–2016 change
	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	
Marijuana	84.9	83.9	83.9	81.1	82.1	82.2	81.6	81.4	81.3	79.5	81.0	+1.5
Amyl/butyl nitrites	18.4	18.1	16.9	15.7	—	—	—	—	—	—	—	
LSD	29.0	28.7	28.5	26.3	25.1	25.1	27.6	24.5	25.9	26.5	28.0	+1.6
Some other hallucinogen ^b	43.9	43.7	42.8	40.5	39.5	38.3	37.8	36.6	33.6	31.4	32.5	+1.0
PCP	23.1	21.0	20.6	19.2	18.5	17.2	14.2	15.3	11.1	13.8	12.6	-1.2
Ecstasy (MDMA) ^c	40.3	40.9	41.9	35.1	36.4	37.1	35.9	35.1	36.1	37.1	32.5	-4.7 s
Cocaine	46.5	47.1	42.4	39.4	35.5	30.5	29.8	30.5	29.2	29.1	28.6	-0.5
Crack	38.8	37.5	35.2	31.9	26.1	24.0	22.0	24.6	20.1	22.0	19.8	-2.2
Cocaine powder	42.5	41.2	38.9	33.9	29.0	26.4	25.1	28.4	22.3	25.8	22.9	-2.8
Heroin	27.4	29.7	25.4	27.4	24.1	20.8	19.9	22.1	20.2	20.4	20.0	-0.4
Some other narcotic (including methadone) ^d	39.6	37.3	34.9	36.1‡	54.2	50.7	50.4	46.5	42.2	39.0	39.3	+0.3
Amphetamines ^e	52.9	49.6	47.9	47.1	44.1‡	47.0	45.4	42.7	44.5	41.9	41.1	-0.7
Crystal methamphetamine (ice)	26.7	25.1	23.3	22.3	18.3	17.1	14.5	17.2	13.7	15.3	14.5	-0.8
Sedatives (barbiturates) ^f	43.8	41.7	38.8	37.9	36.8	32.4	28.7	27.9	26.3	25.0	25.7	+0.8
Tranquilizers	24.4	23.6	22.4	21.2	18.4	16.8	14.9	15.0	14.4	14.9	15.2	+0.3
Alcohol	92.5	92.2	92.2	92.1	90.4	88.9	90.6	89.7	87.6	86.6	85.4	-1.2
Steroids	41.1	40.1	35.2	30.3	27.3	26.1	25.0	28.5	22.0	23.7	21.3	-2.3
<i>Approximate weighted N =</i>												
	2,131	2,420	2,276	2,243	2,395	2,337	2,280	2,092	2,066	2,181	1,958	

Source. The Monitoring the Future study, the University of Michigan.

Notes. Level of significance of difference between the two most recent classes: s = .05, ss = .01, sss = .001. ' — ' indicates data not available. ' ‡ ' indicates that the question changed the following year. See relevant footnote for that drug. Any apparent inconsistency between the change estimate and the prevalence estimates for the two most recent years is due to rounding.

^aAnswer alternatives were: (1) Probably impossible, (2) Very difficult, (3) Fairly difficult, (4) Fairly easy, and (5) Very easy.

^bIn 2001 the question text was changed from other psychedelics to other hallucinogens and shrooms was added to the list of examples. These changes likely explain the discontinuity in the 2001 results.

^cBeginning in 2014 "molly" was added to the question on availability of Ecstasy (MDMA). An examination of the data did not show any effect from this wording change.

^dIn 2010 the list of examples for narcotics other than heroin was changed from methadone, opium to Vicodin, OxyContin, Percocet, etc. This change likely explains the discontinuity in the 2010 results.

^eIn 2011 the list of examples was changed from uppers, pep pills, bennies, speed to uppers, speed, Adderall, Ritalin, etc. These changes likely explain the discontinuity in the 2011 results.

^fIn 2004 the question text was changed from barbiturates to sedatives/barbiturates and the list of examples was changed from downers, goofballs, reds, yellows, etc. to just downers. These changes likely explain the discontinuity in the 2004 results.



Monitoring the Future website:
<http://www.monitoringthefuture.org>

