

# Prevalence and burden of bipolar disorders in European countries<sup>☆</sup>

Stefano Pini<sup>a,\*</sup>, Valéria de Queiroz<sup>a</sup>, Daniel Pagnin<sup>a</sup>, Lukas Pezawas<sup>b</sup>, Jules Angst<sup>c</sup>,  
Giovanni B. Cassano<sup>a</sup>, Hans-Ulrich Wittchen<sup>d,e</sup>

<sup>a</sup>Department of Psychiatry, Neurobiology, Pharmacology and Biotechnology, University of Pisa, Via Roma, 67-56100, Pisa, Italy

<sup>b</sup>Department of General Psychiatry, University Hospital of Psychiatry, Medical University of Vienna, Währinger Gürtel 18-20, 1090 Vienna, Austria

<sup>c</sup>Universität Zürich, Psychiatrische Uniklinik, Switzerland

<sup>d</sup>Technische Universität Dresden, Institute of Clinical Psychology and Psychotherapy, Germany

<sup>e</sup>Max-Planck-Institute of Psychiatry, Munich, Germany

## Abstract

A literature search, supplemented by an expert survey and selected reanalyses of existing data from epidemiological studies was performed to determine the prevalence and associated burden of bipolar I and II disorder in EU countries. Only studies using established diagnostic instruments based on DSM-III-R or DSM-IV, or ICD-10 criteria were considered. Fourteen studies from a total of 10 countries were identified. The majority of studies reported 12-month estimates of approximately 1% (range 0.5–1.1%), with little evidence of a gender difference. The cumulative lifetime incidence (two prospective-longitudinal studies) is slightly higher (1.5–2%); and when the wider range of bipolar spectrum disorders is considered estimates increased to approximately 6%. Few studies have reported separate estimates for bipolar I and II disorders. Age of first onset of bipolar disorder is most frequently reported in late adolescence and early adulthood. A high degree of concurrent and sequential comorbidity with other mental disorders and physical illnesses is common. Most studies suggest equally high or even higher levels of impairments and disabilities of bipolar disorders as compared to major depression and schizophrenia. Few data are available on treatment and health care utilization.

© 2005 Elsevier B.V. and ECNP. All rights reserved.

**Keywords:** Prevalence; Burden; Bipolar disorder; Mania; Hypomania; Depression

## 1. Introduction

Bipolar disorder (previously also labeled manic-depressive illness) is typically referred to as an episodic, yet lifelong and clinically severe affective (or mood) disorder. Bipolar disorder, associated with considerable treatment needs, is associated with tremendous social and occupational burden for both the individual and family in a substantial percentage of cases (Abood et al., 2002; Bebbington and Ramana, 1995;

Bijl and Ravelli, 2000a,b; Fichter et al., 1995; Simon, 2003; Wittchen et al., 2003; Woods, 2000). The term bipolar disorder, however, encompasses several phenotypes of mood disorders, i.e. mania, hypomania or cyclothymia that may present with a puzzling variety of other symptoms and disorders. According to the Fourth Edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV; APA, 1994), the diagnostic classificatory system used in most epidemiological studies, bipolar disorder is defined by a set of specific symptom criteria. Bipolar type I requires the presence or the history of at least one manic or mixed episode. Although, typically, patients with a manic episode also experience major depressive episodes, bipolar disorder can be diagnosed even if only one manic episode and no past major depressive episodes are present. Bipolar disorder type II differs from type I only by presence of hypomanic but no manic episodes. Hypomanic episodes differ from mania by a

<sup>☆</sup> This paper was prepared in the framework of European College of Neuropsychopharmacology (ECNP) Task Force project on Size and Burden of Mental Disorders in Europe (PI: Hans-Ulrich Wittchen). The paper also serves as input for the European Brain Council (EBC; www.ebc-eurobrain.net) Initiative Cost of Disorders of the Brain (Steering committee: Jes Olesen, Bengt Jönsson, Hans-Ulrich Wittchen).

\* Corresponding author. Tel.: +39 050 993 559; fax: +39 050 215 81.

E-mail address: s.pini@psico.med.unipi.it (S. Pini).

Table 1  
The prevalence of bipolar disorders in Europe

Country/region	Sample (n)/ Population	Instruments	Diagnoses	Age (years)	Prevalence (%) Overall (female/male)	Type of prevalence
<i>Belgium</i>						
Bruxelles region 1990 (Baruffol and Thilmany, 1993)	235 121 females 114 males General population	DISSI DSM-III	Mania	3 cohorts: 25,35,45	1.7 (0.8/2.6)	Lifetime
<i>Czech republic</i>						
Unpublished data		CIDI DSM-IV		18–65	0.5 (0.5/0.5)	12-month
<i>Germany</i>						
National 1998–1999 (Jacobi et al., 2004; Wittchen et al., 2000)	6159 3125 females 3034 males General population	M-CIDI DSM-IV	Any bipolar disorder	18–65	1.0 (1.2/0.8) 0.8 (1.1/0.6)	Lifetime 12-month
				18–34	1.1 (1.2/1.0)	12-month
				35–49	1.1 (1.5/0.7)	12-month
				50–65	0.3 (5.0/0.0)	12-month
Former West Germany 1981 (Wittchen et al., 1992)	1366 682 females 684 males General population	DIS DSM-III	Bipolar disorder	25–64	0.2 (0.5/0.0)	Lifetime
				25–44	0.5	Lifetime
				45–64	0	Lifetime
		ICD-9	Manic-depressive psychosis, circular type	25–64	0.1	Lifetime
			Manic- depressive psychosis, manic type	25–64	0.5	Lifetime
Munich region 1995 (Wittchen et al., 1998)	3021 1528 females 1493 males General population	M-CIDI DSM-IV	Bipolar I	14–24	1.4 (1.7/1.1) 1.3 (1.6/0.9)	Lifetime 12-month
			Bipolar II		0.4 (0.7/0.2) 0.4 (0.7/0.1)	Lifetime 12-month
			Single episode mania		0.1 (0.1/0.0) 0.1 (0.1/0.0)	Lifetime 12-month
			Hypomania		1.5 (1.7/1.4) 1.2 (1.5/0.9)	Lifetime 12-month
<i>Hungary</i>						
National 1995–1996 (Szádóczy et al., 1998)	2953 1645 females 1308 males General population	DIS DSM-III-R	Manic episode	18–64	1.5 (0.8/2.2) 0.9 (0.9/1.0) 0.5 (0.5/0.6)	Lifetime 12-month Point
				18–24	2.3 (1.4/3.3) 1.8 (1.1/2.9)	12-month Point
				25–34	1.4 (0.9/1.9) 1.2 (1.1/1.6)	12-month Point
				35–44	0.6 (0.3/0.0) 0.5 (0.6/0.4)	12-month Point
				45–54	0.5 (1.4/0.0) 0.5 (0.7/0.0)	12-month Point
				55–64	0.0 (0.0/0.0) 0.0 (0.0/0.0)	12-month Point
			Bipolar Disorder I	18–64	1.5 (1.6/1.3)	Lifetime
			Bipolar Disorder II or Atypical	18–64	2.0 (2.0/2.0)	Lifetime
			Hypomania	18–64	2.2 (2.1/2.2)	Lifetime
			Any Bipolar Disorder	18–64	5.1 (4.5/5.7)	Lifetime

Table 1 (continued)

Country/region	Sample (n)/ Population	Instruments	Diagnoses	Age (years)	Prevalence (%) Overall (female/male)	Type of prevalence
<i>Iceland</i>						
National 1987–1988 (Stefánsson et al., 1991)	862 421 females 441 males General population	DIS DSM-III	Bipolar disorder Atypical Bipolar Disorder	55–57 Cohort of people born in 1931	0.2 (0.2/0.2) 0.5 (0.5/0.5)	Lifetime Lifetime
<i>Northern Ireland</i>						
District of Derry region 1993–1994 (McConnell et al., 2002)	923 496 females 427 males General population	SCAN ICD-10	Bipolar Disorder	18–64	0.2	12-month
<i>Republic of Ireland</i>						
County of Monaghan region 1996 (Scully et al., 2004)	not provided General population	SCID DSM-III-R	Bipolar Disorder I Bipolar Disorder II	not provided	0.3 (0.3/0.3) 0.1 (0.1/0.1)	Lifetime Lifetime
<i>Italy</i>						
Florence area (Faravelli et al., 1990)	1000 537 females 463 males General population	Structured interview/ DSM-III DSM-III-R	Bipolar Disorder I Bipolar Disorder II Cyclothymic disorder	15 years and older	1.3 (1.9/0.7) 0.4 (0.2/0.2) 0.2 (0.4/0.0) 0.1 (0.2/0.0) 0.4 (1.6/0.2) 0.4 (0.56/0.22)	12-month Point 12-month Point 12-month Point
<i>Netherlands</i>						
National 1996 (Bijl et al., 1998a,b)	7076 3488 females 3588 males General population	CIDI DSM-III-R	Bipolar Disorder I/ Bipolar Disorder NOS	18–64	1.8 (2.1/1.5) 1.1 (1.1/1.1) 0.6 (0.8/0.4)	Lifetime 12-month Point
<i>Spain</i>						
Reus region 1987–1990 (Canals et al., 1997)	290 152 females 138 males General population	SCAN DSM-III-R ICD-10	Hypomania DSM-III-R Hypomania ICD-10	17–18	0.0 (0.0/0.0) 2.4 (3.9/0.7)	Point Point
<i>Spain</i>						
Cantabria region (Vasquez-Barquero et al., 1986, 1987)	1223 642 females 581 males General population	PSE-GHQ ICD-9	Manic depressive psychosis, manic type	17 years and older	0.08 (0.17/0.0)	Point
<i>Switzerland</i>						
Zurich 1978–1993 (Angst, 1995)	4547 2346 females 2201 males General population	SPIKE SCL-90 DSM-IV	Any bipolar disorder Mania	22–35	6.0 4.4	Lifetime Lifetime

shorter duration (at least 4 days instead of 1 week), and less severe impairment (not severe enough to cause marked impairment in social or occupational functioning, psychiatric hospitalization, or psychotic features). The DSM-IV also includes ‘cyclothymia’ as a bipolar spectrum disorder with hypomanic as well as depressive episodes that do not meet criteria for major depression (APA, 1994). More recently, some authors have suggested extending bipolar criteria in

various ways. For instance, expanding the diagnosis to include childhood conditions despite different symptom presentations (Biederman et al., 2003), relaxing duration criteria to include subthreshold manifestations (Angst, 1998; Angst et al., 2003a,b), and more generally by including a wider scope of bipolar spectrum disorders (Akiskal, 1996). Aside from these conceptual considerations, the primary aim of this paper is to review epidemiological surveys in the




Fig. 1. (a) First onset of hypomanic and manic episodes in a community sample ( $n=3021$ ) aged 14–24 years (EDSP). From: Wittchen et al., 2003. (b) First onset of hypomanic and manic episodes in a community sample ( $n=3021$ ) cumulative incidence after 5 years. From: Wittchen et al., 2003.

community that provide data on bipolar disorder I and II in Europe, focusing on identifying similarities and differences of the prevalence rates from various studies. Further, available information on age of onset, comorbidity with physical and mental disorders, and burden associated with bipolar disorders will be reviewed.

## 2. Methods

Studies referenced in Medline, EMBASE and Psycho Info and published after 1980 were included in this review. For details regarding the search process see Wittchen and Jacobi (2005). Only studies meeting the following criteria were included: (i) use of structured or standardized diagnostic interviews, (ii) use of diagnostic criteria for bipolar disorder according to ICD-10 (1992) or the APA (1994), (iii) data published in either English, German, French, Italian, Portuguese, or Spanish language, (iv) study conducted in an European country, (v) findings are based on epidemiological samples either in the community or in clinical populations. The review covered the 25 European Union (EU) membership countries (Austria, Belgium, Cyprus, Czech Republic, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Iceland, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherlands, Norway, Poland, Portugal, Slovakia,

Slovenia, Spain, Sweden and United Kingdom) plus Iceland, Norway and Switzerland.

## 3. Results

### 3.1. Prevalence and incidence

Table 1 displays the findings from a total of 14 studies in 10 EU countries meeting the inclusion criteria. All studies are general population samples. Age ranges covered were quite variable, ranging from birth cohort studies of subjects aged 55–57, over studies in 14–24 years olds, to studies covering a wider age range. The majority of studies included are regional surveys, with the exception of four studies. Five are based on small sample sizes of less than 1,000 subjects. The majority of studies used DSM-III, -III-R or -IV criteria, respectively—three studies were based on ICD-10 criteria.

Overall, the lifetime prevalence rate of mania, respectively bipolar I appears to be very similar across studies, with estimates ranging from 0.1–0.2% for two smaller Spanish studies to 1.8% in the Netherlands. The highest estimate was found in the Zurich study with an estimate of 4.4%. The 12-month rate for adults appears more consistent with estimates of 0.2% in Ireland to 1.1% in Netherlands. The point estimates indicated for some

studies are usually considerably lower. Only a few studies are powered or designed to indicate the prevalence of bipolar II disorder. In Hungary and Iceland, the overall lifetime prevalence of bipolar II disorder and atypical bipolar disorder appears to be higher than for bipolar disorder I. Conversely, in Italy and Ireland higher 12-month rates for bipolar disorder type I than bipolar II were reported. Overall estimates for bipolar II appear to be less consistent, probably due to the use of different diagnostic assessment instruments.

Taking together the findings of six studies that used largely comparable diagnostic assessment tools (namely the Composite international Diagnostic Interview; CIDI; Wittchen, 1994) and its predecessor the Diagnostic Interview Schedule, Robins et al., 1981) and that report 12-month rates (Czech Republic, Germany, Hungary, Netherlands, Norway) quite convergent 12-month rates for having any bipolar disorder were obtained for the age range 18–65. The median of the 12-month prevalence estimates across these studies, comprising altogether 21,848 subjects, was 0.9% (interquartile range: 0.5–0.9).

The female/male ratio of bipolar disorders was relatively similar in all studies, revealing slightly higher rates for females as compared to males. The only exception is a Belgian study with higher rates in men (Baruffol and Thilmany, 1993). Although none of the studies available was sufficiently powered to detect differences in age-specific 12-month prevalences, Table 1 reveals some indications that the highest rates occur in subjects aged 18–34. In contrast, the lowest estimates were found for the highest age groups examined. However, the large confidence intervals do not allow to conclude the existence of significantly different rates in the age range 18–65.

Findings on the age of first onset appear to be quite variable, possibly due to different sample composition. Using retrospective age of onset data and cumulative incidence data from a study in adolescents and young adults (ages up to 29), Fig. 1a reveals that the mean onset of first hypomanic and of manic episodes may be quite early and much earlier than those for depressive episodes. Among 14–24 years old in the Early Developmental Stages of Psychopathology Study (EDSP) the cumulative incidence was 2.3% for hypomania and 1.5% for DSM-IV manic episode. Over two subsequent waves of follow-up, covering a period of 5 years, the cumulative incidence up to age 29 was 4.7% (hypomania) and 2.6% for mania, respectively (Fig. 1b). The mean age of onset in this young birth cohort was 14.8 years for hypomania and 15.4 years for mania. In studies with samples of a wider age range (18–65) the mean age for the first manic or hypomanic episodes is 26.2 years, with 40% having their first episode between 18 and 24 years (NEMESIS, Ten Have et al., 2002). In the German National Survey (age 18–65) the mean age of onset was found to be 18 and 23.8 years, respectively, with 75% of the cases occurring up to 25 years (Jacobi et al., 2002, 2004;

Wittchen et al., 2003). In Hungary, the mean age of onset of bipolar disorders is 19.9 years, the first symptoms are reported slightly earlier by female (17.9 years) and the age interval with highest probability for development of the illness was between 15 and 19 years in both sexes (Szádóczy et al., 1998).

### 3.2. Natural course

Data on the natural course of bipolar disorders from epidemiological studies are available from three studies (Zurich study, EDSP, NEMESIS).

Using data from the EDSP (Table 2) it has been estimated that 20.7% of all subjects (aged 14–24 years at the outset of the study) with an initial hypomanic episode at baseline will experience another hypomanic or manic episode over the next 5 years; the risk for a depressive episode was estimated to be 26.7%. Relatively similar figures were found for those with an initial manic episode (19.3% and 33.1%, respectively). 59% developed neither a manic/hypomanic nor a depressive episode. Similar conditional proportions were reported for mania. Although these data are by and large concordant with observations from one other epidemiological study and clinical studies (Angst and Sellaro, 2000), caution in the generalizability of such results is needed, because such findings depend heavily on the studies' age range, the length of the follow-up and the diagnostic criteria and thresholds used.

The course of bipolar disorder is typically polyphasic with manic, depressive and mixed episodes. The short observation time in available epidemiological studies does not allow an in-depth characterization of the temporal patterns. There is evidence, however, that bipolar disorders experience a greater number of episodes as compared to unipolar depression. The length of episodes has been examined in several studies of both patient as well as community samples. Table 3 indicates that the four

Table 2  
Community sample—EDSP community cohort aged 14–24 years at baseline ( $n=2548$ ): Conditional probability of cases with hypomania or mania or major depression disorder (MDD; DSM-IV criteria) at baseline to develop further episodes over 4.2 years

Baseline condition	Type of further episodes over 5 years <sup>a,b</sup>			
	Hypomania	Mania	MDD	Neither
Hypomania (BIP I) ( $n=54$ )	10 (14.8%)	3 (5.9%)	15 (26.7%)	31 (59.0%)
Mania (BIP II) ( $n=34$ )	4 (13.4%)	3 (5.9%)	10 (33.1%)	19 (56.3%)
MDD ( $n=288$ )	8 (2.0%)	6 (1.8%)	59 (23.8%)	178 (73.4%)
No BIP/MDD ( $n=2162$ )	58 (2.3%)	26 (1.0%)	215 (9.4%)	1937 (88.1%)

From Wittchen et al., 2003.

<sup>a</sup> Numbers indicate conditional probabilities (%) by line.

<sup>b</sup> few cases developed psychotic syndromes ( $n=4$  cases).

Table 3  
Natural length of episodes (in months) in patient and community samples (adapted from Angst and Sellaro, 2000)

Study	Mean	Median	Q1	Q3
Mendel (1881)		5–6	3–4	6–7
Kraepelin (1913)	3–8			
Panse (1924)	7			
Wertham (1929)		4–6	2–4	8–10
Rennie (1942)	3.5–5.8			
Kinkelin (1954)	3.5–4.8			
Keller et al. (1986)	3–6			
Angst and Preisig (1995)	4.3	3	2	5
Eaton et al. (1997)	2–3			
Lewinsohn et al. (1995)	2			
Wittchen et al. (2000; GHS <sup>a</sup> )	1.7			
Wittchen et al. (2003; EDSP <sup>b</sup> )	0.9			

<sup>a</sup> German National Health Interview and Examination Survey.

<sup>b</sup> Early Developmental Stages of Psychopathology.

population based epidemiological studies (bottom of list) reveal shorter mean episode duration (0.9–3 months) as compared to the findings in patient samples (3+ months) and those of less recent studies (7–8 months). It is not clear to what degree these differences are influenced by age and severity.

### 3.3. Comorbidity with mental disorders

The vast majority of the studies reviewed suggest that bipolar disorder is frequently comorbid with a wide range of other disorders. Significant associations (odds ratios controlled for age and gender as a measure for the strength of relationship) were found for all axis I disorders examined. In the German and Dutch studies almost all patients who were diagnosed with bipolar I disorder had a lifetime history of at least one other axis I disorder. More than two third of patients had a history of one or more anxiety disorders and over 70% had a history of a substance abuse disorder. Among bipolar disorder subjects the rates of alcohol use ranges from 21.4 % in adults to 54.5% in adolescents and young adults (Angst, 1998; Wittchen et al., 2003). With regard to anxiety disorders, persons with bipolar disorder have an increased risk of generalized anxiety disorder, panic disorder and specific phobias (Szádóczy et al., 1998; Wittchen et al., 1998). Specifically, bipolar II disorder is associated with a more than 10-fold increased risk of panic disorder and repeated panic attacks than compared to without bipolar (Angst, 1998). In the German EDSP cohort study of adolescents and young adults (up to 29 years), a high degree of comorbidity is evident even in early stages of this disorder. Particularly high associations were evident for the link between bipolar I disorder with substance use, panic disorder, agoraphobia, post-traumatic stress disorder (PTSD) and obsessive-compulsive disorder (OCD), both within as well as in between episodes (Wittchen et al., 2003).

The Zurich Cohort community study is among the few community studies also reporting the prevalence of antisocial behavior and classic measurements of personality, as aggression and neuroticism, and verified that all measures are elevated in bipolar patients when compared with a control group (Angst, 1995). Despite the interesting implications of links between personality, personality disorder and bipolar syndromes (Maier et al., 1995), no epidemiological study to date has examined the association between personality disorders and bipolar disorders in unselected community samples, with the exception of elevated odds for antisocial personality disorder (Robins and Regier, 1991). Evidence from clinical studies usually reveals increased rates for personality disorders; however, there are rarely substantial differences between unipolar and bipolar depressive patients (Rossi et al., 2001). Most studies indicate that both conditions have elevated rates of avoidant and dependent personality disorder (Akiskal, 1996; Battaglia et al., 1996; Rossi et al., 2001). A small number suggest elevated risks for obsessive and/or borderline features are an unnoticed sign of bipolarity (Akiskal, 1996; Savino et al., 1993).

### 3.4. Somatic disorders and mortality

Bipolar disorder is associated with reduced general and mental health perception, emotional and social functioning, and vitality when compared with other types of mental disorder and controls (Angst, 1995; Ten Have et al., 2002). Wunderlich et al. (1998) highlighted that bipolar disorders have an elevated risk of suicide attempts, though it is not clear whether this increase in risk is due to the high degree of comorbidity, or to the manic/hypomanic or depressive component. Further, there have been UK reports that suicide is the cause of death of approximately 20% of persons with bipolar disorder (Ösby et al., 2001) and 60% mention suicidal ideation (Ten Have et al., 2002). The increased risk for suicide attempts is high in the first years of the illness; and clinical studies highlight the highest suicide risk during the first year after the first hospital admission regardless of age and gender (Wunderlich et al., 1998). Further, a decrease in suicidality with increasing duration of illness up to 5 years after first admission has been observed (Hoyer et al., 2000, 2004).

Bipolar disorders are also associated with a poor physical health status and with increased deaths due to natural causes. The most frequent diseases that cause death are cardiovascular, followed by cancer, and respiratory and cerebrovascular diseases. Bipolar patients with a history of hospitalization are more likely to die because of unnatural causes (i.e. suicide), with a higher mortality ratio in patients with their first admission at younger ages and during the first years after the diagnosis (Hoyer et al., 2000; Ösby et al., 2001).

### 3.5. Health services utilization

Little information is available from the review of studies concerning health care utilization and treatment. Individuals with bipolar disorders are more likely to use all forms of care compared to people with somatoform, substance use and some anxiety disorders and less likely compared to panic disorder, GAD, OCD and psychotic disorders (Bijl et al., 1998a,b; Jacobi et al., 2004). Faravelli et al. (1990) observed that patients with bipolar disorder types I and II had a significantly greater probability of seeking any form of medical help, being prescribed medication by their general practitioner, being referred to a psychiatrist and being hospitalized than patients with cyclothymia, dysthymia or minor depression. Ten Have et al. (2002) showed that 70% of the sample with bipolar disorder had sought help for emotional problems from a primary care provider (general practitioner or a community social worker), 56% had called a mental health care provider and 25.5% had never sought any help. Of the 56% who sought mental health care, 20% had never spoken to a professional about their hypomanic or manic episode. Thus, these patients remained under-recognized and did not receive adequate treatment. Epidemiologic parameters such as hospitalization risk and duration, outpatient treatment duration with drugs, as well as other forms of intervention require additional study.

### 3.6. Costs

There are only three studies in Europe that estimated cost-of-illness for bipolar disorder. De Zelicourt et al. (2003) estimated the annual costs of inpatient care associated with manic episodes in France to be around 1.3 billion, with 98% of the total cost referring to hospitalisation. Olié and Lévy (2002) estimated the 3-month direct medical costs for manic episode necessitating hospitalisation and in a 3-month period following hospitalisation in 22297. Cost of hospitalisation emerged as the main cost driver and the cost of institutionalisation (hospitalisation plus nursing home care) accounted for more than 99% of all costs. A limitation of this study is that consists of a 3-month cross-sectional evaluation of this patient population; therefore, extrapolation to long-term management cannot be easily undertaken. Moreover, this study specifically investigated the cost of single manic episodes, not the wider costs of treating bipolar illness. As pointed out by the authors, the study was based on hospital records and, because of the variability of the diligence with which out-patient treatment is recorded by hospitals, might have underestimated the direct costs associated with out-patient treatment. On the other hand, as manic episodes are very disruptive of daily life, work and career development as well as family life and relationships, it is likely that indirect costs would be relatively high in this population of manic patients.

In the UK, the annual cost attributed to bipolar disorder was estimated to be £2 billion, at 1999/2000 currency value or approximately £6900 per person with bipolar disorder (Das Gupta and Guest, 2002). The unemployment rate among people with bipolar disorder was estimated to be 46%. This figure has been compared with the 3% of 1999/2000 unemployed rate among general population in UK. As a result, an excess of 76,500 people annually are unemployed for cause attributable to bipolar disorder. Therefore, the annual indirect cost related to excess unemployment among people with bipolar disorder was estimated to be £1510 million at 1999/2000 value (Das Gupta and Guest, 2002).

## 4. Discussion

### 4.1. Prevalence

The epidemiology of bipolar disorders in Europe has been described in several studies with a remarkable degree of consistency across diverse study designs and countries. The evidence from community studies is highlighted by the clinical description of bipolar disorder as an episodic disorder that usually emerges in early adulthood, with a mean age of onset estimated to be between age 20 and 30. There is fairly convergent evidence that bipolar I and II disorders, according to DSM-IV criteria, have an estimated 12-month prevalence of approx. 1% with no evidence for major differences by age group and gender. This finding is consistent with other studies and reviews. Epidemiological studies carried out in US in the 1990s suggest a lifetime prevalence rate for bipolar I disorder of approximately 1% (Kessler et al., 1994; Jonas et al., 2003; Weissman et al., 1996). Angst (1998) reviewed results from another ten studies between 1985 and 1994 and reported lifetime prevalence rates of 0.7%. He also reviewed nine reports of bipolar II disorder suggesting lifetime prevalence rates of 0.2–3.0%.

Some exceptional prevalence estimates come from are prospective-longitudinal studies with several waves of investigations over many years. Data from Switzerland reported a lifetime prevalence of mania of 4.4%. Similarly, Wittchen et al. (2003) reported a cumulative lifetime incidence risk for community subjects up to age 30 of 2.6%. For bipolar II such longitudinal studies report even higher rates. The findings of five studies in this review report estimates ranging from 2.6% to 6.0%. The reason of such high rates could largely be explained by design and assessment differences. The two highest rates (5.1% and 6.0%), obtained in the studies of Szádóczy et al. (1998) in Hungary and Angst (1998) in Switzerland are probably explained by the adoption of less stringent criteria for diagnosis of bipolar disorder with the inclusion of softer forms along with bipolar type I.

#### 4.2. Disability and treatment

Regarding disability and treatment the existing EU data base is very limited, largely due to the fact, that bipolar I and II disorder are relatively rare disorders; no epidemiological study was powered to describe these disorders in greater detail. Nevertheless, there are a few noteworthy observations. When compared to other mood disorders, anxiety disorders or substance use disorders, persons with bipolar disorder tend to reveal a lower level of functioning, a greater severity of disability and a longer duration of illness. Persons with bipolar I disorder suffer greater losses in productivity, with more bed rest and absenteeism days (Abood et al., 2002; Simon, 2003; Ten Have et al., 2002; Wittchen et al., 1992, 2003). In 1990, the World Health Organization identified bipolar disorder as the sixth leading cause of disability-adjusted life years in the world among people aged 15–44 years with high costs to society (Kleinman et al., 2003; Murray and Lopez, 1996; Woods, 2000). However, studies on bipolar costs in Europe are still scarce, are based on approximate value for direct costs and cannot be generalized to all European countries.

With regard to early recognition and treatment there are even fewer data. Currently, the proportion of bipolar patients in treatment, and how patients are treated and managed through the various stages of their disorder, is not known (Goodwin, 1999; Goodwin and Ghaemi, 1999; Kessing et al., 1998). Furthermore, in clinical samples evidence exists that bipolar disorder may be under-diagnosed (Ghaemi et al., 1999). Treatment choices are particularly important in patients with bipolar disorder where compliance with conventional maintenance treatment is low. For example, some atypical antipsychotics have been shown to have good acute anti-manic and preventive mood-stabilizing properties together with a low incidence of extra-pyramidal effects as compared to typical neuroleptics. These factors may be important for improving compliance. In contrast, significant delays in the diagnosis and treatment of bipolar patients appear to be the rule rather than the exception. The American National Depressive and Manic-Depressive Association (DMDA) survey of bipolar members reported that 50% of bipolar patients did not seek help for 5 years or more after onset of their initial symptoms. The survey also reported that 48% of patients did not receive a bipolar disorder diagnosis until seeing the third professional consulted and that in 34% the interval between seeking help and receiving a bipolar diagnosis was more than 10 years. This delay in receiving care was greatest when symptoms began during childhood and adolescence. Additionally, of those patients who do seek treatment during the initial episode, about one third are likely to have the condition misdiagnosed (Lish et al., 1994). For these reasons, it is important to make a timely, accurate diagnosis and to provide optimal treatment with available pharmacologic options that provide sufficient long-term efficacy while minimizing the risk of side effects (Müller-Oerlinghausen et al., 2002).

#### 4.3. Future research needs

1. There is clearly a marked need for epidemiological studies that provide a fuller account of the degree of met and unmet needs for intervention, treatment and long-term management of bipolar disorders. Because of the low prevalence of bipolar disorder, such studies might either use two-stage designs with screening in the first stage or cohort designs in high risk populations (Mortensen et al., 2003). These studies should not only estimate the degree of under-diagnosing but should also provide data on the type of drug and non-drug treatment received by these patients in various sectors of the health care system.
2. Beyond the assessment of bipolar I and II disorders as currently defined by DSM-IV and ICD-10, these studies should also consider wider definitions. The current diagnostic criteria for bipolar disorders are under discussion and the use of wider criteria for so-called bipolar spectrum disorders (Akiskal and Pinto, 1999) has been stimulated by several comprehensive epidemiological inquiries. Angst et al. (2003a,b) has provided evidence for various alternative diagnostic models that might prove to be more useful and adequate as the current diagnoses. The core role of future epidemiological work in the domain is exploration of more adequate threshold definitions and the provision of data demonstrating their clinical validity and utility.
3. There is a need for studies that provide a better understanding of the complex comorbidity patterns typically associated with bipolar disorders. Most valuable in this respect may be cohort studies in children and adolescents. This is particularly true in light of recent speculations that bipolar disorders might start even before age of 10 (Biederman et al., 2003). It is speculated that a substantial proportion of children and adolescents remain undiagnosed or misdiagnosed, not receiving adequate treatment. In this respect, the relationship to attentional deficit and hyperactivity disorders (ADHD) in childhood and adulthood might be of special interest. Such epidemiological studies may serve a useful double purpose: contributing data that help resolving basic, as well as a clinical, research issues such as threshold and comorbidity, and secondly, a public health issue, namely providing data to improve the situation of patients.

#### References

- Abood, Z., Sharkey, A., Weeb, M., Kelly, A., Gill, M., 2002. Are patients with bipolar affective disorder socially disadvantaged? A comparison with a control group. *Bipolar Disord.* 4, 243–248.
- Akiskal, H.S., 1996. The prevalence clinical spectrum of bipolar disorders: beyond DSM-IV. *J. Clin. Psychopharmacol.* 16 (Suppl. 1), 4–14.
- Akiskal, H.S., Pinto, O., 1999. The evolving bipolar spectrum. Prototypes I, II, III, and IV. *Psychiatr. Clin. North Am.* 22, 517–534.

- American Psychiatric Association, 1994. Diagnostic and Statistical Manual of Mental Disorders, 4th ed. American psychiatric Association, Washington, DC.
- Angst, J., 1995. Epidémiologie du spectre bipolaire. *Encéphale* (Suppl. VI), 37–42.
- Angst, J., 1998. The emerging epidemiology of hypomania and bipolar II disorder. *J. Affect. Disord.* 50, 143–151.
- Angst, J., Preisig, M., 1995. Course of a clinical cohort of unipolar, bipolar and schizoaffective patients: results of a prospective study from 1959 to 1985. *Schweiz. Arch. Neurol. Psychiatr.* 146, 5–16.
- Angst, J., Sellaro, R., 2000. Historical perspectives and natural history of bipolar disorder. *Biol. Psychiatry* 48, 445–457.
- Angst, J., Gamma, A., Benazzi, F., Ajdacic, V., Eich, D., Rössler, H.W., 2003a. Diagnostic issues in bipolar disorder. *Eur. Neuropsychopharmacol.* 13, S43–S50.
- Angst, J., Gamma, A., Benazzi, F., Ajdacic, V., Eich, D., Rössler, W., 2003b. Toward a re-definition of subthreshold bipolarity: epidemiology and proposed criteria for bipolar-II, minor bipolar disorders and hypomania. *J. Affect. Disord.* 73, 133–146.
- Baruffol, E., Thilmany, M.C., 1993. Anxiety, depression, somatization and alcohol abuse. Prevalence rates in a general Belgian community sample. *Acta Psychiatr. Belg.* 93, 136–153.
- Battaglia, M., Przybeck, T.R., Bellodi, L., Cloninger, C.R., 1996. Temperament dimensions explain the comorbidity of psychiatric disorders. *Compr. Psychiatry* 37 (Suppl. 4), 292–298.
- Bebbington, P., Ramana, R., 1995. The epidemiology of bipolar affective disorder. *Soc. Psychiatr. Epidemiol.* 30, 279–292.
- Biederman, J., Mick, E., Faraone, S.V., Spencer, T., Wilens, T.E., Wozniak, J., 2003. Current concepts in the validity, diagnosis and treatment of paediatric bipolar disorder. *Int. J. Neuropsychopharmacol.* 6 (Suppl. 3), 293–300.
- Bijl, R.V., Ravelli, A., 2000a. Current and residual functional disability associated with psychopathology: findings from the Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Psychol. Med.* 30, 657–668.
- Bijl, R.V., Ravelli, A., 2000b. Psychiatric morbidity, service use, and need for care in the general population: results of the Netherlands Mental Health Survey and incidence study. *Am. J. Public Health* 90, 602–607.
- Bijl, R.V., van Zessen, G., Ravelli, A., de Rijk, C., Langendoen, Y., 1998a. The Netherlands Mental Health Survey and Incidence Study (NEMESIS): objectives and design. *Soc. Psychiatry Psychiatr. Epidemiol.* 33, 581–586.
- Bijl, R.V., Ravelli, A., van Zessen, G., 1998b. Prevalence of psychiatric disorder in the general population: results of the Netherlands Mental Health Survey and Incidence Study (NEMESIS). *Soc. Psychiatry Psychiatr. Epidemiol.* 33, 587–595.
- Canals, J., Domènech, E., Carbajo, C., Bladé, J., 1997. Prevalence of DSM-III-R and ICD-10 psychiatric disorders in a Spanish population of 18-year-olds. *Acta Psychiatr. Scand.* 96, 287–294.
- Das Gupta, R., Guest, J.F., 2002. Annual cost of bipolar disorder to UK society. *Br. J. Psychiatry* 180, 227–233.
- De Zelicourt, M., Dardennes, R., Verdoux, H., Gandhi, G., Khoshnood, B., Chomette, E., Papatheodorou, M.L., Edgell, E.T., Even, C., Fagnani, F., 2003. Frequency of hospitalisations and inpatient care costs of manic episodes: inpatients with bipolar I disorder in France. *Pharmacoeconomics* 21, 1081–1090.
- Eaton, W.W., Anthony, J.C., Gallo, J., Cai, G.J., Tien, A., Romanoski, A., Lyketsos, C., Chen, L.S., 1997. Natural history of diagnostic interview schedule DSM-IV major depression—The Baltimore Epidemiologic Catchment Area Follow-up. *Arch. Gen. Psychiatry* 54, 993–999.
- Faravelli, C., Degl'Innocenti, B.G., Aiuzzi, L., Incerpi, G., Pallanti, S., 1990. Epidemiology of mood disorders: a community survey in Florence. *J. Affect. Disord.* 20, 135–141.
- Fichter, M.M., Rehm, J., Elton, M., Dilling, H., Achatz, F., 1995. Mortality risk and mental disorders; longitudinal results from the Upper Bavarian Study. *Psychol. Med.* 25, 297–307.
- Ghaemi, N.S., Sachs, G.S., Chiou, A.M., 1999. Is bipolar disorder still under diagnosed? Are the antidepressants over utilized? *J. Affect. Disord.* 52, 135–144.
- Goodwin, G.M., 1999. Prophylaxis of bipolar disorder: how and who should be treated in the long term? *Eur. Neuropsychopharmacology* 9, 125–129.
- Goodwin, F.K., Ghaemi, S.N., 1999. The impact of the discovery of lithium on psychiatric thought and practice in the EUA and Europe. *Aust. N. Z. J. Psychiatry* 33, 54–64.
- Hoyer, E.H., Mortensen, P.B., Olesen, A.V., 2000. Mortality and causes of death in a total national sample of patients with affective disorders admitted for the first time between 1973 and 1993. *Br. J. Psychiatry* 176, 76–82.
- Hoyer, E.H., Olesen, A.V., Mortensen, P.B., 2004. Suicide risk in patients hospitalised because of an affective disorder: a follow-up study, 1973–1993. *J. Affect. Disord.* 78 (Suppl. 3), 209–217.
- Jacobi, F., Wittchen, H.-U., Hölling, G., Sommer, S., Lieb, R., Höfler, M., Pfister, H., 2002. Estimating the prevalence of mental and somatic disorders in the community: aims and methods of German National Health Interview and Examination Survey. *Int. J. Methods Psychiatr. Res.* 11, 1–18.
- Jacobi, F., Wittchen, H.-U., Hölling, C., Höfler, M., Pfister, N.M., Lieb, R., 2004. Prevalence, comorbidity and correlates of mental disorders in the general population: results from the German Health Interview and Examination Survey (GHS). *Psychol. Med.* 34, 1–15.
- Jonas, B.S., Brody, D., Roper, M., Narrow, W.E., 2003. Prevalence of mood disorders in a national sample of young American adults. *Soc. Psychiatry Psychiatr. Epidemiol.* 3, 618–624.
- Keller, M.B., Lavori, P.W., Coryell, W., Andreasen, N.C., Endicott, J., Clayton, P.J., Klerman, G.L., Hirschfeld, R.M., 1986. Differential outcome of pure manic, mixed/cycling, and pure depressive episodes in patients with bipolar illness. *JAMA* 255, 3138–3142.
- Kessing, L.V., Andersen, P.K., Mortensen, P.B., Bolwig, T.G., 1998. Recurrence in affective disorder: I. Case register study. *Br. J. Psychiatry* 172, 23–28.
- Kessler, R.C., McGonagle, K.A., Zhao, S., Nelson, C.B., Hughes, M., Eshleman, S.M.A., Wittchen, H.-U., Kendler, S., 1994. Lifetime and 12 months prevalence of DSM-III-R psychiatric disorders in the United States: results from National Comorbidity Survey. *Arch. Gen. Psychiatry* 51, 8–19.
- Kinkelin, M., 1954. Verlauf und Prognose des Manisch-Depressiven Irreseins. *Schweiz. Arch. Neurol. Neurochir. Psychiatr.* 73, 100–146.
- Kleinman, L., Lowin, A., Flood, E., Gandhi, G., Edgell, E., Revicki, D., 2003. Costs of bipolar disorder. *Pharmacoeconomics* 21, 601–622.
- Kraepelin, E., 1913. *Psychiatrie*, (8th edn.). Ein Lehrbuch für Studierende und Ärzte, vol. III. Barth, Leipzig.
- Lewinsohn, P.M., Klein, D.N., Seeley, J.R., 1995. Bipolar disorders in a community sample of older adolescents: prevalence, phenomenology, comorbidity, and course. *J. Am. Acad. Child Adolesc. Psych.* 34, 454–463.
- Lish, J.D., Dime-Meenan, S., Whybrow, P.C., Price, P.A., Hirschfeld, R.M.A., 1994. The National Depressive and Manic-depressive Association (DMDA) survey of bipolar members. *J. Affect. Disord.* 31, 281–294.
- Maier, W., Minges, J., Lichtermann, D., Heun, R., 1995. Personality disorders and personality variations in relatives of patients with bipolar affective disorders. *J. Affect. Disord.* 35, 173–181.
- McConnell, P., Bebbington, P., McClelland, R., Gillespie, K., Houghton, S., 2002. Prevalence of psychiatric disorder and the need for psychiatric care in Northern Ireland. *Br. J. Psychiatry* 181, 214–219.
- Mendel, E., 1881. *Die Manie. Eine Monographie.* Urban Schwarzenberg, Vienna.
- Mortensen, P.B., Pedersen, C.B., Melbye, M., Mors, O., Ewald, H., 2003. Individual and familial risk factors for bipolar affective disorders in Denmark. *Arch. Gen. Psychiatry* 60, 1209–1215.
- Müller-Oerlinghausen, B., Berghöfer, A., Bauer, M., 2002. Bipolar disorder. *Lancet* 359, 241–247.
- Murray, C.J.L., Lopez, A.D., 1996. *The Global Burden of Disease.* World Health Organization. Harvard University Press, Cambridge.

- Olié, J.P., Lévy, E., 2002. Manic episodes: the direct cost of a three-month period following hospitalization. *Eur. Psychiatr.* 17, 278–286.
- Ösby, U., Brandt, L., Correia, N., Ekblom, A., Sparen, P., 2001. Excess mortality in bipolar and unipolar disorder in Sweden. *Arch. Gen. Psychiatry* 58, 844–850.
- Pansee, F., 1924. Untersuchungen über Verlauf und Prognose beim Manisch-Depressiven Irresein. *Monatsschr. Psychiatr. Neurol.* 56, 15–82.
- Rennie, T.A.C., 1942. Prognosis in manic-depressive psychoses. *Am. J. Psychiatry* 98, 801–814.
- Robins, L.N., Regier, D.A. (Eds.), 1991. *Psychiatric disorders in America: the Epidemiologic Catchment Area Study*. The Free Press, New York.
- Robins, L.N., Helzer, J.E., Croughan, J., Ratcliff, K.S., 1981. National Institute of Mental Health Diagnostic Interview Schedule: its history, characteristics and validity. *Arch. Gen. Psychiatry* 38, 381–389.
- Rossi, A., Marinangeli, M.G., Butti, G., Scinto, A., DiCicco, L., Kalyvoka, A., Petrucci, C., 2001. Personality disorders in bipolar and depressive disorders. *J. Affect. Disord.* 65 (Suppl. 1), 3–8.
- Savino, M., Perugi, G., Simonini, E., Soriani, A., Cassano, G.B., Akiskal, H.S., 1993. Affective comorbidity in panic disorder: is there a bipolar connection? *J. Affect. Disord.* 28 (Suppl. 3), 155–163.
- Scully, P.J., Owens, J.M., Kinsella, A., Waddington, J.L., 2004. Schizophrenia, schizoaffective and bipolar disorder within an epidemiologically complete, homogeneous population in rural Ireland: small area variation in rate. *Schizophr. Res.* 67, 143–155.
- Simon, G.E., 2003. Social and economic burden of mood disorders. *Biol. Psychiatry* 54, 208–215.
- Stefánsson, J.G., Lindal, E., Björnsson, J.K., Guðmundsdóttir, Á., 1991. Lifetime prevalence of specific mental disorders among people born in Iceland in 1931. *Acta Psychiatr. Scand.* 84, 142–149.
- Szádóczky, E., Papp, Zs., Vitrai, J., Rihmer, Z., Furedi, J., 1998. The prevalence of major depressive and bipolar disorder in Hungary. Results from a national epidemiologic survey. *J. Affect. Disord.* 50, 153–162.
- Ten Have, M., Vollebergh, W., Bijl, R., Nolen, W.A., 2002. Bipolar disorder in a general population in the Netherlands (prevalence, consequences and care utilization): results from the Netherlands Mental Health Survey and Incidence Study (NEMESIS). *J. Affect. Disord.* 68, 203–213.
- Vasquez-Barquero, J.L., Diez-Manrique, J.F., Peña, C., Quintanal, R.G., Labrador López, M., 1986. Two stage design in a community survey. *Br. J. Psychiatry* 149, 88–97.
- Vasquez-Barquero, J.L., Diez-Manrique, J.F., Peña, C., Aldama, J., Rodríguez, C.S., Arango, J.M., Mirapeix, C., 1987. A community mental health survey in Cantabria: a general description of morbidity. *Psychol. Med.* 17, 227–241.
- Weissman, M.M., Bland, R.C., Canino, G.J., Faravelli, C., Greenwald, S., Hwu, H.-G., Joyce, P.R., Karam, E.G., Lee, C.-K., Lellouch, J., Lepine, J.-P., Newman, S.C., Rubio-Stipec, M., Wells, J.E., Wickramaratne, P.J., Wittchen, H.-U., Yeh, E.-K., 1996. Cross-national epidemiology of major depression and bipolar disorder. *JAMA* 276, 293–299.
- Wertham, F.I., 1929. A group of benign chronic psychoses: prolonged manic excitements. With a statistical study of age, duration and frequency in 2000 manic attacks. *Am. J. Psychiatry* 9, 17–78.
- Wittchen, H.-U., 1998. Reliability and validity studies of the WHO-Composite International Diagnostic Interview (CIDI): a critical review. *J. Psychiatr. Res.* 28 (Suppl. 1), 57–84.
- Wittchen, H.-U., Jacobi, F., 2005. Size and Burden of Mental Disorders in Europe—a critical review and appraisal of 27 studies. *Eur. Neuropsychopharmacol.* 15, 357–376.
- Wittchen, H.-U., Essau, C.A., von Zerssen, D., Krieg, J.C., Zaudig, M., 1992. Lifetime and six-month prevalence of mental disorders in the Munich Follow-up Study. *Eur. Arch. Psychiatry Clin. Neurosci.* 241, 247–258.
- Wittchen, H.-U., Nelson, C.B., Lachner, G., 1998. Prevalence of mental disorders and psychosocial impairments in adolescents and young adults. *Psychol. Med.* 28 (Suppl. 1), 109–126.
- Wittchen, H.-U., Pfister, H., Schmidtke, B., Winter, S., Müller, N., 2000. German National Health Interview and Examination Survey-Mental Health Supplement (GHS-MHS): Part 2. Tables and instructions for public use file (supplement to final report BMBFBW 01 EH 9701/8). Munich: Max-Planck-Institute for Psychiatry, Clinical Psychology and Epidemiology. The public use file can be ordered at: [jacobi@psychologie.tu-dresden.de](mailto:jacobi@psychologie.tu-dresden.de).
- Wittchen, H.-U., Mühlhig, S., Pezawas, L., 2003. Natural course and burden of bipolar disorders. *Int. J. Neuropsychopharmacol.* 6, 145–154.
- Woods, S.W., 2000. The economic burden of bipolar disease. *J. Clin. Psychiatry* 61 (Suppl. 13), 38–41.
- Wunderlich, U., Bronisch, T., Wittchen, H.-U., 1998. Comorbidity patterns in adolescents and young adults with suicide attempts. *Eur. Arch. Psychiatry Clin. Neurosci.* 248 (Suppl. 2), 87–95.